

***H(div) Preconditioning for a Mixed Finite
Element Formulation of the Stochastic Diffusion
Problem***

Furnival, Darran and Elman, Howard and Powell,
Catherine

2008

MIMS EPrint: **2008.66**

Manchester Institute for Mathematical Sciences
School of Mathematics

The University of Manchester

Reports available from: <http://eprints.maths.manchester.ac.uk/>

And by contacting: The MIMS Secretary
School of Mathematics
The University of Manchester
Manchester, M13 9PL, UK

ISSN 1749-9097

$H(\text{div})$ PRECONDITIONING FOR A MIXED FINITE ELEMENT FORMULATION OF THE STOCHASTIC DIFFUSION PROBLEM

HOWARD C. ELMAN, DARRAN G. FURNIVAL, CATHERINE E. POWELL

ABSTRACT. We study $H(\text{div})$ preconditioning for the saddle-point systems that arise in a stochastic Galerkin mixed formulation of the steady-state diffusion problem with random data. The key ingredient is a multigrid V-cycle for a weighted, stochastic $H(\text{div})$ operator, acting on a certain tensor product space of random fields with finite variance. We build on the Arnold-Falk-Winther multigrid algorithm presented in [1] by varying the spatial discretization from grid to grid whilst keeping the stochastic discretization fixed. We extend the deterministic analysis to accommodate the modified $H(\text{div})$ operator and establish spectral equivalence bounds with a new multigrid V-cycle operator that are independent of the spatial and stochastic discretization parameters. We implement multigrid within a block-diagonal preconditioner for the full stochastic saddle-point problem, derive eigenvalue bounds for the preconditioned system matrices and investigate the impact of all the discretization parameters on the convergence rate of preconditioned MINRES.

1. INTRODUCTION

In deterministic modeling of physical processes input variables that represent material properties, boundary conditions, or source terms are assumed to be known explicitly. This results in deterministic partial differential equations (PDEs) whose numerical solution can be effected via traditional discretization methods. In situations where complete knowledge of an input variable is lacking, it is more fitting to pose the problem in a probabilistic setting and solve the resulting stochastic PDEs. The present work is concerned with the numerical solution of the stochastic steady-state diffusion equation written in mixed form. Here, the source of uncertainty is the diffusion coefficient which is modeled as a random field with a prescribed mean and correlation function.

In the last few years there has been a great deal of interest in so-called stochastic Galerkin methods for solving elliptic PDEs with random field coefficients (e.g., see [7], [2], [22]). This body of work has developed rigorous analysis for the pioneering work [11] which advocates coupling a polynomial chaos basis for a subspace of random functions with finite variance, with a traditional finite element spatial discretization. The advantage of this so-called stochastic finite element methodology is that a *single* linear system needs to be solved, the solution to which can be post-processed to obtain probabilistic information of interest. However, this system is typically orders of magnitude larger than those obtained from corresponding deterministic models, and specialised solvers are essential.

If the Galerkin discretization of the stochastic PDE in question is based on a primal weak formulation, the resulting linear system is symmetric and positive definite. An efficient multigrid solver for those systems was proposed in [13] and [8]. If the discretization is based on a mixed formulation, then we have to solve a large-scale symmetric, indefinite system of equations which is more challenging. A mean-based preconditioning scheme was suggested in [9]. However, it loses efficiency if the variance of the random field coefficients is large relative to the mean field.

In [1], Arnold et al. pointed out that the saddle-point matrices that arise from mixed finite element discretizations of the *deterministic* steady-state diffusion equation (with unit coefficients) have the same mapping properties as a block-diagonal matrix whose blocks are discrete representations of the deterministic $H(\text{div})$ and L^2 operators. If exact solves are assumed, this matrix is known to provide a preconditioner for MINRES that yields convergence independent of

This work was supported by the U. S. Department of Energy under grant DEFG0204ER25619, and by the U. S. National Science Foundation undergrant CCF0726017.

the spatial discretisation parameter. Computing the action of the inverse of the leading block is not a trivial task, however, and so a multigrid algorithm for $H(\text{div})$ operators was suggested, leading to a practical preconditioner. Other authors have also suggested deterministic $H(\text{div})$ multigrid algorithms (e.g., see [14], [24], [12].) The work of Arnold et al. [1] is attractive, however, since it takes the form of a standard multigrid V-cycle method with a specialised smoother.

In this work, we propose and analyse a new stochastic $H(\text{div})$ multigrid method. We extend the deterministic analysis from [1] to accommodate a stochastic $H(\text{div})$ operator which is weighted by random diffusion coefficients, and establish spectral equivalence bounds with the proposed multigrid V-cycle operator that are totally independent of the spatial and stochastic discretization parameters. We implement multigrid within a block-diagonal preconditioner for the full stochastic saddle-point system and investigate the impact of all the discretization parameters on the convergence rate of preconditioned MINRES. Numerical results are presented in §8 and we include experiments with a cheaper variant of the analyzed method.

2. STOCHASTIC STEADY STATE DIFFUSION PROBLEM

Here we introduce the stochastic PDE to be solved, derive a weak mixed formulation and an inf-sup stable finite-dimensional problem. As noted in §1, the finite-dimensional problem is obtained by application of a Galerkin finite element method with the solution variables being expressed using curtailed polynomial chaos expansions. A semi-discrete formulation of the problem, that will be used in the subsequent analysis, is also introduced.

2.1. Boundary Value Problem. The stochastic steady-state diffusion equation with homogeneous Dirichlet boundary value conditions is given by

$$(1) \quad \begin{cases} -\nabla \cdot (c\nabla p) = f & \text{in } D \times \Omega, \\ p = 0 & \text{on } \partial D \times \Omega, \end{cases}$$

where D is the spatial domain, Ω is a sample space, $c: D \times \Omega \rightarrow \mathbb{R}$ is the diffusion coefficient, and $f: D \times \Omega \rightarrow \mathbb{R}$ is the source function. The sample space in turn belongs to a probability space (Ω, \mathcal{F}, P) where \mathcal{F} is a σ -algebra and P is a probability measure. Note that the divergence and gradient operators are considered to act on spatial components only. We will consider the spatial domain, D , to be a convex two-dimensional bounded open set with piecewise smooth boundary. In particular we take D to be the interior of a polygon.

Often in solving the diffusion problem the quantity $c\nabla p$ is of more interest than the quantity p . Instead of seeking an approximation to p in (1) and then post-processing this to obtain an approximation to $c\nabla p$, a preferred approach is to obtain an approximation for $c\nabla p$ directly by converting the second order system to a first order system. This is achieved by making the substitution $\vec{u} = c\nabla p$ which gives

$$(2) \quad \begin{cases} c^{-1}\vec{u} - \nabla p = 0 & \text{in } D \times \Omega, \\ -\nabla \cdot \vec{u} = f & \text{in } D \times \Omega, \\ p = 0 & \text{on } \partial D \times \Omega. \end{cases}$$

We will refer to p as the displacement and \vec{u} as the flux.

In order to characterize the random aspect of the problem we start by assuming that c^{-1} is expressed as a curtailed Karhunen-Loève expansion, i.e.

$$(3) \quad c(\vec{x}, \vec{\xi}(\omega))^{-1} = c_0(\vec{x}) + \sum_{r=1}^m \sqrt{\lambda_r} c_r(\vec{x}) \xi_r(\omega)$$

where ξ_r , $r = 1, \dots, m$, are assumed to be continuous, independent, and identically distributed random variables, and (λ_r, c_r) are acquired by solving an eigenvalue problem in the form of a Fredholm integral equation as discussed, e.g., in [11]. Introducing the random vector $\vec{\xi} = (\xi_1, \dots, \xi_m)$ and assuming that $f = f(\vec{x}, \vec{\xi})$, we then define $\mathcal{F} = \sigma(\vec{\xi})$ which is understood to be the minimal σ -algebra generated by $\vec{\xi}$. With the problem defined in this manner, then providing random fields p and \vec{u} can be found that satisfy (2) the Doob-Dynkin lemma, which is given e.g., in [5], tells us that they will be expressible as Borel functions of $\vec{\xi}$ for each $\vec{x} \in D$, i.e. $p = p(\vec{x}, \vec{\xi})$ and $\vec{u} = \vec{u}(\vec{x}, \vec{\xi})$.

Now note, that given $\xi_r(\Omega) = \Gamma$, $r = 1, \dots, m$, and denoting the density function of ξ_r by ρ_r we have, for any Borel function $g = g(\vec{y})$,

$$(4) \quad \int_{\Omega} g(\vec{\xi}(\omega)) dP = \int_{\Gamma^m} g(\vec{y}) \rho(\vec{y}) d\vec{y}$$

where $\rho(\vec{y}) = \rho_1(y_1) \cdots \rho_m(y_m)$ and $d\vec{y} = dy_1 \cdots dy_m$. Here ρ is the joint density function of $\vec{\xi}$ and the above integrals represent the expected value of $g(\vec{\xi})$. As all the random fields in this paper are Borel functions of $\vec{\xi}$ for each $\vec{x} \in D$, (4) can be used to evaluate all integrals over the sample space, Ω , that appear in the sequel.

We will also assume that the reciprocal of the diffusion coefficient (and hence the diffusion coefficient itself) is bounded between two positive constants almost everywhere, i.e.

$$(5) \quad a_{\min} \leq c(\vec{x}, \vec{\xi}(\omega))^{-1} \leq a_{\max} \quad \forall \vec{x} \in D \quad P\text{-a.e.}$$

where by P -a.e. we mean that there exists a set $E \in \mathcal{F}$ with $P(E) = 0$ such that the proposition holds on the complement of E .

2.2. Function Space Notation. We will make use of the Lebesgue spaces L_2 and L_∞ on D and Ω , and the Sobolev spaces H^m , where m is an integer, on D . These spaces and their associated inner products and norms are defined in numerous texts, e.g., [10]. Note that when $m = 0$, $H^m(D) = L_2(D)$.

The spaces $H^m(D)^2$ are understood to contain vector fields whose components are in $H^m(D)$, i.e. given $\vec{v} = (v_1, v_2) \in H^m(D)^2$ then $v_1, v_2 \in H^m(D)$. Inner products on these vector function spaces are defined by summing the inner products on the components, i.e.

$$(\vec{v}, \vec{w})_{H^m(D)^2} = (v_1, w_1)_{H^m(D)} + (v_2, w_2)_{H^m(D)},$$

with the induced norms following. The space $H(\text{div}; D)$ is understood to be the space of functions in $L_2(D)^2$ whose divergence is in $L_2(D)$. It has the inner product

$$(\vec{v}, \vec{w})_{H(\text{div}; D)} = (\vec{v}, \vec{w})_{L_2(D)^2} + (\nabla \cdot \vec{v}, \nabla \cdot \vec{w})_{L_2(D)}$$

from which the norm $\|\cdot\|_{H(\text{div}; D)}$ is induced.

The random fields in this paper will live in tensor products of spaces defined on the spatial domain, D , with spaces defined on the sample space, Ω . The space $H^m(D) \otimes L_2(\Omega)$ is given by

$$H^m(D) \otimes L_2(\Omega) = \{v: D \times \Omega \rightarrow \mathbb{R} \mid D^\gamma v \in L_2(D) \otimes L_2(\Omega), |\gamma| \leq m\},$$

with inner product

$$(v, w)_{H^m(D) \otimes L_2(\Omega)} = \sum_{|\gamma| \leq m} \int_{\Omega} \int_D D^\gamma v D^\gamma w,$$

where $D^\gamma v$ represents weak derivatives of v , i.e.

$$D^\gamma v = \frac{\partial^{|\gamma|} v}{\partial x_1^{\gamma_1} \partial x_2^{\gamma_2}}$$

with $\gamma = (\gamma_1, \gamma_2) \in \mathbb{N}^2$ and $|\gamma| = \gamma_1 + \gamma_2$. This inner product induces the norm $\|\cdot\|_{H^m(D) \otimes L_2(\Omega)}$. The spaces $H^m(D)^2 \otimes L_2(\Omega)$ and $H(\text{div}; D) \otimes L_2(\Omega)$ are defined in an analogous fashion. Tensor products of Hilbert spaces are discussed more formally in [2] and [21].

The space $L_\infty(D) \otimes L_\infty(\Omega)$ is the space of all random fields on $D \times \Omega$ whose absolute value has a finite essential supremum. The norm associated with this space is defined via this essential supremum, i.e.

$$\|v\|_{L_\infty(D) \otimes L_\infty(\Omega)} = \text{ess. sup}_{(\vec{x}, \omega) \in D \times \Omega} |v(\vec{x}, \omega)|.$$

From this definition, we see that $\|c^{-1}\|_{L_\infty(D) \otimes L_\infty(\Omega)} \leq a_{\max}$, where a_{\max} is as in (5).

2.3. Weighted $H(\operatorname{div}; D) \otimes L_2(\Omega)$ Bilinear Form. We define the Hilbert space U to consist of the vector space $L_2(D)^2 \otimes L_2(\Omega)$ paired with the inner product

$$(\vec{v}, \vec{w})_U = \int_{\Omega} \int_D \frac{1}{c} \vec{v} \cdot \vec{w}.$$

This induces the norm $\|\vec{v}\|_U$.

We define a weighted $H(\operatorname{div}; D) \otimes L_2(\Omega)$ bilinear form $\Lambda: H(\operatorname{div}; D) \otimes L_2(\Omega) \times H(\operatorname{div}; D) \otimes L_2(\Omega) \rightarrow \mathbb{R}$ by

$$(6) \quad \Lambda(\vec{v}, \vec{w}) = (\vec{v}, \vec{w})_U + (\nabla \cdot \vec{v}, \nabla \cdot \vec{w})_{L_2(\Omega) \otimes L_2(D)}.$$

This differs from the standard $H(\operatorname{div}; D) \otimes L_2(\Omega)$ bilinear form only by the weight c^{-1} in the first term. As c^{-1} is bounded almost everywhere between two positive constants, as noted in (5), $\Lambda(\cdot, \cdot)$ induces a norm, to be denoted as $\|\cdot\|_{\Lambda}$. Note that this norm is equivalent to $\|\cdot\|_{H(\operatorname{div}; D) \otimes L_2(\Omega)}$.

2.4. Weak Formulation. Let $c \in L_{\infty}(D) \otimes L_{\infty}(\Omega)$ and $f \in L_2(D) \otimes L_2(\Omega)$. The weak formulation of the first-order boundary value problem given in §2.1 is: find $\vec{u} \in H(\operatorname{div}; D) \otimes L_2(\Omega)$ and $p \in L_2(D) \otimes L_2(\Omega)$ such that

$$\begin{aligned} a(\vec{u}, \vec{v}) + b(p, \vec{v}) &= 0 & \forall \vec{v} \in H(\operatorname{div}; D) \otimes L_2(\Omega), \\ b(q, \vec{u}) &= -l(q) & \forall q \in L_2(D) \otimes L_2(\Omega) \end{aligned}$$

where

$$a(\vec{v}, \vec{w}) = (\vec{v}, \vec{w})_U, \quad b(q, \vec{v}) = (q, \nabla \cdot \vec{v})_{L_2(D) \otimes L_2(\Omega)}, \quad l(q) = (f, q)_{L_2(D) \otimes L_2(\Omega)}.$$

This will possess a unique solution, as shown in [4], providing that the bilinear forms $a(\cdot, \cdot)$ and $b(\cdot, \cdot)$ are continuous, $a(\cdot, \cdot)$ is coercive on the null-space of $b(\cdot, \cdot)$, and that there exists a constant $\beta > 0$, called the inf-sup constant, such that

$$(7) \quad \sup_{\substack{\vec{v} \in H(\operatorname{div}; D) \otimes \\ L_2(\Omega) \setminus \{0\}}} \frac{b(q, \vec{v})}{\|\vec{v}\|_{H(\operatorname{div}; D) \otimes L_2(\Omega)}} \geq \beta \|q\|_{L_2(D) \otimes L_2(\Omega)} \quad \forall q \in L_2(D) \otimes L_2(\Omega).$$

The continuity and coercivity conditions on $a(\cdot, \cdot)$ and $b(\cdot, \cdot)$ are readily demonstrated, with the conditions on $a(\cdot, \cdot)$ being satisfied as a consequence of (5). The existence of β was established in [9] where, moreover, it was shown to depend only on the spatial domain D and to be equal to its counterpart in the analogous deterministic analysis.

2.5. Polynomial Chaos. The polynomial chaos expansion method, as pioneered in [11] and generalized in [22], consists of constructing a finite-dimensional subspace of $L_2(\Omega)$, here denoted by T , that is the span of the m -variate polynomials from the Askey scheme of hypergeometric polynomials (discussed in [19]) that are orthogonal with respect to the underlying probability measure. That is, $T = \operatorname{span}\{\chi_1, \dots, \chi_M\}$ where

$$\int_{\Omega} \chi_k(\vec{\xi}) \chi_l(\vec{\xi}) dP = \int_{\Gamma^m} \chi_k(\vec{y}) \chi_l(\vec{y}) \rho(\vec{y}) dy = \kappa_k \delta_{kl}$$

and

$$(8) \quad M = \frac{(m+n)!}{m!n!}.$$

Note that if ρ is the density function corresponding to an m -dimensional Gaussian distribution, then the polynomial chaos will consist of m -variate Hermite polynomials. If ρ is the density function corresponding to an m -dimensional uniform distribution, then the polynomial chaos will consist of m -variate Legendre polynomials. For polynomials corresponding to other distributions see [23].

2.6. Finite Element Formulation. Let $\mathcal{T} = \{\Delta_1, \dots, \Delta_K\}$ be a triangulation on D . We assume that the triangulation belongs to a family of triangulations that are quasi-uniform and shape regular. Let h_k be the length of the longest side of Δ_k and define $h = \max h_k$ to be the mesh parameter of \mathcal{T} .

In the following we will use tensor products of finite dimensional vector spaces. Note that such spaces are spanned by products of basis functions from the vector spaces that comprise the tensor product. The zeroth-order Raviart-Thomas space is defined by

$$R = \{ \vec{v} \in H(\text{div}; D) \mid v|_{\Delta_k} \in P_0(\Delta_k)^2 + P_0(\Delta_k)\vec{x} \}$$

where $P_0(\Delta_k)$ denotes polynomials of zero degree, i.e. constants on Δ_k . Let $\{\vec{\phi}_1, \dots, \vec{\phi}_{N_R}\}$ be a basis of R and let $Q = \text{span}\{\psi_1, \dots, \psi_{N_Q}\}$ be the set of piecewise constants on the triangulation \mathcal{T} . Then $R \otimes T \subset H(\text{div}; D) \otimes L_2(\Omega)$ and $Q \otimes T \subset L_2(D) \otimes L_2(\Omega)$. Note that R and Q are known to satisfy a discrete deterministic inf-sup condition, as shown, for example, in [4] and [18].

The finite element formulation of the weak formulation given in §2.4 is given by: find $\vec{u}_{hn} \in R \otimes T$ and $p_{hn} \in Q \otimes T$ such that

$$\begin{aligned} a(\vec{u}_{hn}, \vec{v}) + b(p_{hn}, \vec{v}) &= 0 & \forall \vec{v} \in R \otimes T, \\ b(q, \vec{u}_{hn}) &= -l(q), & \forall q \in Q \otimes T. \end{aligned}$$

This possesses a unique solution under the same conditions that apply for the weak formulation. The discrete continuity and coercivity conditions on $a(\cdot, \cdot)$ and $b(\cdot, \cdot)$ follow in an analogous way to the infinite-dimensional case considered in §2.4. Existence and uniqueness is then assured if it can be shown that there exists a constant $\beta_{hn} > 0$ such that

$$(9) \quad \sup_{\vec{v} \in R \otimes T \setminus \{0\}} \frac{b(q, \vec{v})}{\|\vec{v}\|_{H(\text{div}; D) \otimes L_2(\Omega)}} \geq \beta_{hn} \|q\|_{L_2(D) \otimes L_2(\Omega)} \quad \forall q \in Q \otimes T.$$

The existence of β_{hn} can be demonstrated by employing Fortin's Lemma (see [3]) as follows. First we note that by considering $\nabla \cdot \nabla s = w$ where $w \in L_2(D) \otimes T$, it can be shown that there exists a unique $\vec{z} \in H^1(D)^2 \otimes T$ such that

$$\|\vec{z}\|_{H(\text{div}; D) \otimes L_2(\Omega)} \leq C_D \|w\|_{L_2(D) \otimes L_2(\Omega)}$$

where C_D is only dependent on the spatial domain. Using standard arguments it then follows that the constant $\beta > 0$ satisfying (7) also satisfies a semi-discrete inf-sup condition given by

$$(10) \quad \sup_{\vec{v} \in H^1(D)^2 \otimes T \setminus \{0\}} \frac{b(q, \vec{v})}{\|\vec{v}\|_{H(\text{div}; D) \otimes L_2(\Omega)}} \geq \beta \|q\|_{L_2(D) \otimes L_2(\Omega)} \quad \forall q \in L_2(D) \otimes T.$$

Now suppose there exists an operator $\Pi: H^1(D)^2 \otimes T \rightarrow R \otimes T$ such that for each $\vec{v} \in H^1(D) \otimes T$,

$$\begin{aligned} b(\Pi\vec{v} - \vec{v}, q) &= 0 \quad \forall q \in L_2(D) \otimes L_2(\Omega), \\ \|\Pi\vec{v}\|_{H(\text{div}; D) \otimes L_2(\Omega)} &\leq C_\Pi \|\vec{v}\|_{H(\text{div}; D) \otimes L_2(\Omega)}, \end{aligned}$$

where C_Π is independent of h , m , and n . Then

$$\begin{aligned} \sup_{\vec{v} \in R \otimes T \setminus \{0\}} \frac{b(q, \vec{v})}{\|\vec{v}\|_{H(\text{div}; D) \otimes L_2(\Omega)}} &= \sup_{\vec{v} \in H^1(D)^2 \otimes T \setminus \{0\}} \frac{b(q, \Pi\vec{v})}{\|\Pi\vec{v}\|_{H(\text{div}; D) \otimes L_2(\Omega)}} \\ &= \sup_{\vec{v} \in H^1(D)^2 \otimes T \setminus \{0\}} \frac{b(q, \Pi\vec{v} - \vec{v}) + b(\vec{v}, q)}{\|\Pi\vec{v}\|_{H(\text{div}; D) \otimes L_2(\Omega)}} \\ &\geq \frac{\beta}{C_\Pi} \|q\|_{L_2(D) \otimes L_2(\Omega)} \quad \forall q \in Q \otimes T. \end{aligned}$$

An operator Π satisfying the above properties is given by the Raviart-Thomas interpolation operator described in §2.8. Therefore, the existence of the discrete inf-sup constant is established and, moreover, it is independent of the discretization parameters h , m , and n .

In [6] the well-posedness of a stochastic saddle-point problem arising in the solution of a PDE with stochastic boundary was analysed. In that work, an inf-sup stability result was established under the assumption that a particular discrete stochastic space is chosen. Here, we note that (9) holds independently of the choice of $T \subset L_2(\Omega)$.

2.7. Matrix Formulation. The finite element formulation given in §2.6 can be written as a matrix problem. This is obtained by using the expansions

$$(11) \quad \vec{u}_{hn} = \sum_{j=1}^{N_R} \sum_{l=1}^M u_{jl} \vec{\phi}_j \chi_l \quad p_{hn} = \sum_{j=1}^{N_Q} \sum_{l=1}^M p_{jl} \psi_j \chi_l$$

and allowing \vec{v} to vary over the basis functions of $R \otimes T$ and q to vary over the basis functions of $Q \otimes T$. This leads to a matrix problem of the form

$$(12) \quad \begin{bmatrix} A & B^T \\ B & 0 \end{bmatrix} \begin{bmatrix} \mathbf{u} \\ \mathbf{p} \end{bmatrix} = \begin{bmatrix} \mathbf{0} \\ \mathbf{f} \end{bmatrix}.$$

The matrices that appear in the system matrix can be decomposed as

$$A = G_0 \otimes K_0 + \sum_{r=1}^m \sqrt{\lambda_r} G_r \otimes K_r, \quad B = G_0 \otimes B_0,$$

with entries given by

$$\begin{aligned} [G_r]_{kl} &= \int_{\Omega} \xi_r \chi_k \chi_l \quad \left(= \int_{\Gamma^m} y_r \chi_k \chi_l \rho \right), \\ [K_r]_{ij} &= \int_D c_r \vec{\phi}_i \cdot \vec{\phi}_j, \quad [B_0]_{ij} = \int_D \psi_i \nabla \cdot \vec{\phi}_j, \end{aligned}$$

where $\xi_0 = 1$ in the definition of G_0 . The vector \mathbf{f} has entries given by

$$\mathbf{f} = \begin{bmatrix} \mathbf{f}_1 \\ \vdots \\ \mathbf{f}_M \end{bmatrix}, \quad [\mathbf{f}_k]_i = - \int_{\Omega} \int_D f \psi_i \chi_k.$$

The solution vector contains the coefficients of \vec{u}_{hn} and p_{hn} stacked columnwise. Once it has been computed we have approximations to the random fields \vec{u} and p that satisfy the weak formulation of the diffusion problem. These approximations can then be post-processed to obtain probabilistic information about the solution variables. The means and variances can be trivially computed, and the probability distributions of quantities of interest can be computed via sampling of $\vec{\xi}$.

2.8. Interpolation Operator. We here define a spatial interpolation operator $\Pi: H^1(D)^2 \otimes L_2(\Omega) \rightarrow R \otimes L_2(\Omega)$ such that for each $\omega \in \Omega$, Π maps $v(\vec{x}, \vec{\xi})$ to a function in R in the conventional manner (see e.g., [3, p. 147]), i.e. Π acts only on the spatial components of the random fields in its domain and does so in a manner analogous to its deterministic counterpart. To be more precise, we first define $\Pi_k: H^1(\Delta_k)^2 \otimes L_2(\Omega) \rightarrow R \otimes L_2(\Omega)$ such that for $\vec{v} \in H^1(\Delta_k)^2 \otimes L_2(\Omega)$,

$$\int_{e_i} (\vec{v} - \Pi_k \vec{v}) \cdot \vec{n}_i = 0, \quad i = 1, 2, 3, \quad \forall \omega \in \Omega,$$

where e_i , $i = 1, 2, 3$, are the edges of Δ_k , and \vec{n}_i , $i = 1, 2, 3$ are the respective unit normal vectors to these edges. Then Π is defined such that for $\vec{v} \in H^1(D)^2 \otimes L_2(\Omega)$,

$$(\Pi \vec{v})|_{\Delta_k} = \Pi_k(\vec{v}|_{\Delta_k}).$$

Following [1], we have the approximation property

$$\|\vec{v} - \Pi \vec{v}\|_{L_2(D)^2} \leq C_0 h \|\vec{v}\|_{H^1(D)^2} \quad \forall \omega \in \Omega,$$

where C_0 is a constant. Squaring and integrating over Ω gives

$$(13) \quad \|\vec{v} - \Pi \vec{v}\|_{L_2(D)^2 \otimes L_2(\Omega)} \leq C_0 h \|\vec{v}\|_{H^1(D)^2 \otimes L_2(\Omega)}.$$

2.9. Semi-Discrete Formulation. We here introduce a semi-discrete formulation of the weak formulation of the diffusion problem where only the stochastic space $T \subset L_2(\Omega)$ is finite-dimensional. This is given by: find $\vec{u}_n \in H(\text{div}; D) \otimes T$ and $p_n \in L_2(D) \otimes T$ such that

$$\begin{aligned} a(\vec{u}_n, \vec{v}) + b(p_n, \nabla \cdot \vec{v}) &= 0 & \forall v \in H(\text{div}; D) \otimes T, \\ b(q, \nabla \cdot \vec{u}_n) &= -l(q) & \forall q \in L_2(D) \otimes T. \end{aligned}$$

This possesses a unique solution under an analogous set of conditions as for the weak formulation and the finite element formulation discussed in §2.4 and §2.6 respectively. In particular, the inf-sup condition that is required, over the relevant semi-discrete space, was established in §2.6 and is given by (10).

In order to obtain a bound on the error between the flux solutions to the semi-discrete and the fully discrete problems we make the following assumption. Given $f \in L_2(D) \otimes T$ we assume that $\vec{u}_n \in H^1(D)^2 \otimes T$ and that for each realization of ξ ,

$$\|\vec{u}_n\|_{H^1(D)^2} \leq C_r \|f\|_{L_2(D)},$$

from which it follows that

$$(14) \quad \|\vec{u}_n\|_{H^1(D)^2 \otimes L_2(\Omega)} \leq C_r \|f\|_{L_2(D) \otimes L_2(\Omega)}.$$

Theorem 1. *Given $f \in L_2(D) \otimes T$, then the flux solution to the semi-discrete problem, \vec{u}_n , and the flux solution to the fully discrete problem, \vec{u}_{hn} , satisfy*

$$\|\vec{u}_n - \vec{u}_{hn}\|_U \leq C_* h \|f\|_{L_2(D) \otimes L_2(\Omega)},$$

where C_* is independent of h , m , and n , and a_{\max} is as in (5).

Proof. Using the upper bound on c^{-1} in (5) and following [4] we have

$$\begin{aligned} \|\vec{u}_n - \vec{u}_{hn}\|_U &\leq a_{\max}^{1/2} \|\vec{u}_n - \vec{u}_{hn}\|_{L_2(D) \otimes L_2(\Omega)} \\ &\leq a_{\max}^{1/2} C_* \inf_{\vec{v} \in R \otimes T} \|\vec{u}_n - \vec{v}\|_{L_2(D) \otimes L_2(\Omega)} \end{aligned}$$

where C_* is a constant. Now, as $\Pi \vec{u}_n \in R \otimes T$, using (13) and (14) we have

$$\begin{aligned} \|\vec{u}_n - \vec{u}_{hn}\|_U &\leq a_{\max}^{1/2} C_* C_0 h \|\vec{u}_n\|_{H^1(D)^2 \otimes L_2(\Omega)} \\ &\leq a_{\max}^{1/2} C_* C_0 C_r h \|f\|_{L_2(D) \otimes L_2(\Omega)} \end{aligned}$$

which demonstrates the theorem with $C_* = a_{\max}^{1/2} C_* C_0 C_r$. \square

3. HELMHOLTZ DECOMPOSITION, PROJECTION AND $H(\text{div}; D)$ OPERATORS

In this section we describe a Helmholtz decomposition of $\vec{v} \in R \otimes T$ analogous to its deterministic counterpart given in [1]. We also introduce some projection operators and a weighted $H(\text{div}; D) \otimes L_2(\Omega)$ operator which, along with the Helmholtz decomposition, will be used in §§5, 6 and 7.

3.1. Helmholtz Decomposition. Let $W = \text{span}\{\varphi_1, \dots, \varphi_{N_W}\}$ be the set of piecewise linear functions defined on the triangulation \mathcal{T} . Here N_W will equal the number of nodes in the triangulation. It is well known, e.g, see [1], that

$$\{\vec{v} \in R \mid \nabla \cdot \vec{v} = 0\} = \{\nabla \times w \mid w \in W\}.$$

As divergence and curl are purely spatial operators, i.e. they don't affect the stochastic nature of the random fields they act upon, it follows that

$$(15) \quad \{\vec{v} \in R \otimes T \mid \nabla \cdot \vec{v} = 0\} = \{\nabla \times w \mid w \in W \otimes T\}.$$

Here $\nabla \times w = (-\partial w / \partial x_2, \partial w / \partial x_1)$.

We now define a weighted discrete gradient operator (cf. [1, eq. (3.1)]), denoted $\text{grad}_h^c : Q \otimes T \rightarrow R \otimes T$, such that for $q \in Q \otimes T$,

$$(16) \quad (\text{grad}_h^c q, \vec{v})_U = -(q, \nabla \cdot \vec{v})_{L_2(D) \otimes L_2(\Omega)} \quad \forall \vec{v} \in R \otimes T.$$

The superscript c denotes the dependence of the operator on the random field c through the definition of $(\cdot, \cdot)_U$.

Theorem 2. (*Helmholtz Decomposition*) Given $\vec{v} \in R \otimes T$, there exists $q \in Q \otimes T$ and $w \in W \otimes T$ such that

$$(17) \quad \vec{v} = \text{grad}_h^c q + \nabla \times w.$$

Proof. To establish the given decomposition it is sufficient to show that $\text{grad}_h^c(Q \otimes T)$ and $\nabla \cdot (W \otimes T)$ are orthogonal complements on $R \otimes T$ with respect to $(\cdot, \cdot)_U$. To see this, let $w \in W \otimes T$, then

$$\begin{aligned} \nabla \cdot (\nabla \times w) = 0 &\Leftrightarrow (\nabla \cdot (\nabla \times w), q)_{L_2(D) \otimes L_2(\Omega)} = 0 && \forall q \in Q \otimes T \\ &\Leftrightarrow (\nabla \times w, \text{grad}_h^c q)_U = 0 && \forall q \in Q \otimes T \\ &\Leftrightarrow \nabla \times w \in (\text{grad}_h^c(Q \otimes T))^\perp. \end{aligned}$$

□

Note that, for $q \in Q \otimes T$ and $w \in W \otimes T$, $\text{grad}_h^c q$ and $\nabla \times w$ are also orthogonal with respect to $(\cdot, \cdot)_\Lambda$, where $(\cdot, \cdot)_\Lambda$ is as defined in §2.3.

3.2. Projection Operators. We define the projection operator $\Theta: L_2(D) \otimes L_2(\Omega) \rightarrow Q \otimes T$ such that for each $v \in L_2(D) \otimes L_2(\Omega)$,

$$(\Theta v, q)_{L_2(D) \otimes L_2(\Omega)} = (v, q)_{L_2(D) \otimes L_2(\Omega)} \quad \forall q \in Q \otimes T.$$

The operator Θ affects both the spatial and the stochastic aspect of the random fields on which it operates. It will be in our interest to define an operator that only affects the spatial aspect of the random fields it acts upon. Therefore, we define $\Sigma: L_2(D) \otimes L_2(\Omega) \rightarrow Q \otimes L_2(\Omega)$ such that for each $v \in L_2(D) \otimes L_2(\Omega)$,

$$(\Sigma v, q)_{L_2(D)} = (v, q)_{L_2(D)} \quad \forall q \in Q \quad \forall \omega \in \Omega,$$

that is, for each $\omega \in \Omega$, Σ is the $L_2(D)$ -projection of $v(\vec{x}, \vec{\xi})$ onto Q . It can then be shown that given $\vec{v} \in H^1(D)^2 \otimes L_2(\Omega)$ we have the commutativity property

$$\nabla \cdot \Pi \vec{v} = \Sigma \nabla \cdot \vec{v}, \quad \forall \omega \in \Omega,$$

the proof of which follows that given in [3] for the deterministic case. Using this we also have for $\vec{v} \in R \otimes T$ and $q \in Q \otimes T$,

$$(18) \quad \begin{aligned} (\Theta \nabla \cdot \vec{v}, q)_{L_2(D) \otimes L_2(\Omega)} &= \int_\Omega (\nabla \cdot \vec{v}, q)_{L_2(D)} = \int_\Omega (\Sigma \nabla \cdot \vec{v}, q)_{L_2(D)} \\ &= \int_\Omega (\nabla \cdot \Pi \vec{v}, q)_{L_2(D)} = (\nabla \cdot \Pi \vec{v}, q)_{L_2(D) \otimes L_2(\Omega)}, \end{aligned}$$

which is analogous to the deterministic result given in [1].

We define $\mathcal{P}: H(\text{div}; D) \otimes L_2(\Omega) \rightarrow R \otimes T$ such that for each $\vec{w} \in H(\text{div}; D) \otimes L_2(\Omega)$,

$$\Lambda(\mathcal{P}\vec{w}, \vec{v}) = \Lambda(\vec{w}, \vec{v}) \quad \forall \vec{v} \in R \otimes T.$$

That is, \mathcal{P} is the projection operator onto $R \otimes T$ defined with respect to $\Lambda(\cdot, \cdot)$.

3.3. Weighted $H(\text{div}; D) \otimes L_2(\Omega)$ Operator. We next define the weighted $H(\text{div}; D) \otimes L_2(\Omega)$ operator $\mathcal{H}: R \otimes T \rightarrow R \otimes T$ such that for each $\vec{w} \in R \otimes T$,

$$(\mathcal{H}\vec{w}, \vec{v})_U = \Lambda(\vec{w}, \vec{v}) \quad \forall \vec{v} \in R \otimes T.$$

Note that \mathcal{H} is a positive-definite symmetric operator, therefore we can define its inverse $\mathcal{H}^{-1}: R \otimes T \rightarrow R \otimes T$ such that for each $\vec{w} \in R \otimes T$,

$$\Lambda(\mathcal{H}^{-1}\vec{w}, v) = (\vec{w}, \vec{v})_U \quad \forall \vec{v} \in R \otimes T.$$

Note also that \mathcal{H} maps $\text{grad}_h^c(Q \otimes T)$ onto itself. To see this, let $\vec{w} = \text{grad}_h^c q$ for some $q \in Q \otimes T$. Then, using (6) and (16),

$$\begin{aligned} (\mathcal{H}\vec{w}, \vec{v})_U &= (\text{grad}_h^c q, \vec{v})_U + (\nabla \cdot \text{grad}_h^c q, \nabla \cdot \vec{v})_{L_2(D) \otimes L_2(\Omega)} \\ &= (\text{grad}_h^c q, \vec{v})_U - (\text{grad}_h^c \nabla \cdot \text{grad}_h^c q, \vec{v})_U \\ &= (\text{grad}_h^c (q - \nabla \cdot \text{grad}_h^c q), \vec{v})_U \quad \forall \vec{v} \in R \otimes T. \end{aligned}$$

As $q - \nabla \cdot \text{grad}_h^c q \in Q \otimes T$ it follows that given $q \in Q \otimes T$ there exists $q' \in Q \otimes T$ such that $\mathcal{H} \text{grad}_h^c q = \text{grad}_h^c q'$. Similarly, for $q \in Q \otimes T$, it can be shown that there exists $q' \in Q \otimes T$ such that $\mathcal{H}^{-1} \text{grad}_h^c q = \text{grad}_h^c q'$.

4. SOLVING THE FIRST ORDER DIFFUSION PROBLEM

The coefficient matrix given in (12), denoted here by C , is sparse, symmetric, and indefinite. Hence, a suitable solution scheme is the MINRES algorithm, described e.g., in [15]. This is a Krylov subspace method that minimizes the Euclidean norm of the residual error at each step. In order to ensure that the iteration is efficient, preconditioning is essential. Ideally, we want the number of MINRES iterations required to reduce the error to within a specified tolerance to be independent of all the discretization parameters, and so we seek a preconditioner P such that the eigenvalues of $P^{-1}C$ are independent of the discretization parameters h , n , and m .

4.1. Deterministic $H(\text{div}; D)$ preconditioning. Recall that in the deterministic diffusion problem, where the diffusion coefficient, $c = c_0(\vec{x})$, and the source function, $f = f_0(\vec{x})$, are known (deterministic) fields, the standard mixed Galerkin discretisation gives rise to a linear system of the form

$$(19) \quad \begin{bmatrix} K_0 & B_0^T \\ B_0 & 0 \end{bmatrix} \begin{bmatrix} \mathbf{u}_0 \\ \mathbf{p}_0 \end{bmatrix} = \begin{bmatrix} \mathbf{0} \\ \mathbf{f}_0 \end{bmatrix},$$

where $B_0 \in \mathbb{R}^{N_Q \times N_R}$ and $K_0 \in \mathbb{R}^{N_R \times N_R}$ are defined as in §2.7. The so-called ideal $H(\text{div}; D)$ preconditioner given by

$$(20) \quad P_0 = \begin{bmatrix} K_0 + D_0 & 0 \\ 0 & N_0 \end{bmatrix},$$

where $D_0 \in \mathbb{R}^{N_R \times N_R}$ and $N_0 \in \mathbb{R}^{N_Q \times N_Q}$ are defined via

$$[D_0]_{ij} = \int_D (\nabla \cdot \vec{\phi}_i)(\nabla \cdot \vec{\phi}_j), \quad [N_0]_{ij} = \int_D \psi_i \psi_j,$$

was proposed in [1] for the case $c_0(\vec{x}) = 1$ and subsequently analysed in [17] and [16] for arbitrary diffusion coefficients. This choice of preconditioner is motivated by the observation that $\forall \vec{v} \in R$ and $\forall q \in Q$,

$$\|c^{-\frac{1}{2}} \vec{v}\|_{L^2(D)}^2 + \|\nabla \cdot \vec{v}\|_{L^2(D)}^2 = (\mathbf{v}, (K_0 + D_0) \mathbf{v}), \quad \|q\|_{L^2(D)}^2 = (\mathbf{q}, N_0 \mathbf{q}).$$

Consequently, the diagonal blocks of P_0 provide representations of a pair of norms in which the chosen mixed finite element discretization is known to be inf-sup stable (see [4, 18]). Hence, P_0 mimics the mapping properties of the underlying saddle-point matrix. The following theorem gives bounds for the eigenvalues of $P_0^{-1}C_0$, where C_0 is the system matrix in (19).

Theorem 3. *The $N_R + N_Q$ eigenvalues of*

$$\begin{bmatrix} K_0 & B_0^T \\ B_0 & 0 \end{bmatrix} \begin{bmatrix} \mathbf{v} \\ \mathbf{q} \end{bmatrix} = \nu \begin{bmatrix} K_0 + D_0 & 0 \\ 0 & N_0 \end{bmatrix} \begin{bmatrix} \mathbf{v} \\ \mathbf{q} \end{bmatrix},$$

lie in $(-1, -\mu_0] \cup \{1\}$, where $\mu_0 \in (0, 1)$ is a constant independent of h .

Proof. See [17]. □

To obtain a practical scheme we need to approximate the action of the inverse of $K_0 + D_0$. Suppose then that there exists some (any) matrix V_0 satisfying,

$$(21) \quad \theta_0 \leq \frac{(\mathbf{v}, (K_0 + D_0) \mathbf{v})}{(\mathbf{v}, V_0 \mathbf{v})} \leq \bar{\theta}_0 \leq 1, \quad \forall \mathbf{v} \in \mathbb{R}^{N_R \times N_R} \setminus \{\mathbf{0}\},$$

where (\cdot, \cdot) indicates the Euclidean inner product. The following theorem gives bounds for the eigenvalues of the preconditioned system obtained by replacing $K_0 + D_0$ in (20) with V_0 .

Theorem 4. *The $(N_R + N_Q)$ eigenvalues of*

$$\begin{bmatrix} K_0 & B_0^T \\ B_0 & 0 \end{bmatrix} \begin{bmatrix} \mathbf{v} \\ \mathbf{q} \end{bmatrix} = \nu \begin{bmatrix} V_0 & 0 \\ 0 & N_0 \end{bmatrix} \begin{bmatrix} \mathbf{v} \\ \mathbf{q} \end{bmatrix},$$

lie in

$$\left(-1, \frac{1}{2} \left(\theta_0 (1 - \mu_0) - \sqrt{\theta_0^2 (\mu_0 - 1)^2 + 4\mu_0 \theta_0} \right) \right) \cup [\theta_0, 1],$$

where μ_0 is the quantity defined in Theorem 3 and θ_0 is the constant from (21).

Proof. See [16]. □

In [1], Arnold et al. demonstrated that, in the case $c_0(\vec{x}) = 1$, choosing V_0 to be the matrix that represents the inverse of a certain multigrid V-cycle operator yields $\theta_0 = 1 - \delta_0$ and $\bar{\theta}_0 = 1$ in (21), where δ_0 is a constant independent of h . Theorem 4 then tells us that the preconditioner

$$(22) \quad \begin{bmatrix} V_0 & 0 \\ 0 & N_0 \end{bmatrix}$$

is h -optimal. For a general diffusion coefficient both μ_0 and δ_0 depend on $c_0(\vec{x})$ and in [16] it was shown that (22) is not always c -optimal.

4.2. Stochastic $H(\text{div}; D) \otimes L_2(\Omega)$ preconditioning. We base our preconditioning strategy for (12) on the deterministic preconditioner described in §4.1. That is, we define, first, the analogous ideal preconditioner

$$(23) \quad P = \begin{bmatrix} A + F & 0 \\ 0 & N \end{bmatrix}$$

where A is the $(1, 1)$ block of (12), $F = G_0 \otimes D_0$ and $N = G_0 \otimes N_0$. Now, $\forall \vec{v} \in R \otimes T$ and $\forall \mathbf{q} \in Q \otimes T$, the blocks of the preconditioner provide representations of the tensor product norms,

$$\|\vec{v}\|_\Lambda^2 = (\mathbf{v}, (A + F) \mathbf{v}), \quad \|q\|_{L^2(D) \otimes L^2(\Omega)}^2 = (\mathbf{q}, N \mathbf{q}).$$

The efficiency of this preconditioner can be analysed in exactly the same way as in the deterministic case. Recalling that M is the dimension of the stochastic space T defined in §2.5, and N_R and N_Q are the dimensions of the deterministic spaces R and Q defined in §2.6, we have the following theorem.

Theorem 5. *The $M \times (N_R + N_Q)$ eigenvalues of the generalised eigenvalue problem,*

$$\begin{bmatrix} A & B^T \\ B & 0 \end{bmatrix} \begin{bmatrix} \mathbf{v} \\ \mathbf{q} \end{bmatrix} = \nu \begin{bmatrix} A + F & 0 \\ 0 & N \end{bmatrix} \begin{bmatrix} \mathbf{v} \\ \mathbf{q} \end{bmatrix}$$

lie in $(-1, -\mu] \cup \{1\}$, where $\mu \in (0, 1)$ is a constant independent of h , m , and n .

Proof. The proof follows that for the deterministic case and will be given here in outline. First note that since $\nabla \cdot R = Q$, we have $D_0 = B_0^T N_0^{-1} B_0$ and

$$\begin{aligned} B^T N^{-1} B &= (G_0 \otimes B_0)^T (G_0 \otimes N_0)^{-1} (G_0 \otimes B_0) \\ &= G_0 \otimes (B_0^T N_0^{-1} B_0) = G_0 \otimes D_0 = F. \end{aligned}$$

It immediately follows that there are MN_R eigenvalues equal to one. The remaining eigenvalues are negative and satisfy

$$B(A + F)^{-1} B^T \mathbf{q} = -\nu N \mathbf{q}.$$

Now, using (9) along with the equivalence of $\|\cdot\|_{H(\text{div}; D) \otimes L_2(\Omega)}$ and $\|\cdot\|_\Lambda$ gives

$$\beta_{hn} \|q\|_{L_2(D) \otimes L_2(\Omega)} \leq \frac{1}{\min\{1, a_{\max}^{-1/2}\}} \sup_{\vec{v} \in R \otimes T \setminus \{0\}} \frac{b(q, \nabla \cdot \vec{v})}{\|\vec{v}\|_\Lambda} \quad \forall q \in Q \otimes T.$$

Converting this to matrix notation one then obtains

$$\beta_{hn}^2 \min\{1, a_{\max}^{-1}\} \leq \frac{(\mathbf{q}, B(A + F)^{-1} B^T \mathbf{q})}{(\mathbf{q}, N \mathbf{q})} \quad \forall \mathbf{q} \in \mathbb{R}^{M \times N_Q}.$$

Therefore, we can choose $\mu = \beta_{hn}^2 \min\{1, a_{\max}^{-1}\}$. \square

To obtain a practical preconditioning scheme, we need a computationally optimal way to approximate the inverse of $A + F$ in each MINRES iteration. If we have at our disposal a matrix V that satisfies

$$(24) \quad \theta \leq \frac{(\mathbf{v}, (A + F)\mathbf{v})}{(\mathbf{v}, V\mathbf{v})} \leq \bar{\theta} \leq 1, \quad \forall \mathbf{v} \in \mathbb{R}^{MN_R} \setminus \{\mathbf{0}\},$$

then the analogue of Theorem 4 holds. Specifically, we have the following result.

Theorem 6. *The $M \times (N_R + N_Q)$ eigenvalues of*

$$\begin{bmatrix} A & B^T \\ B & 0 \end{bmatrix} \begin{bmatrix} \mathbf{v} \\ \mathbf{q} \end{bmatrix} = \nu \begin{bmatrix} V & 0 \\ 0 & N \end{bmatrix} \begin{bmatrix} \mathbf{v} \\ \mathbf{q} \end{bmatrix},$$

lie in

$$\left(-1, \frac{1}{2} \left(\theta(1 - \mu) - \sqrt{\theta^2(\mu - 1)^2 + 4\mu\theta} \right) \right] \cup [\theta, 1],$$

where μ is as in Theorem 5 and θ is the constant from (24).

Proof. Follow the proof of Theorem 4.1 in [16] replacing deterministic matrices with their larger Kronecker product counterparts. \square

In the next sections, we will extend the deterministic analysis presented in [1] to construct a hybrid multigrid V-cycle operator, $\mathcal{V} : R \otimes T \rightarrow R \otimes T$ whose matrix representation V provides spectral equivalence bounds (24) for $A + F$ with constant $\bar{\theta} = 1$ and $\theta = 1 - \delta$ where δ is a constant independent of h , m , and n . The efficiency of the resulting preconditioner,

$$(25) \quad \begin{bmatrix} V & 0 \\ 0 & N \end{bmatrix},$$

is then determined by the spectral inclusion bounds in Theorem 6.

5. TWO-GRID RESULTS

In the following, two triangulations on the spatial domain, \mathcal{T}_h and \mathcal{T}_H , are used with mesh parameters h and H respectively, with $H > h$. Throughout the remainder of this section the spaces and operators defined in §2 will carry a subscript h or H to indicate the underlying triangulation they are defined with respect to.

5.1. Two-grid Bounds. Here we obtain some bounds between functions defined on the coarse grid and related functions defined on the fine grid. The bounds are analogous to those given in Lemma 3.1 in [1].

We first note that given an arbitrary function $q_h \in Q_h \otimes T$ and defining $\vec{v}_h = \text{grad}_h^c q_h \in R_h \otimes T$, we can uniquely define q_H and \vec{v}_H such that $\vec{v}_H = \text{grad}_H^c q_H$ and $\nabla \cdot \vec{v}_H = \Theta_H \nabla \cdot \vec{v}_h$. Moreover, given $f = -\nabla \cdot \text{grad}_h^c q_h$, (q_h, \vec{v}_h) and (q_H, \vec{v}_H) will be finite element approximations to the diffusion problem given by (2), as can be demonstrated using the finite element formulation of the diffusion problem given in §2.6.

Lemma 1. *Given $q_h \in Q_h \otimes T$ there exists $\vec{\sigma} \in R_h \otimes T$ such that*

$$\nabla \cdot \vec{\sigma} = q_h - \Theta_H q_h, \quad \|\vec{\sigma}\|_{H^1(D)^2 \otimes L_2(\Omega)} \leq C_1 \|q_h - \Theta_H q_h\|_{L_2(D) \otimes L_2(\Omega)},$$

where C_1 is independent of h , m , and n .

Proof. Consider the semi-discrete variational problem: find $p \in H_0^1(D) \otimes T$ such that

$$\int_{\Omega} \int_D \nabla p \cdot \nabla v = \int_{\Omega} \int_D (q_h - \Theta_H q_h) v \quad \forall v \in H_0^1(D) \otimes T.$$

This can be seen as the weak formulation of the second-order diffusion problem with source function $f = q_h - \Theta_H q_h$. From regularity considerations (cf. [2]) we have

$$\|p\|_{H^2(D)} \leq C_D \|q_h - \Theta_H q_h\|_{L_2(D)} \quad \forall \omega \in \Omega$$

where C_D is only dependent on the boundary of the spatial domain. From this it follows that

$$\|p\|_{H^2(D)\otimes L_2(\Omega)} \leq C_D \|q_h - \Theta_H q_h\|_{L_2(D)\otimes L_2(\Omega)}.$$

Since p solves the diffusion problem we have

$$-\nabla^2 p = q_h - \Theta_H q_h \quad \text{a.e.}$$

Now set $\vec{\sigma} = -\nabla p \in R_h \otimes T$. Then

$$\|\vec{\sigma}\|_{H^1(D)^2 \otimes L_2(\Omega)} = \|\nabla p\|_{H^1(D)^2 \otimes L_2(\Omega)} \leq \|p\|_{H^2(D)\otimes L_2(\Omega)} \leq C_D \|q_h - \Theta_H q_h\|_{L_2(D)\otimes L_2(\Omega)}$$

which, equating C_D with C_1 , establishes the lemma. \square

Theorem 7. For $q_h \in Q_h \otimes T$,

$$\|q_h - \Theta_H q_h\|_{L_2(D)\otimes L_2(\Omega)} \leq C_2 H \|\text{grad}_h^c q_h\|_U$$

where C_2 is independent of h , m , and n .

Proof. Define $\nabla \cdot \vec{\sigma} = q_h - \Theta_H q_h$ as in Lemma 1. Then

$$\begin{aligned} \|q_h - \Theta_H q_h\|_{L_2(D)\otimes L_2(\Omega)}^2 &= (\nabla \cdot \vec{\sigma}, q_h - \Theta_H q_h)_{L_2(D)\otimes L_2(\Omega)} = (\nabla \cdot \vec{\sigma}, (\Theta_h - \Theta_H)q_h)_{L_2(D)\otimes L_2(\Omega)} \\ &= ((\Theta_h - \Theta_H)\nabla \cdot \vec{\sigma}, q_h)_{L_2(D)\otimes L_2(\Omega)} = (\nabla \cdot (\Pi_h - \Pi_H)\vec{\sigma}, q_h)_{L_2(D)\otimes L_2(\Omega)}, \end{aligned}$$

where in the last line we used the commutativity result given in (18). Applying the definition of the discrete gradient operator and the Cauchy-Schwarz inequality gives

$$\begin{aligned} \|q_h - \Theta_H q_h\|_{L_2(D)\otimes L_2(\Omega)}^2 &= ((\Pi_h - \Pi_H)\vec{\sigma}, \text{grad}_h^c q_h)_U \leq \|(\Pi_h - \Pi_H)\vec{\sigma}\|_U \|\text{grad}_h^c q_h\|_U \\ &\leq (\|\vec{\sigma} - \Pi_H \vec{\sigma}\|_U + \|\vec{\sigma} - \Pi_h \vec{\sigma}\|_U) \|\text{grad}_h^c q_h\|_U. \end{aligned}$$

Now applying the equivalence between $\|\cdot\|_U$ and $\|\cdot\|_{L_2(D)^2 \otimes L_2(\Omega)}$, the approximation bound given by (13), and noting that $h < H$ gives

$$\begin{aligned} \|q_h - \Theta_H q_h\|_{L_2(D)\otimes L_2(\Omega)}^2 &\leq a_{\max}^{1/2} (\|\vec{\sigma} - \Pi_H \vec{\sigma}\|_{L_2(D)^2 \otimes L_2(\Omega)} + \\ &\quad \|\vec{\sigma} - \Pi_h \vec{\sigma}\|_{L_2(D)^2 \otimes L_2(\Omega)}) \|\text{grad}_h^c q_h\|_U \\ &\leq a_{\max}^{1/2} (H + h) \|\vec{\sigma}\|_{H^1(D)^2 \otimes L_2(\Omega)} \|\text{grad}_h^c q_h\|_U \\ &\leq a_{\max}^{1/2} C_0 2HC_1 \|q_h - \Theta_H q_h\|_{L_2(D)\otimes L_2(\Omega)} \|\text{grad}_h^c q_h\|_U \end{aligned}$$

which establishes the theorem with $C_2 = 2C_0 C_1 a_{\max}^{1/2}$. \square

Theorem 8. Let $q_h \in Q_h \otimes T$ and $\vec{v}_h = \text{grad}_h^c q_h \in R_h \otimes T$. Define q_H and \vec{v}_H such that $v_H = \text{grad}_H^c q_H$ and $\nabla \cdot \vec{v}_H = \Theta_H \nabla \cdot \vec{v}_h$. Then

$$\|\vec{v}_h - \vec{v}_H\|_U \leq C_3 H \|\nabla \cdot \vec{v}_h\|_{L_2(D)\otimes L_2(\Omega)}$$

where C_3 is independent of h , m , and n .

Proof. As noted above, \vec{v}_h and \vec{v}_H can be considered to be finite element approximations to the flux solution of a first-order diffusion problem with source function $f = -\nabla \cdot \text{grad}_h^c q_h$. Let $\vec{v}_* \in H(\text{div}; D) \otimes T$ be the solution to the related semi-discrete problem, as described in §2.9. Then, using Theorem 1 and noting $H > h$,

$$\begin{aligned} \|\vec{v}_h - \vec{v}_H\|_U &\leq \|\vec{v}_* - \vec{v}_h\|_U + \|\vec{v}_* - \vec{v}_H\|_U \leq 2C_* H \|\nabla \cdot \text{grad}_h^c q_h\|_{L_2(D)\otimes L_2(\Omega)} \\ &= 2C_* H \|\nabla \cdot \vec{v}_h\|_{L_2(D)\otimes L_2(\Omega)}, \end{aligned}$$

which establishes the theorem with $C_3 = 2C_*$. \square

Theorem 9. Let $q_h \in Q_h \otimes T$ and $\vec{v}_h = \text{grad}_h^c q_h \in R_h \otimes T$. Define q_H and \vec{v}_H such that $\vec{v}_H = \text{grad}_H^c q_H$ and $\nabla \cdot \vec{v}_H = \Theta_H \nabla \cdot \vec{v}_h$. Then

$$\|\vec{v}_h - \vec{v}_H\|_\Lambda \leq C_4 H \|\mathcal{H}_h \vec{v}_h\|_U$$

where C_4 is independent of h , m , and n .

Proof. First, note that $\nabla \cdot \vec{v}_h \in Q_h \otimes T$. Applying Theorem 7 gives

$$\|\nabla \cdot \vec{v}_h - \nabla \cdot \vec{v}_H\|_{L_2(D) \otimes L_2(\Omega)} = \|\nabla \cdot \vec{v}_h - \Theta_H \nabla \cdot \vec{v}_h\|_{L_2(D) \otimes L_2(\Omega)} \leq C_2 H \|\text{grad}_h^c \nabla \cdot \vec{v}_h\|_U.$$

Then, using this along with Theorem 8 gives

$$\begin{aligned} \|\vec{v}_h - \vec{v}_H\|_\Lambda^2 &= \|\vec{v}_h - \vec{v}_H\|_U^2 + \|\nabla \cdot \vec{v}_h - \nabla \cdot \vec{v}_H\|_{L_2(D) \otimes L_2(\Omega)}^2 \\ &\leq C_3^2 H^2 \|\nabla \cdot \vec{v}_h\|_{L_2(D) \otimes L_2(\Omega)}^2 + C_2^2 H^2 \|\text{grad}_h^c \nabla \cdot \vec{v}_h\|_U^2 \end{aligned}$$

It can be shown, using the definition of the weighted $H(\text{div}; D) \otimes L_2(\Omega)$ operator given in §3.3, that

$$\|\mathcal{H}_h \vec{v}_h\|_U^2 = \|\vec{v}_h\|_\Lambda^2 + \|\nabla \cdot \vec{v}_h\|_{L_2(D) \otimes L_2(\Omega)}^2 + \|\text{grad}_h^c \nabla \cdot \vec{v}_h\|_U^2.$$

Therefore,

$$\|\vec{v}_h - \vec{v}_H\|_\Lambda \leq \max\{C_2, C_3\} H \|\mathcal{H}_h \vec{v}_h\|_U$$

which establishes the theorem with $C_4 = \max\{C_2, C_3\}$. \square

5.2. Λ -Projection Bounds. In this section we obtain two bounds involving a function in $R_h \otimes T$ and its projection with respect to $\Lambda(\cdot, \cdot)$ onto $R_H \otimes T$. These two bounds comprise the stochastic analogue of Lemma 3.2 in [1].

Theorem 10. *Let $\vec{v} \in R_h \otimes T$ and define $\vec{w} = \vec{v} - \mathcal{P}_H \vec{v} \in R_h \otimes T$ with Helmholtz Decomposition $\vec{w} = \text{grad}_h^c q + \nabla \times z$. Then*

$$\|\text{grad}_h^c q\|_U \leq C_4 H \|\vec{w}\|_\Lambda.$$

where C_4 is as in Theorem 9.

Proof. First note that

$$\Lambda(\text{grad}_h^c q, \mathcal{H}_h^{-1} \text{grad}_h^c q) = (\text{grad}_h^c q, \text{grad}_h^c q)_U = \|\text{grad}_h^c q\|_U^2.$$

Let $\vec{\tau}_h = \mathcal{H}_h^{-1} \text{grad}_h^c q$. Now, using (15), we have

$$\begin{aligned} \Lambda(\nabla \times z, \vec{\tau}_h) &= \int_\Omega \int_D \frac{1}{c} \nabla \times z \cdot \vec{\tau}_h + \int_\Omega \int_D (\nabla \cdot \nabla \times z)(\nabla \cdot \vec{\tau}_h) \\ &= \int_\Omega \int_D \frac{1}{c} \nabla \times z \cdot \vec{\tau}_h = (\nabla \times z, \vec{\tau}_h)_U. \end{aligned}$$

As \mathcal{H}_h (and hence \mathcal{H}_h^{-1}) maps $\text{grad}_h^c(Q_h \otimes T)$ onto itself (as noted in §3.3) there exists $q' \in Q_h \otimes T$ such that $\vec{\tau}_h = \text{grad}_h^c q'$. So, using (16),

$$\Lambda(\nabla \times z, \vec{\tau}_h) = (\nabla \times z, \text{grad}_h^c q')_U = -(\nabla \cdot \nabla \times z, q')_{L_2(D) \otimes L_2(\Omega)} = 0,$$

on account of (15) again. Define $\vec{\tau}_H$ in relation to $\vec{\tau}_h$ as in Theorem 9. Then

$$\begin{aligned} \Lambda(\text{grad}_h^c q, \vec{\tau}_h) &= \Lambda(\vec{v} - \mathcal{P}_H \vec{v}, \vec{\tau}_h) = \Lambda(\vec{v} - \mathcal{P}_H \vec{v}, \vec{\tau}_h - \vec{\tau}_H) \\ &\leq \|\vec{v} - \mathcal{P}_H \vec{v}\|_\Lambda \|\vec{\tau}_h - \vec{\tau}_H\|_\Lambda \leq C_4 H \|\vec{w}\|_\Lambda \|\text{grad}_h^c q\|_U, \end{aligned}$$

which completes the proof. \square

Theorem 11. *Let $\vec{v} \in R_h \otimes T$ and define $\vec{w} = \vec{v} - \mathcal{P}_H \vec{v} \in R_h \otimes T$ with Helmholtz Decomposition $\vec{w} = \text{grad}_h^c q + \nabla \times z$. Then*

$$\|z\|_{L_2(D) \otimes L_2(\Omega)} \leq C_5 H \|\vec{w}\|_\Lambda$$

where C_5 is independent of h , m , and n .

Proof. A standard result from the deterministic analysis (cf. [1]) yields

$$\|z\|_{L_2(D)} \leq C_6 H \|\nabla \times z\|_{L_2(D)^2} \quad \forall \omega \in \Omega$$

for some constant C_6 . Therefore,

$$\|z\|_{L_2(D) \otimes L_2(\Omega)} \leq C_6 H \|\nabla \times z\|_{L_2(D)^2 \otimes L_2(\Omega)} \leq C_6 a_{\min}^{-1/2} H \|\nabla \times z\|_U.$$

Now, using the properties of the Helmholtz decomposition noted in §3.1,

$$\begin{aligned} \|\nabla \times z\|_U^2 &= (\vec{w} - \text{grad}_h^c q, \nabla \times z)_U = (\vec{w}, \nabla \times z)_U \\ &= (\vec{w}, \nabla \times z)_U + (\nabla \cdot \vec{w}, \nabla \cdot \nabla \times z)_{L_2(D) \otimes L_2(D)} \\ &= \Lambda(\vec{w}, \nabla \times z) \leq \|\vec{w}\|_\Lambda \|\nabla \times z\|_U. \end{aligned}$$

Therefore $\|\nabla \times z\|_U \leq \|\vec{w}\|_\Lambda$ which combined with the above result establishes the theorem with $C_5 = C_6 a_{\min}^{-1/2}$. \square

6. ADDITIVE SCHWARZ METHOD

A constitutive part of multigrid is the smoother, where by smoother we mean an iterative method that successively smoothes the error associated with the iterates of the system that multigrid is applied to. Following [1] we will use the *additive Schwarz method* (extended to incorporate the stochastic nature of the problem) as our smoother. This method is defined with respect to a decomposition of the spatial domain D . In this section we will define the additive Schwarz operator and establish some results that will ultimately be used to demonstrate the convergence of the multigrid algorithm defined in §7.1.

6.1. Domain Decomposition Results. Let $\mathcal{D} = \{D_1, \dots, D_L\}$ be an over lapping covering of D . Let γ be an integer such that no point in \bar{D} occurs in more than γ elements of \mathcal{D} . Then $\sum_k \int_{D_k} \leq \gamma \int_D$ and γ is called the overlap parameter. Though we need not be specific here we note that in practice we will take each element of \mathcal{D} to be a set of triangles in \mathcal{T} that share a common node. For the partitions that we use to generate the numerical results in §8 this gives $\gamma = 3$. Now define

$$R^k \otimes T = \{ \vec{v} \in R \otimes T \mid \text{supp}(v) \subset D_k \times \Omega \}.$$

Given $\vec{v} \in R \otimes T$ there exists a decomposition $\vec{v} = \sum_k \vec{v}_k$ where $\vec{v}_k \in R^k \otimes T$. Let $\{\theta^k: \bar{D} \rightarrow \mathbb{R}\}_{k=1}^L$ be a partition of unity subordinate to the covering \mathcal{D} , i.e. $\sum_k \theta^k = 1$ and $\text{supp}(\theta^k) \subset D_k$.

Theorem 12. *Let $\vec{v} \in R \otimes T$ and $\vec{v}_k = \Pi \theta^k \vec{v} \in R^k \otimes T$. Then $\vec{v} = \sum_k \vec{v}_k$ and*

$$\sum_{k=1}^L \Lambda(\vec{v}_k, \vec{v}_k) \leq C_7 ((1 + h^{-2}) \|\vec{v}\|_U^2 + \|\nabla \cdot \vec{v}\|_{L_2(D) \otimes L_2(\Omega)}^2)$$

where C_7 is a constant.

Proof. Following [1], there exists a constant C_8 such that

$$\|\vec{v}_k\|_{L_2(D)^2} \leq C_8 \|\vec{v}\|_{L_2(D_k)^2} \quad \forall \omega \in \Omega.$$

Introducing the notation $\|\cdot\|_{U_k}$ to denote a norm defined in the same manner as $\|\cdot\|_U$ but restricted to D_k , we have

$$\|\vec{v}_k\|_U^2 \leq a_{\max} \|\vec{v}_k\|_{L_2(D)^2 \otimes L_2(\Omega)}^2 \leq C_8^2 a_{\max} \|\vec{v}\|_{L_2(D_k)^2 \otimes L_2(\Omega)}^2 \leq \frac{C_8^2 a_{\max}}{a_{\min}} \|\vec{v}\|_{U_k}^2.$$

Also following [1], there exists a constant C_9 such that

$$\|\nabla \cdot \vec{v}_k\|_{L_2(D)} \leq C_9 (h^{-1} \|\vec{v}\|_{L_2(D_k)^2} + \|\nabla \cdot \vec{v}\|_{L_2(D_k)}) \quad \forall \omega \in \Omega.$$

Therefore,

$$\|\nabla \cdot \vec{v}_k\|_{L_2(D) \otimes L_2(\Omega)}^2 \leq 2C_9^2 (h^{-2} a_{\min}^{-1} \|\vec{v}\|_{U_k}^2 + \|\nabla \cdot \vec{v}\|_{L_2(D) \otimes L_2(\Omega)}^2).$$

Then, using these two results, we have

$$\begin{aligned} \sum_{k=1}^L \Lambda(\vec{v}_k, \vec{v}_k) &= \sum_{k=1}^L (\|\vec{v}_k\|_U^2 + \|\nabla \cdot \vec{v}_k\|_{L_2(D) \otimes L_2(\Omega)}^2) \\ &\leq \sum_{k=1}^L \left(\frac{C_8^2 a_{\max}}{a_{\min}} \|\vec{v}_k\|_{U_k}^2 + 2C_9^2 \left(\frac{1}{h^2 a_{\min}} \|\vec{v}\|_{U_k}^2 + \|\nabla \cdot \vec{v}\|_{L_2(D) \otimes L_2(\Omega)}^2 \right) \right) \end{aligned}$$

which establishes the theorem with

$$C_7 = \gamma \max \left\{ \frac{C_8^2 a_{\max}}{a_{\min}}, 2C_9 \max \left\{ \frac{1}{a_{\min}}, 1 \right\} \right\},$$

where γ is the overlap parameter noted above. \square

6.2. Additive Schwarz Operator. We define the projection operator $\mathcal{P}^k: R \otimes T \rightarrow R^k \otimes T$ such that for $\vec{v} \in R \otimes T$,

$$\Lambda(\mathcal{P}^k \vec{v}, \vec{w}) = \Lambda(\vec{v}, \vec{w}) \quad \forall \vec{w} \in R^k \otimes T.$$

Note the similarity between \mathcal{P}^k and the projection operator \mathcal{P} defined in §3.2. The additive Schwarz operator is now defined by

$$\mathcal{S} = \eta \sum_{k=1}^L \mathcal{P}^k \mathcal{H}^{-1}$$

where η is some constant chosen such that Theorem 13 below holds. The additive Schwarz operator thus defined is symmetric with respect to $(\cdot, \cdot)_U$. To see this let $\vec{v}, \vec{w} \in R \otimes T$, then

$$\begin{aligned} (\mathcal{S}\vec{v}, \vec{w})_U &= \eta \sum_{k=1}^L (\mathcal{P}^k \mathcal{H}^{-1} \vec{v}, \vec{w})_U = \eta \sum_{k=1}^L \Lambda(\mathcal{P}^k \mathcal{H}^{-1} \vec{v}, \mathcal{H}^{-1} \vec{w}) \\ &= \eta \sum_{k=1}^L \Lambda(\mathcal{H}^{-1} \vec{v}, \mathcal{P}^k \mathcal{H}^{-1} \vec{w}) = \Lambda(\mathcal{H}^{-1} \vec{v}, \mathcal{S}\vec{w}) = (\vec{v}, \mathcal{S}\vec{w})_U. \end{aligned}$$

The operator \mathcal{S} can also be shown to be positive definite with respect to $(\cdot, \cdot)_U$ providing that, given $\vec{v} \in R \otimes T$ there exists a decomposition $\vec{v} = \sum_{k=1}^L \vec{v}_k$, $\vec{v}_k \in R^k \otimes T$ such that

$$(26) \quad \sum_{k=1}^L \Lambda(\vec{v}_k, \vec{v}_k) \leq C_\diamond \Lambda(\vec{v}, \vec{v})$$

where C_\diamond is a positive constant. To see this, we employ a standard argument as given, e.g., in [20]. Let $\mathcal{P}_S = \sum_{k=1}^L \mathcal{P}^k$. Then

$$\begin{aligned} \Lambda(\vec{v}, \vec{v}) &= \sum_{k=1}^L \Lambda(\vec{v}, \vec{v}_k) = \sum_{k=1}^L \Lambda(\mathcal{P}^k \vec{v}, \vec{v}_k) \leq \sum_{k=1}^L \|\mathcal{P}^k \vec{v}\|_\Lambda \|\vec{v}_k\|_\Lambda \\ &\leq \left(\sum_{k=1}^L \Lambda(\mathcal{P}^k \vec{v}, \mathcal{P}^k \vec{v}) \right)^{1/2} \left(\sum_{k=1}^L \Lambda(\vec{v}_k, \vec{v}_k) \right)^{1/2} \leq \left(\Lambda(\mathcal{P}_S \vec{v}, \vec{v}) \right)^{1/2} \left(C_\diamond \Lambda(\vec{v}, \vec{v}) \right)^{1/2}. \end{aligned}$$

Therefore,

$$\Lambda(\mathcal{P}_S \vec{v}, \vec{v}) \geq \frac{1}{C_\diamond} \Lambda(\vec{v}, \vec{v}).$$

Returning to the additive Schwarz operator and letting $\vec{w} = \mathcal{H}^{-1} \vec{v}$ gives

$$\begin{aligned} (\mathcal{S}\vec{v}, \vec{v})_U &= \eta \sum_{k=1}^L (\mathcal{P}^k \mathcal{H}^{-1} \vec{v}, \vec{v})_U = \eta \sum_{k=1}^L \Lambda(\mathcal{P}^k \mathcal{H}^{-1} \vec{v}, \mathcal{H}^{-1} \vec{v}) \\ &= \eta \sum_{k=1}^L \Lambda(\mathcal{P}^k \vec{w}, \vec{w}) = \eta \Lambda(\mathcal{P}_S \vec{w}, \vec{w}) \geq \frac{1}{C_\diamond} \eta \Lambda(\vec{w}, \vec{w}). \end{aligned}$$

As $\vec{v} = 0 \Leftrightarrow \vec{w} = 0$ this shows that \mathcal{S} is positive definite and hence invertible. That (26) holds for all $v \in R \otimes T$ can be shown by following a similar line of reasoning as in [1]. However, here we show that (26) holds on a necessary subspace of $R \otimes T$. This is done in the following section.

7. MULTIGRID

In the following we consider a family of triangulations \mathcal{T}_j , $j = 1, \dots, J$, with mesh parameters h_j , where $h_i > h_j$ for $i < j$. We assume these triangulations give rise to a nested sequence of finite-dimensional spaces denoted $R_1 \otimes T \subset \dots \subset R_J \otimes T \subset U$, as defined in §2.6. For the remainder of this section the various operators and spaces defined in §§2,3 will carry a subscript to denote the underlying triangulation on which they are defined. We let \mathcal{D}_j be a covering of D defined with respect to \mathcal{T}_j , as discussed in §6.1, and we let \mathcal{S}_j , $j = 2, \dots, J$, be the additive Schwarz operator defined with respect to \mathcal{D}_j .

7.1. Multigrid V-Cycle. We now define a multigrid algorithm, analogous to that given in [1], for solving the vector equation $\mathcal{H}_J \vec{v} = \vec{z}$. Let $\vec{z} \in R_k \otimes T$. Then we define $\mathcal{V}_k^{-1}: R_k \otimes T \rightarrow R_k \otimes T$ such that $\mathcal{V}_1^{-1} \vec{z} = \mathcal{H}_1^{-1} \vec{z}$ and $\mathcal{V}_k^{-1} \vec{z}$, $k = 2, \dots, J$, by the recursive algorithm:

```

 $\vec{v} = 0$ 
for  $j = 1, \dots, k$ 
   $\vec{v} \leftarrow \vec{v} + \mathcal{S}_j(\vec{z} - \mathcal{H}_j \vec{v})$ 
end
 $\vec{v} \leftarrow \vec{v} + \mathcal{V}_{j-1}^{-1} \Theta_{j-1}(\vec{z} - \mathcal{H}_j \vec{v})$ 
for  $j = 1, \dots, k$ 
   $\vec{v} \leftarrow \vec{v} + \mathcal{S}_j(\vec{z} - \mathcal{H}_j \vec{v})$ 
end

```

This corresponds to a multigrid V-cycle with one multigrid iteration at each level. Moreover, at each level there are k pre-smoothing and post-smoothing steps.

The matrix representation of the operator \mathcal{H}_J is $A_J^{-1}(A_J + F_J)$ where A_J and F_J are as in §4.2. Therefore, the vector equation $\mathcal{H}_J \vec{v} = \vec{z}$ can be expressed as the linear system $(A_J + F_J)\mathbf{v} = A_J \mathbf{z}$, where \mathbf{v} and \mathbf{z} are the coefficient vectors of \vec{v} and \vec{z} . This relationship along with the convergence properties of the above algorithm will be used in §7.2 to demonstrate that the spectral bounds in (24) will be independent of the parameters h , m , and n if the matrix V_J is chosen such that V_J^{-1} is the matrix representation of the multigrid operator \mathcal{V}_J^{-1} .

7.2. Multigrid Convergence. We here establish that the multigrid algorithm given in §7.1 converges and that \mathcal{V}_J and \mathcal{H}_J are spectrally equivalent.

Theorem 13. *The inequality*

$$\Lambda((I - \mathcal{S}_j \mathcal{H}_j) \vec{v}, \vec{v}) \geq 0 \quad \forall \vec{v} \in R_j \otimes T$$

holds for $j = 1, \dots, J$.

Proof. For $j = 1$, $\mathcal{S}_j = \mathcal{H}_j^{-1}$ and the result holds as $\Lambda(\cdot, \cdot)$ is an inner product on $R_1 \otimes T$. Let $2 < j \leq J$. Then

$$\begin{aligned} \Lambda((I - \mathcal{S}_j \mathcal{H}_j) \vec{v}, \vec{v}) &= \Lambda(\vec{v}, \vec{v}) - \Lambda(\mathcal{S}_j \mathcal{H}_j \vec{v}, \vec{v}) = \Lambda(\vec{v}, \vec{v}) - \eta \sum_{k=1}^{L_j} \Lambda(\mathcal{P}_j^k \vec{v}, \vec{v}) \\ &= \Lambda(\vec{v}, \vec{v}) - \eta \sum_{k=1}^{L_j} \Lambda(\mathcal{P}_j^k \vec{v}, \mathcal{P}_j^k \vec{v}) = \Lambda(\vec{v}, \vec{v}) - \eta \sum_{k=1}^{L_j} \|\mathcal{P}_j^k \vec{v}\|_{\Lambda}^2. \end{aligned}$$

Let $\|\cdot\|_{\Lambda_k}$ be the norm induced on $D_k \times \Omega$ by $\|\cdot\|_{\Lambda}$. Then

$$\|\mathcal{P}_j^k \vec{v}\|_{\Lambda}^2 = \Lambda(\mathcal{P}_j^k \vec{v}, \mathcal{P}_j^k \vec{v}) = \Lambda(\mathcal{P}_j^k \vec{v}, \vec{v}) \leq \|\mathcal{P}_j^k \vec{v}\|_{\Lambda_k} \|\vec{v}\|_{\Lambda_k} = \|\mathcal{P}_j^k \vec{v}\|_{\Lambda} \|\vec{v}\|_{\Lambda_k}.$$

Therefore,

$$\Lambda((I - \mathcal{S}_j \mathcal{H}_j) \vec{v}, \vec{v}) \leq (1 - \eta \gamma) \Lambda(\vec{v}, \vec{v})$$

where γ is the overlap parameter discussed in §6.1. So the theorem holds providing $\eta \leq \gamma^{-1}$. \square

Lemma 2. Let $\vec{v} \in R_j \otimes T$ and let $\vec{w} \in (I - \mathcal{P}_{j-1})\vec{v} \in R_j \otimes T$. Then, for $j = 2, \dots, J$, there exists a decomposition $\vec{w} = \sum_{k=1}^{L_j} \vec{w}_k$, $\vec{w}_k \in R_j^k \otimes T$ such that

$$(27) \quad \sum_{k=1}^{L_j} \Lambda(\vec{w}_k, \vec{w}_k) \leq C_{10} \Lambda(\vec{w}, \vec{w})$$

where C_{10} is independent of h , m , and n .

Proof. Let \vec{w} have the Helmholtz decomposition $\vec{w} = \text{grad}_{h_j}^c q + \nabla \times s$, where $q \in Q_j \otimes T$ and $s \in W_j \otimes T$. Define $\vec{w}' = \text{grad}_{h_j}^c q$ and $\vec{w}'' = \nabla \times s$. Then given the decompositions $\vec{w}' = \sum_k \vec{w}'_k$ and $\vec{w}'' = \sum_k \vec{w}''_k$ and taking into account that the Helmholtz decomposition is orthogonal with respect to $\Lambda(\cdot, \cdot)$, it is sufficient to show that

$$\sum_{k=1}^{L_j} \Lambda(\vec{w}'_k, \vec{w}'_k) \leq C'_{10} \Lambda(\vec{w}', \vec{w}'), \quad \sum_{k=1}^{L_j} \Lambda(\vec{w}''_k, \vec{w}''_k) \leq C''_{10} \Lambda(\vec{w}'', \vec{w}'')$$

for some constants C'_{10} and C''_{10} . This will result in $C_{10} = 2 \max\{C'_{10}, C''_{10}\}$. To show the first of these, let $w'_k = \Pi_j \theta_j^k w'$ where θ_j^k is as in §6.2. Invoking Theorems 10 and 12 gives

$$\begin{aligned} \sum_{k=1}^{L_j} \Lambda(\vec{w}'_k, \vec{w}'_k) &\leq C_7((1 + h_j^{-2})\|\vec{w}'\|_U^2 + \|\nabla \cdot \vec{w}'\|_{L_2(D) \otimes L_2(\Omega)}^2) \\ &= C_7(\|\vec{w}'\|_\Lambda^2 + h_j^{-2}\|\vec{w}'\|_U^2) \leq C_7(\|\vec{w}'\|_\Lambda^2 + C_4^2 h_j^{-2} h_{j-1}^2 \|\vec{w}\|_\Lambda^2). \end{aligned}$$

Assume that for $j = 2, \dots, J$, there exists $\rho > h_{j-1}/h_j$, then, as $\|\vec{w}'\|_\Lambda \leq \|\vec{w}\|_\Lambda$,

$$\sum_{k=1}^{L_j} \Lambda(\vec{w}'_k, \vec{w}'_k) \leq 2C_7 \max\{1, C_4^2 \rho^2\} \Lambda(\vec{w}, \vec{w}).$$

Next, following [1], there exists a decomposition $\vec{w}'' = \sum \vec{w}''_k$ such that

$$\sum_{k=1}^{L_j} (\vec{w}''_k, \vec{w}''_k)_{L_2(D)^2} \leq C_{11}(\|\vec{w}''\|_{L_2(D)^2}^2 + h_j^{-2}\|s\|_{L_2(D)}^2) \quad \forall \omega \in \Omega,$$

where C_{11} is a constant. From this it follows that

$$\sum_{k=1}^{L_j} \Lambda(\vec{w}''_k, \vec{w}''_k) \leq a_{\max} C_{11} (a_{\min}^{-1} \|w''\|_\Lambda^2 + h_j^{-2} \|s\|_{L_2(D) \otimes L_2(\Omega)}^2).$$

Applying Theorem 11 and noting that $\|\vec{w}''\|_\Lambda \leq \|\vec{w}\|_\Lambda$ gives

$$\sum_{k=1}^{L_j} \Lambda(\vec{w}''_k, \vec{w}''_k) \leq 2 a_{\max} C_{11} \max\{a_{\min}^{-1}, C_5^2 \rho^2\} \Lambda(\vec{w}, \vec{w}).$$

Therefore the theorem is established. \square

Lemma 3. Let $\vec{v} \in R_j \otimes T$ and $\vec{w} = (I - \mathcal{P}_{j-1})\vec{v}$. Then for $j = 2, \dots, J$,

$$(\mathcal{S}_j^{-1} \vec{w}, \vec{w})_U = \frac{1}{\eta} \inf \sum_{k=1}^{L_j} \Lambda(\vec{w}_k, \vec{w}_k)$$

where the infimum is taken over all decompositions of the form $\vec{w} = \sum_k \vec{w}_k$, $\vec{w}_k \in R_j^k \otimes T$.

Proof. Let $E_j = \{\vec{w} \in R_j \otimes T \mid \vec{w} = (I - \mathcal{P}_{j-1})\vec{v}, \vec{v} \in R_j \otimes T\}$ for $j = 2, \dots, J$. As a consequence of Lemma 2, (24) is satisfied on each of these sets and, therefore, \mathcal{S}_j is invertible on E_j . The rest of the proof follows in an analogous fashion to the proof for the deterministic case given in Appendix B of [1]. \square

Theorem 14. *The inequality*

$$(\mathcal{S}_j^{-1}(I - \mathcal{P}_{j-1})\vec{v}, (I - \mathcal{P}_{j-1})\vec{v})_U \leq C_{12} \Lambda((I - \mathcal{P}_{j-1})\vec{v}, (I - \mathcal{P}_{j-1})\vec{v}) \quad \forall \vec{v} \in R_j \otimes T$$

holds for $j = 2, \dots, J$, where C_{12} is independent of h , m , and n .

Proof. Let $\vec{w} = (I - \mathcal{P}_{j-1})\vec{v}$. Then for each $\vec{v} \in R_j \otimes T$ it is required to show that

$$(\mathcal{S}^{-1}\vec{w}, \vec{w})_U \leq C_{12}\Lambda(\vec{w}, \vec{w}).$$

From Lemma 2 we know there exists a decomposition $\vec{w} = \sum_k \vec{w}_k$, $\vec{w}_k \in R_j^k \otimes T$ such that

$$\sum_{k=1}^{L_j} \Lambda(\vec{w}_k, \vec{w}_k) \leq C_{10}\Lambda(\vec{w}, \vec{w}).$$

Now using Lemma 3 gives the required inequality with $C_{12} = \eta^{-1}C_{10}$. \square

Theorem 15. *The eigenvalues of $\mathcal{V}_J^{-1}\mathcal{H}_J$ are contained in the interval $[1 - \delta, 1]$ where*

$$\delta = \frac{C_{12}}{C_{12} + 2k}.$$

Proof. Given Theorem 13 and Theorem 15, the proof follows that for Corollary 5.2 in [1]. \square

Note that δ , thus defined, depends on k , which denotes the number of pre-smoothing and post-smoothing steps in the multigrid algorithm, and $C_{12} = \eta^{-1}C_{10}$, where C_{10} is independent of the discretization parameters, h , m , and n , but does depend on the lower and upper bounds of the diffusion coefficient, a_{\min} and a_{\max} .

From Theorem 15 we have

$$1 - \delta \leq \frac{(\mathcal{H}_J\vec{v}, \vec{v})_U}{(\mathcal{V}_J\vec{v}, \vec{v})_U} \leq 1 \quad \forall \vec{v} \in V_J.$$

As noted in §7.1 the multigrid algorithm finds an approximation, \vec{v}_1 say, to the solution of $\mathcal{H}_J\vec{v} = \vec{z}$. Moreover, $\mathcal{H}_J\vec{v} = \vec{z}$ can be expressed as $(A_J + F_J)\mathbf{v} = A_J\mathbf{z}$ where A_J and F_J are as in §4.2 and \mathbf{v} and \mathbf{z} are the coefficient vectors of \vec{v} and \vec{z} . Now we define the matrix V_J such that $V_J^{-1}A_J\mathbf{z} = \mathbf{v}_1$ where \mathbf{v}_1 is the coefficient vector of \vec{v}_1 . That is to say, V_J^{-1} is the matrix representation of the multigrid operator \mathcal{V}_J^{-1} . With V_J defined in this way we have

$$\frac{(\mathcal{H}_J\vec{v}, \vec{v})_U}{(\mathcal{V}_J\vec{v}, \vec{v})_U} = \frac{\Lambda(\vec{v}, \vec{v})}{(\mathcal{V}_J\vec{v}, \vec{v})_U} = \frac{((A_J + F_J)\mathbf{v}, \mathbf{v})}{(A_J^{-1}V_J\mathbf{v}, \mathbf{v})_{A_J}} = \frac{((A_J + F_J)\mathbf{v}, \mathbf{v})}{(V_J\mathbf{v}, \mathbf{v})}.$$

Therefore, (24) is satisfied with $\theta = 1 - \delta$ and $\bar{\theta} = 1$. Consequently, as θ governs the values of the eigenvalue bounds described in Theorem 6, we expect the preconditioned MINRES scheme given in §4.2 to converge independently of the discretization parameters but not of the upper and lower bounds of the diffusion coefficient.

8. NUMERICAL IMPLEMENTATION

Our aim is to solve (12) via MINRES using the preconditioner (25). Recall that N is a diagonal matrix and the action of V^{-1} is achieved by applying a V-cycle of the multigrid algorithm analysed in §7 to a system with the coefficient matrix

$$A + F = G_0 \otimes (K_0 + D_0) + \sum_{r=1}^m \sqrt{\lambda_r} G_r \otimes K_r.$$

The cost of computing the action of V^{-1} is clearly a crucial consideration and we now offer some practical insight into this.

The intergrid transfer operators are defined in the usual way. That is, prolongation is achieved via natural inclusion and restriction is defined via transposition. To describe the additive Schwarz smoother, let $H_j = A_j + F_j$ denote the stochastic weighted $H(\text{div})$ matrix associated with the triangulation \mathcal{T}_j . This triangulation is decomposed into L_j overlapping patches which constitute the elements of \mathcal{D}_j as described in §6.1. Then the matrix representation of \mathcal{S}_j is given by

$$S_j = \eta \sum_{k=1}^{L_j} P_j^k H_j^{-1}, \quad P_j^k = (I \otimes J_j^k)^T (H_j^k)^{-1} (I \otimes J_j^k) H_j,$$

where $(I \otimes J_j^k)$ is the patch restriction matrix, I is the $M \times M$ identity matrix, and $J_j^k \in \mathbb{R}^{N_{R_j} \times N_{R_j}}$ is the standard matrix with entries zero or one that extracts the components of

$\mathbf{w} \in \mathbb{R}^{N_{R_j}}$ which are associated with the k -th patch in \mathcal{D}_j . Following the advice in [1], we choose the scaling parameter in the definition of S_j to be $\eta = 1/2$.

Now, computing $S_j \mathbf{v}$, where $\mathbf{v} \in \mathbb{R}^{MN_{R_j}}$, requires local solves with the matrices

$$(28) \quad H_j^k = G_0 \otimes (K_{0,j}^k + D_{0,j}^k) + \sum_{r=1}^m \sqrt{\lambda_r} G_r \otimes K_{r,j}^k, \quad k = 1, \dots, L_j,$$

where $K_{0,j}^k, D_{0,j}^k$, and $K_{r,j}^k$ are small principal submatrices of $K_{0,j}, D_{0,j}$, and $K_{r,j}$, respectively. The major strength of this method is that the L_j solves are decoupled and can (and should) be performed in parallel. The size of H_j^k is approximately $6M \times 6M$ where M is defined in (8) and six is the typical number of flux degrees of freedom lying on a patch in \mathcal{D}_j . We assume that these L_j systems are solved via direct methods and so a few thousand stochastic degrees of freedom can be handled comfortably in a parallel computing environment.

If parallel computing facilities are not available or M is in the order of hundreds of thousands or more, then a cheaper smoother may have to be considered. We outline one possibility. Suppose that we define a modified smoother \hat{S}_j via

$$\hat{S}_j = \eta \sum_{k=1}^{L_j} P_j^k \hat{H}_j^{-1}, \quad P_j^k = (I \otimes J_j^k)^T (\hat{H}_j^k)^{-1} (I \otimes J_j^k) \hat{H}_j$$

where, now,

$$\hat{H}_j^k = G_0 \otimes (K_{0,j}^k + D_{0,j}^k), \quad k = 1, \dots, L_j.$$

Then, \hat{H}_j^k is a local version of the positive definite matrix

$$\hat{H}_j = G_0 \otimes (K_{0,j} + D_{0,j})$$

which is a stochastic $H(\text{div})$ matrix with a deterministic weight, that provides a representation of the norm,

$$\|c_0^{-\frac{1}{2}} \bar{v}\|_{L^2(D)^2 \otimes L^2(\Omega)}^2 + \|\nabla \cdot \bar{v}\|_{L^2(D) \otimes L^2(\Omega)}^2$$

on $R_j \otimes T$. Crucially, since G_0 is diagonal, then

$$(\hat{H}_j^k)^{-1} = G_0^{-1} \otimes (K_{0,j}^k + D_{0,j}^k)^{-1}$$

is block-diagonal and computing $\hat{S}_j \mathbf{v}$ now requires $L_j M$ decoupled solves with matrices of dimension approximately six. Each of these solves is a trivial cost and, moreover, there are two levels of parallelism to exploit.

The analysis in §7 is valid only for the multigrid method that uses the smoother (28). Consequently, we have no theoretical bounds for the constants θ and $\bar{\theta}$ in (24) when V corresponds to the modified method. However, a preliminary analysis suggests that if $c_0(\bar{x})$ dominates in the expression (2.1), the resulting multigrid approximation is efficient.

8.1. Model problem. We now solve (2) with $f = 1$ using uniformly refined meshes of triangular finite elements on $D = (-1, 1) \times (-1, 1)$. The spatial grid consists of $d \times d$ squares each of which is further subdivided into two equal triangles. The random diffusion coefficient $c(\bar{x}, \bar{\xi}(\omega))^{-1}$ is chosen to be of the form (2.1) with the covariance function

$$(29) \quad C(\bar{x}, \bar{y}) = \sigma^2 \exp(-|x_1 - y_1| - |x_2 - y_2|).$$

Note that the eigenvalues and eigenfunctions of this function are known explicitly (see [11]). The random variables $\{\xi_r\}_{r=1}^m$ in (2.1) are assumed to be independent and identically distributed with zero mean and unit variance. If uniform random variables are selected then $\Omega = [-\sqrt{3}, \sqrt{3}]^m$ and multivariate Legendre polynomials of total degree n are used to construct the polynomials chaos basis for T . Alternatively, if Gaussian random variables are selected then $\Omega = \mathbb{R}^m$ and multivariate Hermite polynomials are employed.

We investigate the performance of the preconditioner (25) with respect to the discretisation parameters h, m and n , the distribution of the underlying random variables, ξ_r , and the mean $c_0(\bar{x})$ and variance σ^2 of $c(\bar{x}, \bar{\xi}(\omega))^{-1}$. In each experiment, we report preconditioned MINRES iteration counts and assess the efficiency of the multigrid approximation by computing the

constants θ and $\bar{\theta}$ from (24). In each MINRES iteration, a single V-cycle of multigrid is applied with a single pre- and post-smoothing step per grid level. The iteration is terminated when the relative residual reaches a tolerance of 10^{-6} . All experiments were performed on a modest single-processor Linux machine with 2GB RAM.

8.2. Robustness with respect to discretisation parameters. First we fix the mean and standard deviation of $c(\vec{x}, \vec{\xi}(\omega))^{-1}$ to be $c_0 = 1$ and $\sigma = 0.1$. Iteration counts for solving this problem, using two types of random variables, are listed in Tables 1 and 2. The numbers in parentheses in Table 1 indicate iterations obtained with the exact preconditioner (23).

	n	Uniform				Gaussian			
		$m = 1$	$m = 2$	$m = 3$	$m = 4$	$m = 1$	$m = 2$	$m = 3$	$m = 4$
$d = 16$	1	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)
-	2	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)
-	3	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (6)
-	4	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)
$d = 32$	1	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)
-	2	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)
-	3	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (6)	17 (5)
-	4	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (5)	17 (6)	17 (5)

TABLE 1. Preconditioned MINRES iterations using multigrid with original smoother

	n	Uniform				Gaussian			
		$m = 1$	$m = 2$	$m = 3$	$m = 4$	$m = 1$	$m = 2$	$m = 3$	$m = 4$
$d = 16$	1	17	17	17	18	17	17	17	17
-	2	17	18	19	19	17	19	19	19
-	3	17	19	19	19	18	19	19	19
-	4	17	19	19	19	18	19	19	19
$d = 32$	1	17	17	17	18	17	17	17	18
-	2	17	18	18	18	18	18	18	19
-	3	17	18	18	18	18	18	18	19
-	4	17	18	18	18	18	18	19	19

TABLE 2. Preconditioned MINRES iterations using multigrid with modified smoother

The results for the preconditioner based on the original smoother confirm the analysis of §7 and §4.2. Iteration counts are independent of all the discretisation parameters, and in addition, the distribution of the underlying random variables. In this example, modifying the multigrid smoother also has little to no effect on the performance of the preconditioning scheme. To reinforce this, in Table 3, we list the numerically computed constants from (24) for both variants of the multigrid method.

8.3. Robustness with respect to a_{max} . The experiment above is not challenging from a statistical point of view. We know from Theorems 5 and 6 that iteration counts are likely to be sensitive to the value a_{max} in (5), even if the multigrid approximation is not. Here, we vary the mean c_0 and fix $\sigma = c_0/10$ so that c_0 is the dominant term in (2.1) and $a_{max} = O(c_0)$. We fix the discretisation parameters and choose uniform random variables.

The results in Table 4 tell us that both multigrid approximations improve as $c_0 \rightarrow \infty$ and neither one deteriorates as $c_0 \rightarrow 0$. The deterioration in the MINRES iteration counts as $c_0 \rightarrow \infty$ is not caused by a deficiency in the multigrid approximation but rather is a feature of the underlying exact preconditioner (23), as indicated by the eigenvalue bound established in Theorem 5. This is a known feature of the corresponding deterministic preconditioner. Ideally, (2) should be rescaled so that for the re-scaled diffusion coefficient, $a_{max} \leq 1$.

	n	$m = 1$	$m = 2$	$m = 3$
Original (Uniform)	1	[0.4589,1.0000]	[0.4589,1.0000]	[0.4589,1.0000]
-	2	[0.4586,1.0000]	[0.4587,1.0000]	[0.4587,1.0000]
Original (Gaussian)	1	[0.4589,1.0000]	[0.4589,1.0000]	[0.4578,1.0000]
-	2	[0.4583,1.0000]	[0.4584,1.0000]	[0.4584,1.0000]
Modified (Uniform)	1	[0.4235,1.0003]	[0.4217,1.0009]	[0.4205,1.0013]
-	2	[0.4149,1.0000]	[0.4103,1.0008]	[0.4079,1.0012]
Modified (Gaussian)	1	[0.4235,1.0003]	[0.4217,1.0009]	[0.4205, 1.0013]
-	2	[0.4049,1.0004]	[0.4024,1.0008]	[0.3862, 1.0009]

TABLE 3. Multigrid constants, $d = 8$, $[\theta, \bar{\theta}]$, Uniform random variables

c_0	σ	Exact	Original		Modified	
		iter	iter	$[\theta, \bar{\theta}]$	iter	$[\theta, \bar{\theta}]$
10^{-3}	10^{-4}	3	15	[0.4552,1.0000]	16	[0.4068,1.0008]
10^{-2}	10^{-3}	3	16	[0.4550,1.0000]	16	[0.4068,1.0008]
10^{-1}	10^{-2}	4	16	[0.4556,1.0000]	17	[0.4071,1.0008]
10^0	10^{-1}	5	17	[0.3956,1.0000]	18	[0.4103,1.0008]
10^1	10^0	8	21	[0.4864,1.0000]	22	[0.4387,1.0005]
10^2	10^1	16	40	[0.6453,1.0000]	45	[0.5990,1.0000]
10^3	10^2	46	99	[0.9172,1.0000]	107	[0.8660,1.0013]

TABLE 4. MINRES iterations and multigrid constants $m = 4$, $n = 2$, $d = 32$

8.4. Robustness with respect to the ratio $c_0^{-1}\sigma$. It is known that preconditioners for stochastic finite element systems that are based on the mean-component of the diffusion coefficient are cheap but lose efficiency when $c_0^{-1}\sigma \rightarrow \infty$. Theorems 6 and 15 tell us that if V is based on our original multigrid method, then the efficiency of (25) is affected by a_{max} , and possibly a_{min} , but not directly by the ratio $c_0^{-1}\sigma \rightarrow \infty$ (if a_{max} remains bounded). To illustrate this, we fix $c_0 = 1$ and vary only σ . We fix all the discretisation parameters and choose uniform random variables.

σ/c_0	0.1	0.2	0.3	0.4	0.5	0.6	0.7
Exact	5	5	5	5	5	5	5
Original	17	17	17	17	17	19	19
Modified	18	20	21	24	28	*	*

TABLE 5. MINRES iteration counts, $m = 4$, $n = 2$, $c_0 = 1$, $d = 32$

The iteration counts listed in Table 5 confirm that the preconditioning scheme based on the original multigrid algorithm is insensitive to $c_0^{-1}\sigma$. Iteration counts deteriorate, however, for the modified version and MINRES stagnates when $c_0^{-1}\sigma$ exceeds a certain threshold (indicated by *). This is to be expected since the matrix

$$\sum_{r=1}^m \sqrt{\lambda_r} G_r \otimes K_r$$

which encapsulates the deviation of the diffusion coefficient from its mean value is not taken into account in the smoothing. This causes the quality of the multigrid approximation to deteriorate. To reinforce this, we list the associated multigrid constants in Table 6 for $d = 8$. These results reinforce our earlier comment that the modified method should only be considered if the deviation of $c(\vec{x}, \vec{\xi}(\omega))^{-1}$ from the mean field is small.

σ/c_0		0.4	0.5	0.6
$m = 2$	$n = 1$	[0.2657, 1.0000]	[0.2268, 1.0000]	[0.1915, 1.0000]
	$n = 2$	[0.2155, 1.0000]	[0.1651, 1.0000]	[0.1187, 1.0000]
$m = 3$	$n = 1$	[0.2574, 1.0000]	[0.2179, 1.0000]	[0.1822, 1.0000]
	$n = 2$	[0.2072, 1.0000]	[0.1529, 1.0000]	[0.1035, 1.0000]
$m = 4$	$n = 1$	[0.2194, 1.0000]	[0.2090, 1.0000]	[0.1727, 1.0000]
	$n = 2$	[0.1949, 1.0000]	[0.1078, 1.0000]	[-0.5426, 1.0000]

TABLE 6. Multigrid constants $[\theta, \bar{\theta}]$ for modified smoother, $c_0 = 1$, $d = 8$

9. ACKNOWLEDGEMENTS

We would like to thank John Osborn, Richard Falk, Douglas Arnold, and Ragnar Winther for several useful discussions and in particular for their assistance in establishing Lemma 1.

REFERENCES

- [1] Douglas N. Arnold, Richard S. Falk, and R. Winther. Preconditioning in $H(\text{div})$ and applications. *Math. Comp.*, 66:957–984, 1997.
- [2] Ivo Babuška, Raúl Tempone, and Georgios E. Zouraris. Galerkin finite element approximation of stochastic elliptic partial differential equations. *SIAM J. Numer. Anal.*, 42:800–825, 2004.
- [3] Dietrich Braess. *Finite Elements: Theory, Fast Solvers, and Applications in Solid Mechanics*. Cambridge University Press, second edition, 2001.
- [4] Franco Brezzi and Michel Fortin. *Mixed and Hybrid Finite Element Methods*. Springer-Verlag, 1991.
- [5] Zdzisław Brzeźniak and Tomasz Zastawniak. *Basis Stochastic Processes*. Springer-Verlag, 1999.
- [6] Claudio Canuto and Tomas Kozubek. A fictitious domain approach to the numerical solution of pdes in stochastic domains. *Numer. Math.*, 107:257–293, 2007.
- [7] Manas K. Deb, Ivo M. Babuška, and J. Tinsley Oden. Solution of stochastic partial differential equations using Galerkin finite element techniques. *Comput. Methods Appl. Mech. Eng.*, 190:6359–6372, 2001.
- [8] Howard Elman and Darran Furnival. Solving the stochastic steady-state diffusion problem using multigrid. *IMA. J. Numer. Anal.*, 27:675–688, 2007.
- [9] O. G. Ernst, C. E. Powell, D. J. Silvester, and E. Ullman. Efficient solvers for a linear stochastic galerkin mixed formulation of diffusion problems with random data. Technical Report 2007.126, University of Manchester, 2007.
- [10] Avner Friedman. *Foundations of Modern Analysis*. Dover, 1982.
- [11] Roger G. Ghanem and Pol D. Spanos. *Stochastic Finite Elements: A Spectral Approach*. Springer-Verlag, 1991.
- [12] R. Hiptmair. Multigrid method for $H(\text{div})$ in three dimensions. *ETNA*, 6:133–152, 1997.
- [13] O. P. Le Maître, O. M. Knio, B. J. Debuschere, H. N. Najm, and R. G. Ghanem. A multigrid solver for two-dimensional stochastic diffusion equations. *Comput. Methods Appl. Mech. Engrg.*, 192:4723–4744, 2003.
- [14] P. Vassilevski P and J. Wang. Multilevel iterative methods for mixed finite element discretizations of elliptic problems. *Numer. Math.*, 63:503–520, 1992.
- [15] C. C. Paige and M. A. Saunders. Solution of sparse indefinite systems of linear equations. *SIAM J. Numer. Anal.*, 12:617–629, 1975.
- [16] Catherine E. Powell. Parameter-free $H(\text{div})$ preconditioning for mixed finite element formulation of diffusion problems. *IMA J. Num. Anal.*, 25:783–796, 2005.
- [17] Catherine Elizabeth Powell and David Silvester. Optimal preconditioning for Raviart-Thomas mixed formulation of second-order problems. *SIAM J. Matrix Anal. Appl.*, 25:718–738, 2003.
- [18] P. A. Raviart and J. M. Thomas. A mixed finite element method for second order elliptic problems. In I. Galligani and E. Magenes, editors, *Mathematical Aspects of the Finite Element Method*, no. 606 in Lect. Notes in Math., pages 292–315. Springer-Verlag, New York, 1977.
- [19] Wim Schoutens. *Stochastic Processes and Orthogonal Polynomials*. Springer-Verlag, 2000.
- [20] A. Toselli and O. Widlund. *Domain Decomposition Methods - Algorithms and Theory*. Springer-Verlag, 2005.
- [21] François Trèves. *Topological Vector Spaces, Distributions and Kernels*. Academic Press, 1967.
- [22] Dongbin Xiu. *Generalized (Wiener-Askey) Polynomial Chaos*. PhD thesis, Brown University, 2004.
- [23] Dongbin Xiu and George Em Karniadakis. Modeling uncertainty in steady state diffusion problems via generalized polynomial chaos. *Comput. Methods Appl. Engrg.*, 191:4927–4948, 2002.
- [24] C. I. Goldstein Z. Cai and J. E. Pasciak. Multilevel iteration for mixed finite element systems with penalty. *SIAM J. Sci. Comput.*, 14(5):1072–1088, 1993.