

Lacunarity and Period-doubling

Glendinning, Paul and Smith, Leonard A.

2012

MIMS EPrint: **2012.73**

Manchester Institute for Mathematical Sciences
School of Mathematics

The University of Manchester

Reports available from: <http://eprints.maths.manchester.ac.uk/>

And by contacting: The MIMS Secretary
School of Mathematics
The University of Manchester
Manchester, M13 9PL, UK

ISSN 1749-9097

Lacunarity and Period-doubling

Paul Glendinning^a, Leonard A. Smith^b

^a*Centre for Interdisciplinary Computational and Dynamical Analysis (CICADA) and
School of Mathematics, University of Manchester, Manchester, M13 9PL, U.K.*

Email: p.a.glendinning@manchester.ac.uk

^b*Centre for the Analysis of Time Series (CATS), London School of Economics &
Political Science Houghton Street, London, WC2A 2AE, U.K.*

Email: l.smith@lse.ac.uk

Abstract

We show that the deviation from power laws of the scaling of chaotic measures such as Lyapunov exponents and topological entropy is periodic in the logarithm of the distance from the accumulation of period doubling. Moreover, this periodic function is asymptotically universal for each measure (for functions in the appropriate universality class). This is related to the concept of lacunarity known to exist for scaling functions describing the mass distribution of self-similar fractal sets.

Keywords: lacunarity, fractal, period-doubling, universality class

1. Lacunarity

In self-similar sets, lacunarity describes the way in which voids or lacunae are distributed at all scales. The presence of these voids is reflected in scaling functions which characterize the spatial distribution of points. In self-similar sets, the regular appearance of voids results in a periodic component of the scaling functions; such oscillations, however, are often found in scaling quantities, an observation going back at least to DeBruin ([6], see [17] for a historical review). Guckenheimer [12] suggests that the oscillations in scaling functions seriously hamper numerical attempts to approximate scaling exponents, such as the capacity of fractal sets. These arguments are supported by observations [1, 19, 23, 20]. For a more recent discussion see [24] and references therein. In these studies the effect of lacunarity is confined to problems involving spatial structures; here we show how the same ideas are useful in describing scaling behaviour in families of dynamical systems,

where the (self-similar) fractal structure occurs in parameter space rather than in state space. In particular, we show that universal functions describe the asymptotic structure of quantities such as the topological entropy of a map just after the accumulation of period-doubling ¹.

The source of the lacunarity described here is the existence of self-similar scaling relations of the form $af(x) = f(bx)$ for the quantities we describe. This equation has general solution

$$f(x) = \psi(\log x)x^{\frac{\log a}{\log b}},$$

where $\psi(\log x)$ is periodic with period $\log b$, and the lacunarity of the set is reflected in the oscillation due to ψ .

As an introduction to lacunarity we shall consider a simple example in which all the functions involved can be calculated explicitly (see [18] for additional discussion). The “middle thirds” Cantor set, Λ , is perhaps the most easy fractal set to work with, and the periodic oscillation of the scaling function can be calculated explicitly. The set

$$\Lambda = \{x \in [0, 1] \mid x = \sum_{i=1}^{\infty} \frac{a_i}{3^i}, a_i \in \{0, 2\}\} \quad (1)$$

is a Cantor set since it is uncountable, closed, contains no intervals, and every point is an accumulation point of the set. The standard construction of Λ is inductive: take the interval $[0, 1]$ and remove the middle third, $(\frac{1}{3}, \frac{2}{3})$, leaving two intervals $[0, \frac{1}{3}]$ and $[\frac{2}{3}, 1]$. Now remove the middle third of each of these intervals and repeat the construction, removing the middle third of each remaining interval at each stage. In the limit, this process defines the Cantor set, Λ . Let $N(\epsilon)$ be the smallest number of ϵ -balls needed to cover Λ . The dimension or capacity of Λ , $C(\Lambda)$, is then defined as

$$C(\Lambda) = \lim_{\epsilon \rightarrow 0} -\frac{\log N(\epsilon)}{\log \epsilon} \quad (2)$$

At the n^{th} stage of its construction there are 2^n intervals, each of length 3^{-n} , and Λ is contained in these intervals. Hence

$$N(\epsilon) = 2^n \quad \text{if} \quad \frac{1}{3^n} \leq \epsilon < \frac{1}{3^{n-1}}. \quad (3)$$

¹After completing this work we learned that Robert MacKay had noted the universal nature of the modulation function for period-doubling in his Ph.D. thesis, though he did not compute the functions [15].

Furthermore

$$N(\epsilon) = 2N(3\epsilon) \quad (4)$$

This functional equation has solutions

$$N(\epsilon) = \phi(\log \epsilon) \epsilon^{-\frac{\log 2}{\log 3}} \quad (5)$$

where

$$\phi(\log \epsilon) = \phi(\log \epsilon + \log 3) \quad (6)$$

which gives $C(\Lambda) = \frac{\log 2}{\log 3}$. Since $N(\epsilon)$ is known explicitly from (3), we can calculate ϕ to obtain the saw-tooth function (for $\epsilon < 1$, i.e. $\log \epsilon < 0$)

$$\log \phi(\log \epsilon) = \frac{\log 2}{\log 3} \log \epsilon + \log 2 \quad 0 > \log \epsilon \geq -\log 3 \quad (7)$$

and $\phi(\log \epsilon - \log 3) = \phi(\log \epsilon)$. This function reflects two aspects of the Cantor set. First, the self-similarity of the set is expressed in the periodicity of ϕ , and second, the non-uniformity or bunched aspect of the set is expressed by deviations of ϕ from a constant (hence the term lacunarity, referring to the holes or non-uniformity of the set). Numerically computed estimates of $\log N(\epsilon)$ against $-\log \epsilon$ show periodic variations from a straight line of slope $\frac{\log 2}{\log 3}$ and any attempt to measure the capacity of a set must take this into account. Plots of the lacunarity oscillations of related scaling functions for a number of different Cantor sets are given in [19]. Note that the modulation function, ϕ , of (7) is *not* what we would find in the standard numerical computations of ϕ . $N(\epsilon)$ is the smallest number of ϵ -balls needed to cover Λ , whilst numerical computations do not, in general, compute this minimal cover (see [11], and references thereof).

Similar effects arise in the fractal sets which arise in deterministic dynamical systems [23]. For example, the scaling behaviour of strange attractors near hyperbolic periodic points can be studied in this way (examples from the Hénon map [14] are given [18] and [22]). In this case the stable eigenvalue of the periodic point determines the period of oscillation in the (local) lacunarity.

2. Universality and Period-doubling

The work mentioned above concentrates on the spatial structure of complex sets. It is equally possible to find systems with complex dependence

upon parameters and it is this side of the problem that we wish to comment upon. To do this we shall concentrate on scaling behaviour associated with cascades of period-doubling bifurcations, although at the end of this note we suggest some further situations where the idea of parametric lacunarity may be useful.




Figure 1: The Lyapunov exponent for the logistic map: (a) base 2 logarithm of the Lyapunov exponent against logarithm $r - r_\infty$. The line reflects the power law $(r - r_\infty)^{\frac{1}{\log_2(\delta)}}$. (b) modulation function $\psi(\log_2(r - r_\infty))$ obtained by factoring out the power law.

Period-doubling cascades in unimodal (or one-hump) maps such as the logistic map, $f_\mu(x) = \mu x(1 - x)$, have been studied extensively, both from a topological and a functional point of view [2, 3, 7]. It has been observed (and understood theoretically) that the envelope of the Lyapunov exponents above the point of accumulation of period-doubling in a family of unimodal maps, $\{f_\mu\}$, scales like

$$|\mu - \mu_\infty|^{\frac{\log 2}{\log \delta}} \quad (8)$$

as shown in Figure 1a, where μ_∞ is the accumulation value and δ is a constant (the Feigenbaum constant [8], depending only upon the universality class of $\{f_\mu\}$). In Figure 1b, the underlying power law behaviour of the Lyapunov exponents has been factored out of the signal. The resulting curve certainly looks periodic, and it is this feature of the problem that will be explained below.

The parameter dependence of Lyapunov exponents is not continuous, so we will concentrate on the scaling of another measure of complexity, the topological entropy, since this varies continuously for C^1 families of unimodal maps (see e.g. [7] chapter II.9, for technical details – the topological entropy for unimodal maps is essentially the asymptotic growth rate (limsup) of the number of periodic points of period p as $p \rightarrow \infty$). A similar argument to that presented below should hold (with certain extra assumptions) for the Lyapunov exponents except that the modulating functions cannot be continuous.




Figure 2: The function f and f^2 showing the geometry of renormalization and the map $\mathcal{T}f$.

To understand this scaling, and to take it a little further, we need to recall some basic facts about period-doubling cascades. In what follows we assume that the family $\{f_\mu\}$ has a quadratic maximum for each μ and that there exists a parameter value μ_∞ for which f_{μ_∞} is at the accumulation of period-doubling. It is also convenient to choose a normalization condition: we assume that the maximum is at $x = 0$ and that $f_\mu(0) = 1$ for all μ . Furthermore, we assume that the maps are even ($f_\mu(-x) = f_\mu(x)$), so the maps are defined on the interval $[-1, 1]$ as shown in Figure 2. The analysis of the period-doubling cascades of unimodal maps uses the idea of renormalization [5, 8, 3, 7]: if $f_\mu(1) = -\alpha^{-1} < 0$ then define an induced map on the interval $[-\alpha^{-1}, \alpha^{-1}]$ by taking the second iterate of f in that interval. This new map

is again a unimodal map, and maps the interval $[-\alpha^{-1}, \alpha^{-1}]$ into itself provided the map is sufficiently close to the accumulation of period-doubling. This new map can be rescaled to give a unimodal map of $[-1, 1]$, $\mathcal{T}f$, defined by

$$\mathcal{T}f(x) = -\alpha f \circ f(-x/\alpha). \quad (9)$$

The operator \mathcal{T} can now be thought of as a map on a suitably defined function space (see [7], for full details). In the universality class we are considering, \mathcal{T} has a fixed point $f_*(x)$,

$$\mathcal{T}f_* = f_*,$$

with a one-dimensional unstable manifold, S , with associated eigenvalue $\delta > 1$ and a codimension one stable manifold Σ as shown in Figure 3. The fixed point, f_* also fixes a value of α : $-\alpha_*^{-1} = f_*(1)$. For unimodal maps with quadratic maxima $\delta \approx 4.6692$ and $\alpha_* \approx 0.3995$.




Figure 3: The structure of function space near the fixed point f_* of \mathcal{T} , after [4].

The standard argument for the universality of period-doubling in quadratic families assumes that there exists a codimension one manifold, Σ_n , beneath Σ and intersecting S transversely, such that if $f \in \Sigma_n$ then f has a non-hyperbolic periodic orbit of period 2^{n-1} which satisfies the conditions for a period-doubling bifurcation. Hence, just below Σ_n maps have a stable periodic orbit of period 2^{n-1} , and just above Σ_n this orbit has lost stability

and there is a stable periodic orbit of period 2^n . Now consider maps in the manifold $\mathcal{T}^{-1}\Sigma_n = \Sigma_{n+1}$, which lies between Σ_n and Σ (distances having been contracted by a factor δ^{-1} in the direction of S and expanded in the other directions). If $\mathcal{T}f$ period-doubles from period 2^{n-1} to 2^n then f period-doubles from period 2^n to 2^{n+1} , so Σ_{n+1} consists of maps with non-hyperbolic periodic orbits of period 2^n , about to period-double. Repeated application of \mathcal{T}^{-1} produces a sequence of manifolds accumulating on Σ at the rate δ^{-1} . By a similar argument, there is a sequence of surfaces $\tilde{\Sigma}_n$ accumulating on Σ from above at the same rate, such that if $f \in \tilde{\Sigma}_n$ then f^{2^n} restricted to a suitable interval is a unimodal map which maps the interval onto itself twice (as is the case for $x \rightarrow 4x(1-x)$), and hence f has topological entropy $2^{-n} \log 2$ [3].

Now consider a family of maps, $\{f_\mu\}$, which intersects Σ transversely at $\mu = 0$, with period-doubling cascades occurring as μ tends to zero from below. This family is represented by a curve in function space as shown in Figure 3, and in terms of the description of the previous paragraph, the sequence of period-doubling bifurcations occurs at the parameter values $\mu_n < 0$ on which f_{μ_n} intersects Σ_n , and $\mu_n \rightarrow 0$ as $n \rightarrow \infty$. Similarly there is a sequence of parameters $\tilde{\mu}_n > 0$, with $\tilde{\mu}_n \rightarrow 0$ as $n \rightarrow \infty$ such that $f_{\tilde{\mu}_n}$ intersects $\tilde{\Sigma}_n$. Assuming that μ is a reasonable measure of distance between maps in function space, this immediately gives the famous result

$$\lim_{n \rightarrow \infty} \frac{\mu_{n-1} - \mu_n}{\mu_n - \mu_{n+1}} = \lim_{n \rightarrow \infty} \frac{\tilde{\mu}_{n-1} - \tilde{\mu}_n}{\tilde{\mu}_n - \tilde{\mu}_{n+1}} = \delta \quad (10)$$

[8, 5]. If $\mu \in (\mu_n, \tilde{\mu}_n)$ then f_μ can be renormalized n times, i.e. $\mathcal{T}^n f_\mu$ is well defined, and the curve of maps $\mathcal{T}^n f_\mu$ has been contracted towards S and stretched in the direction of S . For sufficiently large n the curve representing $\mathcal{T}^n f_\mu$, $\mu_n < \mu < \tilde{\mu}_n$, is thus arbitrarily close to S . Thus, again assuming that μ is a reasonable measure of distance in function space and $|\mu|$ is sufficiently small,

$$\mathcal{T}f_\mu \sim f_{\delta\mu}. \quad (11)$$

But if $h(f)$ is the topological entropy of f then (e.g. [2])

$$h(\mathcal{T}f) = h(f^2) = 2h(f),$$

so, using (11) and writing $h(\mu)$ for $h(f_\mu)$,

$$h(\delta\mu) \sim 2h(\mu). \quad (12)$$

Solving this equation gives

$$h(\mu) \sim \psi(\log \mu) \mu^{\frac{\log 2}{\log \delta}}. \quad (13)$$

where the function $\psi(\log \mu)$ is periodic with period $\log \delta$. We have already noted that the curve in function space given by $\{\mathcal{T}^n f_\mu \mid \mu \in (0, \tilde{\mu}_n)\}$ converges to the part of S (the unstable manifold of f_*) above Σ . Hence as $\mu \downarrow 0$ we can expect the modulation ψ in (13) to tend to the modulation on S . In this sense, then, the function ψ in (13) is asymptotically universal within the universality class of the renormalization operator (so a map with a different order maximum would have a different asymptotic modulation and, of course, a different δ).




Figure 4: The topological entropy for the logistic map: (a) logarithm of the topological entropy against logarithm of the parameter showing power law behaviour; (b) modulation function obtained by factoring out the power law.

The asymptotically universal function describes the fine structure of the entropy between $\tilde{\mu}_{n+1}$ and $\tilde{\mu}_n$; since we know that there are intervals of μ values in this range on which the entropy is constant, $\psi(\log \mu)$ is not constant and reflects the variation of $h(\mu)$ from the simple power law $\mu^{\frac{\log 2}{\log \delta}}$.

As mentioned earlier, the entropy, $h(f)$, is a continuous function of parameters whereas the Lyapunov exponent is not. However, the Lyapunov exponent is easier to calculate directly from iterates of the map numerically. To find the topological entropy of a map (and hence to compute ψ) we have

used a result due to Milnor and Thurston [16]. If the critical point of the map is at $x = c$, then we define $a_0(x) = 1$ if $x \leq c$ and $a_0(x) = -1$ if $x > c$. Now define $a_i(x)$ inductively by the relation $a_i(x) = a_{i-1}(x)a_0(f^i(x))$, $i > 0$. The kneading sequence (or itinerary) of a point is the sequence $I(x) = a_0(x)a_1(x)a_2(x)\dots$ and the kneading invariant of a map, f , is the sequence

$$k(f) = \lim_{x \uparrow c} I(x). \quad (14)$$

Now, if $k(f) = k_0k_1k_2\dots$, then we can form the formal power series

$$K(t) = \sum_{i=0}^{\infty} k_i t^i. \quad (15)$$

Milnor and Thurston [16] prove that if $h(f) > 0$ and s is the smallest positive zero of $K(t)$, i.e. $K(s) = 0$ and $K(t) > 0$ for all $0 \leq t < s$, then

$$h(f) = \log(s^{-1}). \quad (16)$$

The first symbols of the kneading invariant of a map are easy to calculate numerically, and we then solved the polynomial approximation of the power series $K(t)$ to find an approximate value of the topological entropy of f . The results are shown in Figure 4. Figure 4a shows a graph of the entropy of f_μ against μ for the logistic map, and Figure 4b shows the (asymptotically universal) function $\psi(\log \mu)$ over three periods. Figure 1b shows the corresponding function for the Lyapunov exponents of the map over four periods (that is, the horizontal extent of the graph corresponds to $4\log_2(\delta)$). We consider this figure to be strong evidence that a similar argument holds for the Lyapunov exponents, or at least that it holds to a good approximation (for the argument to work for the Lyapunov exponents, (11) must hold with differentiable conjugacy; we do not know whether this is the case).

Of course, this same universal function should exist near the accumulation of period-doubling in other types of dynamical systems, in differential equations for example. To illustrate this we have calculated the largest Lyapunov exponent for the Lorenz equations near a window of stable periodic orbits identified by Sparrow [21]. The Lorenz equations are

$$\dot{x} = \sigma(y - x), \quad \dot{y} = rx - y - xz, \quad \dot{z} = -bz + xy \quad (17)$$

where we take the standard parameter values $\sigma = 10$ and $b = \frac{8}{3}$ and treat r as the bifurcation parameter. The system shows a series of period-doubling

bifurcations as r decreases through $r_\infty \approx 99.5247$. Figure 5 shows the results for the Lorenz equations which parallel those of the logistic map shown in Figure 1. Although the estimated² Lyapunov exponents for the Lorenz equations are more uncertain than those of the logistic map, one can still discern the similarity of the oscillation in Figures 1(b) and 5(b).




Figure 5: The leading Lyapunov exponent, Λ_1 , for the Lorenz equations near $r_\infty \approx 99.5247$: (a) logarithm of the Lyapunov exponent against logarithm of $|r_\infty - r|$; (b) modulation function obtained by removing the power law $(r_\infty - r)^{\frac{1}{\log_2(\delta)}}$. There are no free parameters in extracting this oscillation. Note the similarity to the result of Figure 1(a).

3. The period-doubling Cantor set

We began this note by discussing lacunarity in the context of spatial structure and have ended with an approximate function associated with intermittency. At the accumulation of period-doubling, a unimodal map has an invariant Cantor set on which the motion is not chaotic. Once again, there is an approximately self-similar structure here.

The invariant Cantor set at the accumulation of period-doubling is not self-similar. As Figure 2 suggests, the Cantor set can be constructed by

²These numerical estimates were obtained from using the ϵ -returns algorithm. The relative uncertainty in the estimate increases (visibly) for smaller $r_\infty - r$, as the value of $\Lambda(r) \rightarrow 0$.

successively dividing an interval into two smaller intervals of different length. One of these is a factor of α times the parent interval, whilst the other (the right hand invariant interval in Figure 2) can be calculated approximately using a quadratic approximation to f_* to be roughly a factor of α^2 times smaller than the parent interval. So each interval at the n^{th} generation splits into two intervals whose lengths are respectively α and α^2 times the length of the parent interval. From this observation it is easy to see that at the n^{th} generation there are 2^n intervals of which $\binom{n}{k}$ are of length $\alpha^k \alpha^{2n-2k}$.

Thus we have

$$N(\alpha^n) = N(\alpha^{n-1}) + N(\alpha^{n-2}), \quad (18)$$

(cf. $F_n = F_{n-1} + F_{n-2}$, which generates the Fibonacci numbers), and so we obtain the approximation

$$N(\alpha_*^2 \epsilon) = N(\alpha_* \epsilon) + N(\epsilon), \quad (19)$$

where α has been replaced by the asymptotic value α_* defined above.

It is now easy to check that if

$$d_1 = \frac{\log\left(\frac{\sqrt{5}+1}{2}\right)}{\log \alpha_*} \quad \text{and} \quad d_2 = \frac{\log\left(\frac{\sqrt{5}-1}{2}\right)}{\log \alpha_*}$$

and $\phi_i(\log x)$, $i = 1, 2$, are arbitrary periodic functions with period $2 \log \alpha_*$ then

$$\phi_1(\log \epsilon) \epsilon^{d_1} \cos \frac{2m\pi \log \epsilon}{\log \alpha_*} \quad \text{and} \quad \phi_2(\log \epsilon) \epsilon^{d_2} \cos \frac{(2m+1)\pi \log \epsilon}{\log \alpha_*} \quad (20)$$

are solutions, for each $m \in \mathbf{Z}$. Noting that $d_1 < 0$ and $d_2 > 0$ (as $\alpha_* < 1$), and that the cosines are also periodic with period $2 \log \alpha_*$ we find that

$$N(\epsilon) \sim \chi(\log \epsilon) \epsilon^{d_1} \quad (21)$$

as ϵ tends to zero, where $\chi(\log \epsilon) = \chi(\log \epsilon + 2 \log \alpha_*)$. This is only an approximate result: the invariant Cantor set is not strictly self-similar in the way described and the recursion relation for $N(\epsilon)$ is also an approximation. For a fuller, and more precise treatment, see [9, 10].

4. Conclusion

We have given four examples of lacunarity: two for sets in state space and two for sets in parameter space. For the middle third Cantor set the analysis is exact and the modulation function can be derived explicitly. In the second example, for the scaling of topological entropy, the modulation function was computed numerically, and in this case the theory is asymptotically exact, in the sense that the modulation function is universal sufficiently close to the accumulation of period-doubling. The third example, for the Lyapunov exponents above the accumulation of period-doubling, mirrors the analysis of the topological entropy except the function is much less regular (it is not continuous). The analysis of the final example is approximate. Here we defined a scaling for the Cantor set at the accumulation of period-doubling which is asymptotically exact at small length scales, but we have used a rough approximation of this scaling law to obtain approximate scaling functions.

There are other situations in which it is possible to obtain approximate self-similarity over some set in parameter space or state space. In these cases it is possible to repeat the argument used above, but the periodic modulation functions will only be approximately valid. The most obvious example of approximate self-similarity in parameter space is intermittency. The standard analysis (e.g. [13]) uses an approximate renormalization scheme to describe the length of time spent close to the location of a saddlenode bifurcation, and this can be used to derive scaling and *approximate* modulation functions. We have not considered this case in detail, but it would be interesting to know how much relevant information these approximations hold.

Acknowledgements: The work on the logistic map was done at D.A.M.T.P., University of Cambridge, in 1991 supported by a research grant from SERC (now renamed as EPSRC). We are grateful to P. McSharry for computing the Lyapunov exponents for the Lorenz system in finite time.

References

- [1] Badii, R. and A. Politi (1984) Hausdorff Dimension and Uniformity Factor of Strange Attractors, *Phys. Rev. Lett.* **52** 1661–1664.
- [2] Block, L.S. and W.A. Coppel (1991) *Dynamics in One Dimension*, LNM 1513, Springer, Berlin.

- [3] Collet, P. and J.P. Eckmann (1980) *Iterated maps on the interval as dynamical systems*, Birkhauser, Boston.
- [4] Collet, P., J.P. Eckmann and O.E. Lanford (1980), Universal properties of maps on an interval, *Comm. Math. Phys.* **76** 211–254.
- [5] Couillet, P. and C. Tresser (1978) Iterations d’endomorphismes et groupe de renormalisation, *J. de Physique C5*, **9** C5.25–C5.27.
- [6] De Bruijn, N.G. (1948). On Mahler’s partition problem, *Proc. Kon. Ned. Akad. van Wetenschappen*, **51**, 659–669.
- [7] de Melo, W. and S. van Strien (1993) *One-dimensional dynamics*, Springer, Berlin.
- [8] Feigenbaum, M.J. (1978) Quantitative universality for a class of nonlinear transformations, *J. Stat. Phys.* **19** 25–52.
- [9] Feigenbaum, M.J. (1987) Some characterizations of strange sets, *J. Stat. Phys.* **46** 919–924.
- [10] Feigenbaum, M.J. (1988) Presentation functions, fixed-points, and a theory of scaling function dynamics, *J. Stat. Phys.* **52** 527–569.
- [11] Grassberger, P. and I. Procaccia, (1983) Characterization of Strange Attractors, *Phys. Rev. Lett.* **50** 346–349.
- [12] Guckenheimer, J. (1984) Dimension estimates for attractors, in *Contemp. Math.* **28**, *Fluids and Plasmas: Geometry and Dynamics*, ed. J.E. Marsden, 357–367.
- [13] Guckenheimer, J. and P. Holmes (1983) *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*, Springer, New York.
- [14] Hénon, M., (1976) A Two-Dimensional Mapping with a Strange Attractor, *Commun. Math. Phys.* **50** 69–77.
- [15] MacKay, R.S. (1982) *Renormalization in area-preserving maps*, Ph.D. thesis, Princeton.
- [16] Milnor, J. and W. Thurston (1988) On iterated maps of the interval, in *Dynamical Systems* ed. J.C. Alexander, Springer LNM 1342.

- [17] Mullhaupt, A. P. (1987) Delay Equations, DeBruijn Oscillations, and Intermittency, *Phys. Lett. A*, **124** 151–158.
- [18] Smith, Leonard A. (1987) Lacunarity and Chaos in Nature, PhD Thesis, Columbia University, New York.
- [19] Smith, L. A., J.-D. Fournier and E. A Spiegel (1986) Lacunarity and intermittency in fluid turbulence. *Phys. Lett. A* **114** 465–468.
- [20] Smith, Leonard A. (1997) *The Maintenance of Uncertainty*, In Proceedings of the International School of Physics “Enrico Fermi”, Course **CXXXIII**, pages 177–246, Italian Physical Society, Bologna.
- [21] Sparrow, C.T. (1982) *The Lorenz equations: bifurcations, chaos and strange attractors*, Applied Math. Sci. 41, Springer, New York.
- [22] Theiler, J. (1987) *Quantifying chaos*, PhD thesis, California Institute of Technology, Pasadena, USA.
- [23] Theiler, J. (1989) Lacunarity in a best estimator of fractal dimension, *Phys. Lett. A* **133** 195–200.
- [24] Zhou, W.-X. and D. Sornette (2009) Numerical Investigations of discrete scale invariance in fractals and multifractal measures, *Physica A*, **388** 2623–2639.