

*On the local Langlands correspondence for
non-tempered representations*

Aubert, Anne-Marie and Baum, Paul and Plymen,
Roger and Solleveld, Maarten

2013

MIMS EPrint: **2013.77**

Manchester Institute for Mathematical Sciences
School of Mathematics

The University of Manchester

Reports available from: <http://eprints.maths.manchester.ac.uk/>

And by contacting: The MIMS Secretary
School of Mathematics
The University of Manchester
Manchester, M13 9PL, UK

ISSN 1749-9097

On the local Langlands correspondence for non-tempered representations

Anne-Marie Aubert, Paul Baum, Roger Plymen,
and Maarten Solleveld

Dedicated to Peter Schneider on the occasion of his 60th birthday

Abstract. Let G be a reductive p -adic group. We study how a local Langlands correspondence for irreducible tempered G -representations can be extended to a local Langlands correspondence for all irreducible smooth representations of G . We prove that, under a natural condition involving compatibility with unramified twists, this is possible in a canonical way.

To this end we introduce analytic R-groups associated to non-tempered essentially square-integrable representations of Levi subgroups of G . We establish the basic properties of these new R-groups, which generalize Knapp–Stein R-groups.

CONTENTS

Introduction	999
1. Analytic R-groups for non-tempered representations	1001
2. Algebraic families of irreducible representations	1007
3. Algebraic families of Langlands parameters	1009
4. From a tempered to a general local Langlands correspondence	1012
5. Geometric R-groups	1016
6. The principal series of a split group	1018
References	1020

INTRODUCTION

Let F be a local nonarchimedean field and let G be the group of F -rational points of a connected reductive group which is defined over F . Let $\text{Irr}(G)$ be the space of irreducible smooth G -representations and let $\Phi(G)$ be the space of conjugacy classes of Langlands parameters for G . The local Langlands

correspondence (LLC) conjectures that there exists an explicit map

$$\mathrm{Irr}(G) \rightarrow \Phi(G)$$

which satisfies several naturality properties [6]. The collection of representations that correspond to a fixed $\phi \in \Phi(G)$ is known as the L-packet $\Pi_\phi(G)$ and should be finite. A more subtle version of the LLC [31, 3], which for unipotent representations stems from [21], asserts that the members of $\Pi_\phi(G)$ can be parametrized by some irreducible representations ρ of a finite group \mathcal{S}_ϕ . This leads to a space $\Phi^e(G)$ of enhanced Langlands parameters (ϕ, ρ) , and the LLC then should become an injection

$$\mathrm{Irr}(G) \rightarrow \Phi^e(G).$$

The proofs of the LLC for $\mathrm{GL}_n(F)$ [20, 14, 16] are major results. Together with the Jacquet–Langlands correspondence these provide the LLC for inner forms of $\mathrm{GL}_n(F)$, see [17, 5]. (This has been known for a long time already, but was apparently not published earlier.) Recently there has been considerable progress on the LLC for inner forms of $\mathrm{SL}_n(F)$ [17] and for quasi-split classical groups [4, 23]. The LLC has been established for a large class of representations of these groups, including the collection $\mathrm{Irr}^t(G)$ of irreducible tempered representations.

In general it is expected that is easier to prove the LLC for tempered representations of a p -adic group G than for all irreducible representations. The main reason is that every irreducible tempered G -representation is unitary and appears as a direct summand of the parabolic induction of some essentially square-integrable representation.

Therefore a method to generalize the LLC from $\mathrm{Irr}^t(G)$ to $\mathrm{Irr}(G)$ is useful. The aim of this paper is to provide such a method, which is simple in comparison with the aforementioned papers. The idea is based on the Langlands classification and to some extent already present in [10, 3, 28]. It applies to all reductive groups over local non-archimedean fields. Recall that a part of Langlands’ conjectures is that $\mathrm{Irr}^t(G)$ corresponds to the set $\Phi_{\mathrm{bdd}}(G)$ of bounded Langlands parameters (modulo conjugacy).

Theorem 4.2. *Suppose that a tempered local Langlands correspondence is given as an injective map $\mathrm{Irr}^t(G) \rightarrow \Phi_{\mathrm{bdd}}^e(G)$, which is compatible with twisting by unramified characters whenever this is well-defined. Then the map extends canonically to a local Langlands correspondence $\mathrm{Irr}(G) \rightarrow \Phi^e(G)$.*

The main novelty of the paper is the introduction of analytic R-groups for non-tempered representations (see Definition 1.4 and Theorem 1.6). These objects, natural generalizations of R-groups defined (in the p -adic case) by Silberger [29], open up new ways to compare $\mathrm{Irr}^t(G)$ with $\mathrm{Irr}(G)$. Roughly speaking, $\mathrm{Irr}(G)$ is obtained from $\mathrm{Irr}^t(G)$ by ”complexification” (Proposition 2.1).

We show that the relation between $\Phi_{\text{bdd}}^e(G)$ and $\Phi^e(G)$ is similar (Proposition 3.2). As these spaces are not algebraic varieties, a large part of the proof consists of making the term "complexification" precise in this context. We do this by constructing suitable algebraic families of irreducible representations and of enhanced Langlands parameters.

In Section 5 we conjecture how our analytic R-groups are related to geometric R-groups. This should enable one to produce a LLC for $\text{Irr}(G)$ if the Langlands parameters corresponding to essentially square-integrable representations of Levi subgroups of G are known.

With this in mind we check that the hypotheses of Theorem 4.2 are fulfilled in some known cases, in particular for the principal series of a split reductive p -adic group.

1. ANALYTIC R-GROUPS FOR NON-TEMPERED REPRESENTATIONS

Let F be a local nonarchimedean field and let \mathcal{G} be a connected reductive algebraic group defined over F . We consider the group $G = \mathcal{G}(F)$ of F -rational points. Let P be a parabolic subgroup of G with Levi factor M , and let A be the maximal F -split torus in the centre of M . Then $M = Z_G(A)$ and $N_G(M) = N_G(A)$. The Weyl group of M and A is

$$W(M) = W(A) = N_G(M)/M = N_G(A)/M.$$

It acts on equivalence classes of M -representations by

$$(1) \quad (w \cdot \pi)(m) = (\bar{w} \cdot \pi)(m) = \pi(\bar{w}^{-1}m\bar{w}),$$

for any representative $\bar{w} \in N_G(M)$ of $w \in W(M)$. The isotropy group of π is

$$W_\pi := \{w \in W(M) : w \cdot \pi \cong \pi\}.$$

Let M^1 be the subgroup of M generated by all compact subgroups of M . Then M/M^1 is a lattice and a character of M is unramified if and only if it factors through M/M^1 . Let $X_{\text{nr}}(M)$ be the group of unramified characters of M and let $X_{\text{unr}}(M)$ be the subgroup of unitary unramified characters. The above provides $X_{\text{nr}}(M)$ with the structure of a complex torus, such that $X_{\text{unr}}(M)$ is its maximal compact subgroup.

In this paper all representations of p -adic groups are tacitly assumed to be smooth. Let I_P^G be the functor of smooth, normalized parabolic induction, from M -representations to G -representations. The following result is well-known, we include the proof for a lack of a good reference.

Lemma 1.1. *Let π be a finite length M -representation and take $w \in W(M)$. Let $P' \subset G$ be another parabolic subgroup with Levi factor M . Then the G -representations $I_P^G(\pi)$, $I_{P'}^G(w \cdot \pi)$ and $I_{P'}^G(\pi)$ have the same trace and the same irreducible constituents, counted with multiplicity.*

Proof. Conjugation with a representative $\bar{w} \in N_G(M)$ for w yields an isomorphism $I_P^G(w \cdot \pi) \cong I_{\bar{w}^{-1}P\bar{w}}^G(\pi)$. The parabolic subgroup $w^{-1}Pw \subset G$ has

$M = w^{-1}Mw$ as a Levi factor, so without loss of generality we may assume that it equals P' .

Since $I_P^G(\pi)$ and $I_{P'}^G(\pi)$ have finite length [11, 6.3.8] their irreducible constituents (and multiplicities) are determined by their traces [11, 2.3.3]. Therefore it suffices to show that the function

$$\begin{aligned} C_c^\infty(G) \times X_{\text{nr}}(M) &\rightarrow \mathbb{C}, \\ (f, \chi) &\mapsto \text{tr}(f, I_P^G(\pi \otimes \chi)) - \text{tr}(f, I_{P'}^G(\pi \otimes \chi)) \end{aligned}$$

is identically zero. For a fixed $f \in C_c^\infty(G)$ this is a rational function on $X_{\text{nr}}(M)$, which by [32, Théorème IV.1.1] vanishes on a Zariski-dense subset of $X_{\text{nr}}(M)$. Hence it vanishes everywhere. \square

Let $X^*(A)$ and $X_*(A)$ be the character (respectively cocharacter) lattice of A . Since $A/(A \cap M^1) \cong X_*(A)$ is of finite index in M/M^1 , the restriction map $X_{\text{nr}}(M) \rightarrow X_{\text{nr}}(A)$ is surjective and has finite kernel. In particular there are natural isomorphisms

$$\{\chi \in X_{\text{nr}}(M) : \chi(M) \subset \mathbb{R}_{>0}\} \xrightarrow{\text{res}} \text{Hom}_{\mathbb{Z}}(X_*(A), \mathbb{R}_{>0}) \xrightarrow{\log} X^*(A) \otimes_{\mathbb{Z}} \mathbb{R} := \mathfrak{a}^*.$$

We note that \mathfrak{a}^* is a real vector space containing the root system $R(G, A)$. We say that $\chi \in X_{\text{nr}}(M)$ is positive with respect to P if

$$(2) \quad \langle \alpha^\vee, \log |\chi| \rangle \geq 0 \text{ for all } \alpha \in R(P, A).$$

Let $M(\chi)$ be the maximal Levi subgroup of G such that

- $M(\chi) \supset M$ and the split part of $Z(M(\chi))$ is contained in A ;
- χ is unitary on $M \cap M(\chi)_{\text{der}}$.

Assume that π is irreducible and tempered. In particular it is unitary. Then $I_{M(\chi) \cap P}^{M(\chi)}(\pi \otimes \chi)$ is completely reducible, because its restriction to $M(\chi)_{\text{der}}$ is unitary. For every irreducible summand τ of $I_{M(\chi) \cap P}^{M(\chi)}(\pi \otimes \chi)$ the pair $(PM(\chi), \tau)$ satisfies the hypothesis of the Langlands classification [9, 18], so $I_{PM(\chi)}^G(\tau)$ is indecomposable and has a unique irreducible quotient $L(PM(\chi), \tau)$. We call the $L(PM(\chi), \tau)$, for all eligible τ , the Langlands quotients of $I_P^G(\pi \otimes \chi)$. This subset of $\text{Irr}(G)$ depends only $(M, \pi \otimes \chi)$, because $M(\chi)$ and $PM(\chi)$ are uniquely determined by $\log |\chi|$. We denote it by $\text{Irr}_{M, \pi \otimes \chi}(G)$.

In fact $I_P^G(\pi \otimes \chi)$ is completely reducible for χ in a Zariski-dense subset of $X_{\text{nr}}(M)$. In that case $\text{Irr}_{M, \pi \otimes \chi}(G)$ consists of all the constituents of $I_P^G(\pi \otimes \chi)$.

The uniqueness part of the Langlands classification tells us that $L(PM(\chi), \tau)$ is tempered if and only if $M(\chi) = G$ and τ is tempered. This is so if and only if χ is unitary, in which case actually all members of $\text{Irr}_{M, \pi \otimes \chi}(G)$ are tempered.

By Lemma 1.1 the elements of $\text{Irr}_{M, \pi \otimes \chi}(G)$ are also constituents of $I_{P'}^G(\pi \otimes \chi)$, so it is justified to call them the Langlands constituents of $I_{P'}^G(\pi \otimes \chi)$ for any parabolic subgroup $P' \subset G$ containing M .

Harish-Chandra showed that every irreducible tempered representation can be obtained as a direct summand of the parabolic induction of a square-integrable (modulo centre) representation, in an essentially unique way [32, Proposition III.4.1]. These considerations lead to the following result.

Theorem 1.2. [28, Theorem 2.15]

- (a) For every $\pi \in \text{Irr}(G)$ there exist P, M, χ as above and a square-integrable (modulo centre) representation $\omega \in \text{Irr}(M)$, such that $\pi \in \text{Irr}_{M, \omega \otimes \chi}(G)$.
- (b) The groups P and M are unique up to conjugation. Once they are chosen, $\omega \otimes \chi$ is uniquely determined up to $W(M)$ -equivalence.
- (c) π is tempered if and only if χ is unitary.

Thus $\text{Irr}(G)$ is partitioned in disjoint packets $\text{Irr}_{M, \omega \otimes \chi}(G)$, parametrized by conjugacy classes of Levi subgroups M and $W(M)$ -equivalence classes of essentially square-integrable representations $\omega \otimes \chi \in \text{Irr}(M)$.

We remark that Theorem 1.2 is stronger than the Langlands classification as formulated in [9, IV.2] and [18]. There the passage is from smooth representations to tempered representations, whereas in Theorem 1.2 the passage is from smooth representations to essentially square-integrable representations (all assumed irreducible of course). On the other hand, the Langlands classification is one-to-one but Theorem 1.2 is only finite-to-one.

Let $\omega \in \text{Irr}(M)$ be square-integrable modulo centre and write

$$(3) \quad \begin{aligned} \mathcal{O} &= \{\omega \otimes \chi \in \text{Irr}(M) : \chi \in X_{\text{unr}}(M)\}, \\ \mathcal{O}_{\mathbb{C}} &= \{\omega \otimes \chi \in \text{Irr}(M) : \chi \in X_{\text{nr}}(M)\}. \end{aligned}$$

The irreducible constituents of the G -representations $I_P^G(\omega \otimes \chi)$ with $\omega \otimes \chi \in \mathcal{O}$ make up a Harish-Chandra component $\text{Irr}_{\mathcal{O}}(G)$ of $\text{Irr}^t(G)$, see [27, §1]. The group

$$X_{\text{nr}}(M)_{\omega} := \{\chi \in X_{\text{nr}}(M) : \omega \otimes \chi \cong \omega\}$$

is finite, and in particular consists of unitary characters. The bijection

$$(4) \quad X_{\text{nr}}(M)/X_{\text{nr}}(M)_{\omega} \rightarrow \mathcal{O}_{\mathbb{C}} : \chi \mapsto \omega \otimes \chi$$

provides $\mathcal{O}_{\mathbb{C}}$ with the structure of a complex torus, and \mathcal{O} can be identified with its maximal real compact subtorus. However, in general there is no natural multiplication on \mathcal{O} or $\mathcal{O}_{\mathbb{C}}$.

Let $W(\mathcal{O})$ be the stabilizer of \mathcal{O} in $W(M)$, with respect to the action (1). It is also the stabilizer of $\mathcal{O}_{\mathbb{C}}$, and it acts on \mathcal{O} and $\mathcal{O}_{\mathbb{C}}$ by algebraic automorphisms.

Recall from [2, §2] that for every $w \in W(M)$ and every $\omega \otimes \chi \in \mathcal{O}$ there exists a unitary intertwining operator

$$(5) \quad J(w, \omega \otimes \chi) \in \text{Hom}_G(I_P^G(\omega \otimes \chi), I_P^G(w \cdot (\omega \otimes \chi))).$$

It is unique up to a complex number of norm 1. If χ_G is the restriction to M of an unramified character of G , then the right hand side of (5) is $\text{Hom}_G(I_P^G(\omega), I_P^G(w \cdot (\omega)))$, so then we may take

$$(6) \quad J(w, \omega \otimes \chi_G) = J(w, \omega).$$

These operators can be normalized so that $\chi \mapsto J(w, \omega \otimes \chi)$ extends to a rational function on $\mathcal{O}_{\mathbb{C}}$. We fix such a normalization. It determines a rational function $\kappa : W(M) \times W(M) \times \mathcal{O}_{\mathbb{C}} \rightarrow \mathbb{C} \cup \{\infty\}$ by

$$(7) \quad J(w, w' \cdot (\omega \otimes \chi)) \circ J(w', \omega \otimes \chi) = \kappa(w, w', \omega \otimes \chi) J(w w', \omega \otimes \chi).$$

On \mathcal{O} this function is regular and takes values of norm 1. By (7) this holds more generally for all twists of ω by unramified characters of M which are unitary on $M \cap G_{\text{der}}$. We let

$$\kappa_{\omega \otimes \chi} : W_{\omega \otimes \chi} \times W_{\omega \otimes \chi} \rightarrow \mathbb{C} \cup \{\infty\}$$

be the restriction of κ to $W_{\omega \otimes \chi} \times W_{\omega \otimes \chi} \times \{\omega \otimes \chi\}$.

Lemma 1.3. $\kappa_{\omega \otimes \chi}$ has neither poles nor zeros.

Proof. Let $w \in W_{\omega \otimes \chi}$ and recall the definition of $M(\chi)$, below (2). It shows that $w \in N_{M(\chi)}(M)/M$. As we saw above, the operator

$$J_{M(\chi)}(w, \omega \otimes \chi) \in \text{End}_{M(\chi)}(I_{P \cap M(\chi)}^{M(\chi)}(\omega \otimes \chi))$$

is regular and invertible. Hence $I_{P \cap M(\chi)}^G(J_{M(\chi)}(w, \omega \otimes \chi))$ is invertible as well. But all the positive roots of $R(G, A)$ that are made negative by w belong to $R(M(\chi), A)$, so

$$J(w, \omega \otimes \chi) = z I_{P \cap M(\chi)}^G(J_{M(\chi)}(w, \omega \otimes \chi)) \text{ for some } z \in \mathbb{C}^{\times}.$$

Now (7) shows that $\kappa_{\omega \otimes \chi}(w, w') \in \mathbb{C}^{\times}$ for all $w, w' \in W_{\omega \otimes \chi}$. □

The associativity of the multiplication in $\text{End}_G(I_P^G(\omega \otimes \chi))$ implies that $\kappa_{\omega \otimes \chi}$ is a 2-cocycle of $W_{\omega \otimes \chi}$. It gives rise to a twisted group algebra $\mathbb{C}[W_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi}]$. By definition this algebra has a basis $\{J_w : w \in W_{\omega \otimes \chi}\}$ and its multiplication is given by

$$(8) \quad J_w \cdot J_{w'} = \kappa_{\omega \otimes \chi}(w, w') J_{w w'}.$$

Let $R_{\text{red}}(G, A)$ be the reduced root system of (G, A) . Harish-Chandra's μ -function determines a subset

$$R_{\omega \otimes \chi} := \pm \{\alpha \in R_{\text{red}}(G, A) : \mu_{\alpha}(\omega \otimes \chi) = 0\},$$

which is known to be a root system itself [29, §1]. Its Weyl group $W(R_{\omega \otimes \chi})$ is a normal subgroup of $W_{\omega \otimes \chi}$. The parabolic subgroup P determines a set of positive roots $R_{\omega \otimes \chi}^+$. Since $W_{\omega \otimes \chi}$ acts on $R_{\omega \otimes \chi}$, it is known from the general theory of Weyl groups that the subgroup

$$(9) \quad \mathfrak{R}_{\omega \otimes \chi} := \{w \in W_{\omega \otimes \chi} : w(R_{\omega \otimes \chi}^+) = R_{\omega \otimes \chi}^+\}$$

satisfies

$$(10) \quad W_{\omega \otimes \chi} = \mathfrak{R}_{\omega \otimes \chi} \ltimes W(R_{\omega \otimes \chi}).$$

Definition 1.4. The group $\mathfrak{R}_{\omega \otimes \chi}$ is the analytic R -group attached to the essentially square-integrable representation $\omega \otimes \chi \in \text{Irr}(M)$.

Lemma 1.5. *Let Y be a connected subset of $\mathcal{O}_{\mathbb{C}}$ such that $W_{\omega \otimes \chi}$ is the same for all $\omega \otimes \chi \in Y$.*

- (a) *$R_{\omega \otimes \chi}$ and $\mathfrak{R}_{\omega \otimes \chi}$ are independent of $\omega \otimes \chi \in Y$, up to a natural isomorphism.*
- (b) *$\mathbb{C}[\mathfrak{R}_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi}]$ and the projective representation of $W_{\omega \otimes \chi}$ on $I_P^G(\omega \otimes \chi)$ are independent of $\omega \otimes \chi \in Y$, up to an isomorphism which is determined by the normalization of the intertwining operators J_w .*

Proof. (a) Since μ_α depends only on the values of the coroot α^\vee on $\mathcal{O}_{\mathbb{C}}$, it is constant on the connected components of $\mathcal{O}_{\mathbb{C}}^{s_\alpha}$. Hence $R_{\omega \otimes \chi} = R_{\omega \otimes \chi'}$ for all $\omega \otimes \chi, \omega \otimes \chi' \in Y$. This implies the corresponding statement for the R-groups, by their very definition.

(b) The action of $W_{\omega \otimes \chi}$ via the $J(w, \omega \otimes \chi)$ defines a projective representation on $I_P^G(\omega \otimes \chi)$. By [32, §IV.1] the vector space underlying $I_P^G(\omega \otimes \chi)$ is independent of $\chi \in X_{\text{nr}}(M)/X_{\text{nr}}(M)_\omega$. By Lemma 1.3 the $J(w, \omega \otimes \chi)$ depend algebraically on χ , so we have a continuous family of projective representations of the finite group $W_{\omega \otimes \chi}$. Given the dimension, there are only finitely many equivalence classes of such representations, so all the $I_P^G(\omega \otimes \chi)$ with $\omega \otimes \chi \in Y$ are isomorphic as projective $W_{\omega \otimes \chi}$ -representations. In particular the 2-cocycles $\kappa_{\omega \otimes \chi}$ of $W_{\omega \otimes \chi}$ for different $\omega \otimes \chi \in Y$ are in the same cohomology class. Moreover, since the $\kappa_{\omega \otimes \chi}$ are defined in terms of the $J(w, \omega \otimes \chi)$, they vary continuously as functions on Y . Now (8) shows that there is a unique family algebra isomorphisms

$\mathbb{C}[W_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi}] \rightarrow \mathbb{C}[W_{\omega \otimes \chi'}, \kappa_{\omega \otimes \chi'}]$ of the form $J_w \mapsto a_w(\omega \otimes \chi, \omega \otimes \chi') J_w$ with $a_w : Y^2 \rightarrow \mathbb{C}^\times$ continuous and $a_w(\omega \otimes \chi, \omega \otimes \chi) = 1$. In view of part (a) these isomorphisms restrict to

$$\mathbb{C}[\mathfrak{R}_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi}] \rightarrow \mathbb{C}[\mathfrak{R}_{\omega \otimes \chi'}, \kappa_{\omega \otimes \chi'}]. \quad \square$$

The following result generalizes the theory of R-groups [2, §2] to non-tempered representations. It also provides an explanation for the failure of some properties of R-groups observed in [8], see Example 5.3.

Theorem 1.6. *Let $\omega \in \text{Irr}(M)$ be square-integrable modulo centre and let $\chi \in X_{\text{nr}}(M)$.*

- (a) *There exists an injective algebra homomorphism*

$$\mathbb{C}[\mathfrak{R}_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi}] \rightarrow \text{End}_G(I_P^G(\omega \otimes \chi)),$$

which is bijective if χ is positive with respect to P . It is canonical up to twisting by characters of $\mathfrak{R}_{\omega \otimes \chi}$.

- (b) *Part (a) determines bijections*

$$\begin{array}{ccccc} \text{Irr}(\mathbb{C}[\mathfrak{R}_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi}]) & \rightarrow & \text{Irr}_{M, \omega \otimes \chi}(M(\chi)) & \rightarrow & \text{Irr}_{M, \omega \otimes \chi}(G) \\ \rho & \mapsto & \pi(M, \omega \otimes \chi, \rho) & \mapsto & L(M, \omega \otimes \chi, \rho), \end{array}$$

where $\pi(M, \omega \otimes \chi, \rho) = \text{Hom}_{\mathbb{C}[\mathfrak{R}_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi}]}(\rho, I_{P \cap M(\chi)}^{M(\chi)}(\omega \otimes \chi))$ and $L(M, \omega \otimes \chi, \rho)$ is the unique Langlands constituent of $I_{P_{M(\chi)}}^G(\pi(M, \omega \otimes \chi, \rho))$.

(c) $I_P^G(\omega \otimes \chi) \cong \bigoplus_{\rho} I_{PM(\chi)}^G(\pi(M, \omega \otimes \chi, \rho)) \otimes \rho$ as $G \times \mathbb{C}[\mathfrak{R}_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi}]$ -representations.

Proof. For $\chi \in X_{\text{unr}}(M)$ this is well-known, see [2, §2]. By (6) it holds more generally for $\chi \in X_{\text{nr}}(M)$ which are unitary on $M \cap G_{\text{der}}$.

(a) Since $W(\mathcal{O})$ acts on $\mathcal{O}_{\mathbb{C}}$ by algebraic automorphisms, we can find a set Y as in Lemma 1.5 which contains both $\omega \otimes \chi$ and some $\omega' \in \mathcal{O}$. By [29] the intertwining operator $J(w, \omega') \in \text{End}_G(I_P^G(\omega'))$ is scalar if and only if $w \in W(R_{\omega'})$, and by the aforementioned result of [2] the operators $J(w, \omega')$ with $w \in \mathfrak{R}_{\omega'}$ span a subalgebra of $\text{End}_G(I_P^G(\omega'))$ isomorphic to $\mathbb{C}[\mathfrak{R}_{\omega'}, \kappa_{\omega'}]$. By Lemma 1.5.b the same holds for all elements of Y , and in particular for $\omega \otimes \chi$. By Harish-Chandra's commuting algebra theorem [30, Theorem 5.5.3.2]

$$(11) \quad \mathbb{C}[\mathfrak{R}_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi}] \cong \text{End}_G(I_P^G(\omega \otimes \chi)) \quad \text{for } \chi \in X_{\text{unr}}(M).$$

Since both sides are invariant under twisting by unramified characters of G , (11) holds whenever $\chi \in X_{\text{nr}}(M)$ is unitary on $M \cap G_{\text{der}}$.

Every element of $W(G)$ that stabilizes $(M, \omega \otimes \chi)$ already lies in $W(M(\chi))$. Therefore it does not matter whether we compute $W_{\omega \otimes \chi}$ in G or in $M(\chi)$. The definitions of $R_{\omega \otimes \chi}^+$, $W(R_{\omega \otimes \chi})$ and $\mathfrak{R}_{\omega \otimes \chi}$ are also the same for (G, P) as for $(M(\chi), P \cap M(\chi))$. Now it follows from [28, Proposition 2.14.c] and (11) that for χ positive with respect to P

$$\text{End}_G(I_P^G(\omega \otimes \chi)) \cong \text{End}_{M(\chi)}(I_{P \cap M(\chi)}^{M(\chi)}(\omega \otimes \chi)) \cong \mathbb{C}[\mathfrak{R}_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi}].$$

The construction of the isomorphism (11) is unique up to algebra automorphisms of $\mathbb{C}[\mathfrak{R}_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi}]$ which preserve each of the one-dimensional subspaces $\mathbb{C}w$. Every such automorphism comes from twisting by a character of $\mathfrak{R}_{\omega \otimes \chi}$.

(c) In view of the remarks at the start of the proof, this holds with respect to the group $M(\chi)$ (instead of G). But $\mathbb{C}[\mathfrak{R}_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi}]$ is the same for (G, P) and $(M(\chi), P \cap M(\chi))$, so we obtain the result for G by applying the functor $I_{PM(\chi)}^G$ to the result for $M(\chi)$.

(b) For the same reason as (c), this holds on the level of $M(\chi)$. Choose a parabolic subgroup P' containing M , with respect to which χ is positive. Then $P' \cap M(\chi) = P \cap M(\chi)$, so

$$\pi(M, \omega \otimes \chi, \rho) = \text{Hom}_{\mathbb{C}[\mathfrak{R}_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi}]}(\rho, I_{P' \cap M(\chi)}^{M(\chi)}(\omega \otimes \chi)).$$

By Lemma 1.1 $I_P(M, \omega \otimes \chi, \rho)$ and $I_{P'}^G(M, \omega \otimes \chi, \rho)$ have the same irreducible constituents and by the Langlands classification there is a unique Langlands quotient among them. This provides the bijection $\text{Irr}_{M, \omega \otimes \chi}(M(\chi)) \rightarrow \text{Irr}_{M, \omega \otimes \chi}(G)$. \square

We remark that, since parabolic induction preserves irreducibility of representations in most cases, $L(M, \omega \otimes \chi, \rho) = I_{PM(\chi)}^G(\pi(M, \omega \otimes \chi, \rho))$ for χ in a Zariski-open subset of $\mathcal{O}_{\mathbb{C}}$. For $\omega \otimes \chi \in \mathcal{O}$ the stronger $L(M, \omega \otimes \chi, \rho) = \pi(M, \omega \otimes \chi, \rho)$ holds, because then $M(\chi) = G$.

Theorem 1.6 gives rise to a conjectural parametrization of L-packets. Suppose that ϕ is a Langlands parameter for G , which is elliptic for a Levi subgroup

$M \subset G$. By [6, §10.3] the L-packet $\Pi_\phi(M)$ should consist of essentially square-integrable representations. If $N_G(M, \phi)$ denotes the stabilizer of this L-packet in G , Theorem 1.2 shows that $N_G(M, \phi)$ -associate elements of $\Pi_\phi(M)$ yield the same parabolically induced representations. The conjectural compatibility of the local Langlands correspondence with parabolic induction and with the formation of Langlands quotients make it reasonable to expect that

$$(12) \quad \begin{aligned} \Pi_\phi(G) &= \bigsqcup_{\omega \otimes \chi \in \Pi_\phi(M)/N_G(M, \phi)} \text{Irr}_{M, \omega \otimes \chi}(G) \\ &= \bigsqcup_{\omega \otimes \chi \in \Pi_\phi(M)/N_G(M, \phi)} \{L(M, \omega \otimes \chi, \rho) : \rho \in \text{Irr}(\mathbb{C}[\mathfrak{R}_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi}])\}. \end{aligned}$$

2. ALGEBRAIC FAMILIES OF IRREDUCIBLE REPRESENTATIONS

Let X be a real or complex algebraic variety. By an algebraic family of G -representations we mean a family $\{\pi_x : x \in X\}$ such that all the π_x are realized on the same vector space (up to some natural isomorphism) and the matrix coefficients depend algebraically on x .

Lemma 1.5 and Theorem 1.6 can be used to give a rough description of the geometric structure of the Bernstein component of $\text{Irr}(G)$ determined by \mathcal{O} , in terms of algebraic families. For any subset $Y \subset \mathcal{O}_{\mathbb{C}}$ we define

$$\text{Irr}_{M, Y}(G) := \bigcup_{\omega \otimes \chi \in Y} \text{Irr}_{M, \omega \otimes \chi}(G).$$

Proposition 2.1. *Let Y be a maximal connected subset of $\mathcal{O}_{\mathbb{C}}$ on which $W_{\omega \otimes \chi}$ is constant.*

- (a) *Y is of the form $X \setminus X^*$, where X is a coset of a complex subtorus of $\mathcal{O}_{\mathbb{C}}$ and X^* is a finite union of cosets of complex subtori of smaller dimension than X .*
- (b) *Let $W(\mathcal{O})_X$ be the (setwise) stabilizer of X in $W(\mathcal{O})$. Theorem 1.6 determines a natural bijection*

$$(X \setminus X^*)/W(\mathcal{O})_X \times \text{Irr}_{M, \omega \otimes \chi}(G) \rightarrow \text{Irr}_{M, X \setminus X^*}(G),$$

for any $\omega \otimes \chi \in X \setminus X^$.*

- (c) *Representations in $\text{Irr}_{M, X \setminus X^*}(G)$ are tempered if and only if the parameter $\omega \otimes \chi$ is in $X_{\text{cpt}} \setminus X_{\text{cpt}}^*$, the canonical real form of $X \setminus X^*$.*

Proof. Consider $\mathcal{O}_{\mathbb{C}}$ as an algebraic group via the bijection (4). The invertible elements in the coordinate ring $\mathbb{C}[\mathcal{O}_{\mathbb{C}}] \cong \mathbb{C}[X^*(\mathcal{O}_{\mathbb{C}})]$ are

$$\mathbb{C}[\mathcal{O}_{\mathbb{C}}]^\times = \{zx : x \in X^*(\mathcal{O}_{\mathbb{C}}), z \in \mathbb{C}^\times\}.$$

Hence the action of $W(\mathcal{O})$ on $\mathcal{O}_{\mathbb{C}}$ induces a group action on $\mathbb{C}[\mathcal{O}_{\mathbb{C}}]^\times / \mathbb{C}^\times \cong X^*(\mathcal{O}_{\mathbb{C}})$, say $(w, x) \mapsto \lambda_w(x)$. Then

$$w \cdot zx = zt_w(x)\lambda_w(x) \quad \text{for a unique } t_w(x) \in \mathbb{C}^\times.$$

Clearly t_w determines a group homomorphism $X^*(\mathcal{O}_{\mathbb{C}}) \rightarrow \mathbb{C}^\times$, so it can be regarded as an element of $\mathcal{O}_{\mathbb{C}}$. Thus we decomposed the action of $w \in W(\mathcal{O})$

on $\mathcal{O}_{\mathbb{C}}$ as $t_w \lambda_w$, where λ_w is an automorphism of $\mathcal{O}_{\mathbb{C}}$ as an algebraic group and t_w is translation by an element of $\mathcal{O}_{\mathbb{C}}$. The fixed points of such a transformation are of the form

$$\mathcal{O}_{\mathbb{C}}^w = (\mathcal{O}_{\mathbb{C}}^{\lambda_w})^{\circ} F_w \text{ for some finite subset } F_w \subset \mathcal{O}_{\mathbb{C}}^w.$$

Furthermore $(\mathcal{O}_{\mathbb{C}}^{\lambda_w})^{\circ}$ is an algebraic torus, since it is the image of the λ_w -invariants in the Lie algebra of $\mathcal{O}_{\mathbb{C}}$ under the exponential map. More generally, for any subgroup $W \subset W(\mathcal{O})$,

$$\mathcal{O}_{\mathbb{C}}^W = (\mathcal{O}_{\mathbb{C}}^{\lambda^{(W)}})^{\circ} F_W.$$

The subset $(\mathcal{O}_{\mathbb{C}}^W)^* \subset \mathcal{O}_{\mathbb{C}}^W$ of points with a stabilizer strictly larger than W arises from sets of the same shape, so it is union of cosets of algebraic tori of $(\mathcal{O}_{\mathbb{C}}^{\lambda^{(W)}})^{\circ}$. For $W = W_Y$ we get

$$X = (\mathcal{O}_{\mathbb{C}}^{\lambda^{(W_Y)}})^{\circ} (\omega \otimes \chi) \quad \text{and} \quad X^* = (\mathcal{O}_{\mathbb{C}}^{\lambda^{(W_Y)}})^* \cap X,$$

which are of the required form.

(b) The bijection is constructed with Theorems 1.2, 1.6 and Lemma 1.5. To see that it is natural, consider a $\omega \otimes \chi \in \mathcal{O} \cap X \setminus X^*$ and abbreviate $A = \text{End}_G(I_P^G(\omega \otimes \chi))$. Since all the representations $I_P^G(\omega \otimes \chi')$ are realized on the same vector space, Theorem 1.6.a shows that $A \subset \text{End}_G(I_P^G(\omega \otimes \chi'))$ for all $\omega \otimes \chi' \in X \setminus X^*$, and that A determines the decomposition of $I_P^G(\omega \otimes \chi')$ into indecomposable representations. If we substitute A for $\mathbb{C}[\mathfrak{R}_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi}]$ in Theorem 1.6.b we obtain the same bijection as in part (b) of the current proposition. This makes it clear that twisting by characters in Theorem 1.6.a does not effect the bijection, so it is natural.

(c) is merely a restatement of Theorem 1.2.c. \square

For $\rho \in \text{Irr}(\mathbb{C}[\mathfrak{R}_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi}])$ we put

$$(13) \quad \text{Irr}_{M, X \setminus X^*, \rho}(G) = \{L(M, \omega \otimes \chi, \rho) : \omega \otimes \chi \in X \setminus X^*\}.$$

By Proposition 2.1.b this set is in bijection with the complex quasi-affine variety $(X \setminus X^*)/W(\mathcal{O})_X$. By Proposition 2.1.c

$$(14) \quad \text{Irr}^t(G) \cap \text{Irr}_{M, X \setminus X^*, \rho}(G) = \text{Irr}_{M, X_{\text{cpt}} \setminus X_{\text{cpt}}^*, \rho}(G) := \{\pi(M, \omega \otimes \chi, \rho) : \omega \otimes \chi \in X_{\text{cpt}} \setminus X_{\text{cpt}}^*\},$$

which is in bijection with the real form $(X_{\text{cpt}} \setminus X_{\text{cpt}}^*)/W(\mathcal{O})_X$ of $(X \setminus X^*)/W(\mathcal{O})_X$. By Theorem 1.2.b two such families $\text{Irr}_{M_1, X_1 \setminus X_1^*, \rho_1}(G)$ and $\text{Irr}_{M_2, X_2 \setminus X_2^*, \rho_2}(G)$ are either disjoint or equal. The latter happens if and only if there is a $g \in G$ such that $gM_2g^{-1} = M_1$ and $(g \cdot X_2 \setminus X_2^*, g \cdot \rho_2)$ is $W(M_1)$ -equivalent with $(X_1 \setminus X_1^*, \rho_1)$. In this way $\text{Irr}(G)$ can be regarded as the complexification of $\text{Irr}^t(G)$.

For $\text{Irr}_{M, X \setminus X^*, \rho}(G)$ as in (13), let X^{\sharp} be the union of X^* and the $\omega \in X \setminus X^*$ for which the Langlands quotient $L(M, \omega \otimes \chi, \rho)$ is not the whole of $I_{PM(\chi)}^G(\pi(M, \omega \otimes \chi, \rho))$. Then $\text{Irr}_{M, X \setminus X^{\sharp}, \rho}(G)$ is an algebraic family of irreducible G -representations.

3. ALGEBRAIC FAMILIES OF LANGLANDS PARAMETERS

Let $\check{G} = \check{\mathcal{G}}(\mathbb{C})$ be the complex dual group of $G = \mathcal{G}(F)$. Let E/F be a finite Galois extension over which \mathcal{G} splits. The choice of a pinning (also known as a splitting) for G determines an action of the Galois group $\text{Gal}(E/F)$ on \check{G} . As Langlands dual group we take

$${}^L G = \check{G} \rtimes \text{Gal}(E/F).$$

Recall that the Weil group of F can be written as $\mathbf{W}_F = \mathbf{I}_F \rtimes \langle \text{Frob} \rangle$, where \mathbf{I}_F is the inertia subgroup and Frob is a Frobenius element of \mathbf{W}_F . A Langlands parameter for G is a continuous group homomorphism

$$\phi: \mathbf{W}_F \times \text{SL}_2(\mathbb{C}) \rightarrow {}^L G$$

such that:

- $\phi(x) = \phi^\circ(x) \rtimes \text{pr}(x)$, with $\phi^\circ: \mathbf{W}_F \times \text{SL}_2(\mathbb{C}) \rightarrow \check{G}$ and $\text{pr}: \mathbf{W}_F \times \text{SL}_2(\mathbb{C}) \rightarrow \mathbf{W}_F \rightarrow \text{Gal}(E/F)$ the natural projection;
- $\phi(w)$ is semisimple for $w \in \mathbf{W}_F$;
- $\phi|_{\text{SL}_2(\mathbb{C})}: \text{SL}_2(\mathbb{C}) \rightarrow \check{G}$ is a homomorphism of algebraic groups.

We say that ϕ is relevant for G if, whenever the image of ϕ is contained in a parabolic subgroup ${}^L P$ [6, §3], ${}^L P^\circ$ corresponds to a parabolic subgroup of \mathcal{G} which is defined over F . (This condition is empty if G is quasi-split.) We define $\Psi(G)$ to be the set of relevant Langlands parameters for G and $\Phi(G)$ to be $\Psi(G)/\check{G}$ with respect to the conjugation action.

We say that $\phi \in \Psi(G)$ is bounded if $\phi(\mathbf{W}_F)$ is bounded. Since \mathbf{I}_F is compact and ϕ is continuous, ϕ is bounded if and only if $\phi^\circ(\text{Frob})$ lies in a compact subgroup of \check{G} . We denote the subsets of bounded elements in $\Psi(G)$ and $\Phi(G)$ by $\Psi_{\text{bdd}}(G)$ and $\Phi_{\text{bdd}}(G)$.

Lemma 3.1. *Every $\phi \in \Psi(G)$ can be written as $\phi = \phi_{\text{nr}}\phi_f$ with $\phi_f \in \Psi(G)$, $\phi_f(\mathbf{W}_F)$ finite and*

$$\phi_{\text{nr}}: \mathbf{W}_F \times \text{SL}_2(\mathbb{C})/\mathbf{I}_F \times \text{SL}_2(\mathbb{C}) \rightarrow Z_{\check{G}}(\text{im } \phi_f)^\circ.$$

Proof. Heiermann [15, Lemma 5.1] proved the corresponding result for "admissible homomorphisms" $\mathbf{W}_F \rightarrow {}^L G$. His proof remains valid for our Langlands parameters. Although [15] says only that $\phi(\text{Frob}) \in Z_{\check{G}}(\text{im } \phi_f)$, the proof shows that $\phi(\text{Frob})$ lies in the identity component of the latter group. \square

We remark that in general ϕ_f is not uniquely determined by ϕ , there can be finitely many choices for $\phi_f(\text{Frob})$.

Suppose now that $\phi_f \in \Psi(G)$, with $\phi_f(\mathbf{W}_F)$ finite, is given. For $s \in Z_{\check{G}}(\text{im } \phi_f)^\circ$ the element $s\phi_f(\text{Frob})$ is semisimple if and only if s is semisimple. In this case there is a Langlands parameter

$$\phi_{f,s} := \phi_{\text{nr},s}\phi_f \quad \text{with} \quad \phi_{\text{nr},s}(\text{Frob}) = s.$$

Every parabolic subgroup that contains $\text{im } \phi_{f,s}$ also contains $\text{im } \phi_f$, so the relevance of ϕ_f implies that $\phi_{f,s}$ is relevant for G . We put

$$\Psi(G, \phi_f) = \{\phi' \in \Psi(G) : \phi'_f \sim \phi_f\},$$

where \sim means that ϕ_f is a possible choice for ϕ'_f . Let $\Phi(G, \phi_f)$ be the image of $\Psi(G, \phi_f)$ in $\Phi(G)$.

Since $\mathrm{im}(\phi_f)^\circ = \phi(\mathrm{SL}_2(\mathbb{C}))$ is reductive, so is $Z_{\tilde{G}}(\mathrm{im} \phi_f)^\circ$. Lemma 3.1 and the above show that $\Psi(G, \phi_f)$ is naturally parametrized by the set of semisimple elements $Z_{\tilde{G}}(\mathrm{im} \phi_f)_{\mathrm{ss}}^\circ$. Clearly $\Psi(G)$ is the union (usually not disjoint) of the subsets $\Psi(G, \phi_f)$. Since $Z_{\tilde{G}}(\mathrm{im} \phi_f)_{\mathrm{ss}}^\circ$ is the union of the tori T in $Z_{\tilde{G}}(\mathrm{im} \phi_f)^\circ$, we can write

$$(15) \quad \begin{aligned} \Psi(G) &= \bigcup_{\phi_f, T} \Psi(G, \phi_f, T) := \bigcup_{\phi_f, T} \{\phi_{f,s} : s \in T\}, \\ \Phi(G) &= \bigcup_{\phi_f, T} \Phi(G, \phi_f, T) := \bigcup_{\phi_f, T} (\text{image of } \Psi(G, \phi_f, T) \text{ in } \Phi(G)). \end{aligned}$$

Because all maximal tori of the complex reductive group $Z_{\tilde{G}}(\mathrm{im} \phi_f)^\circ$ are conjugate, we need only one maximal torus T for each choice of ϕ_f to obtain the whole of $\Phi(G)$. Conjugation by any element of \tilde{G} sends any family $\Psi(G, \phi_f, T)$ to another such family, via an isomorphism of tori. Consequently

$$(16) \quad \Phi(G, \phi_{1f}, T_1) \cap \Phi(G, \phi_{2f}, T_2) \text{ is empty or } \Phi(G, \phi_{1f}, T_1 \cap T'_2)$$

for some torus $T'_2 \subset Z_{\tilde{G}}(\mathrm{im} \phi_f)^\circ$. We remark that here and below we allow tori of dimension zero, which are just points.

Let T_{cpt} denote the maximal compact subgroup of a complex torus T , so in particular T is the complexification of T_{cpt} . Then

$$(17) \quad \begin{aligned} \Psi_{\mathrm{bdd}}(G) &= \bigcup_{\phi_f, T} \Psi(G, \phi_f, T_{\mathrm{cpt}}) := \bigcup_{\phi_f, T} \{\phi_{f,s} \in \Psi(G) : s \in T_{\mathrm{cpt}}\}, \\ \Phi_{\mathrm{bdd}}(G) &= \bigcup_{\phi_f, T} \Phi_{\mathrm{bdd}}(G, \phi_f, T) := \bigcup_{\phi_f, T} (\text{image of } \Psi_{\mathrm{bdd}}(G, \phi_f, T) \text{ in } \Phi(G)). \end{aligned}$$

For T_2 and T'_2 as in (16)

$$(18) \quad \Phi(G, \phi_{1f}, T_{1\mathrm{cpt}}) \cap \Phi(G, \phi_{2f}, T_{2\mathrm{cpt}}) \text{ is empty or } \Phi(G, \phi_{1f}, (T_1 \cap T'_2)_{\mathrm{cpt}}).$$

By (16) and (18) the intersections between such sets, which are partially caused by the ambiguity of $\phi \mapsto \phi_f$, do not pose any problems for this way of decomposing the space of Langlands parameters. In the sense of (15) and (17) $\Psi(G)$ can be regarded as the complexification of $\Psi_{\mathrm{bdd}}(G)$. The action of \tilde{G} preserves the structure introduced above, which enables us to see $\Phi(G)$ as the complexification of $\Phi_{\mathrm{bdd}}(G)$.

Now we include the S-groups from [3] in the picture. These are improved versions of the usual component groups. Let \tilde{G}_{sc} be the simply connected cover of the derived group of \tilde{G} . It acts on \tilde{G} by conjugation. For $\phi \in \Psi(G)$ consider the groups

$$C(\phi) := Z_{\tilde{G}_{sc}}(\mathrm{im} \phi) \quad \text{and} \quad \mathcal{S}_\phi := C(\phi)/C(\phi)^\circ.$$

(Arthur calls these groups $S_{\phi, sc}$ and \tilde{S}_ϕ .) Enhanced Langlands parameters for G are pairs (ϕ, ρ) with $\phi \in \Psi(G)$ and $\rho \in \mathrm{Irr}(\mathcal{S}_\phi)$. We call the set of such

parameters $\Psi^e(G)$. The conjugation action of \check{G} on $\Psi(G)$ extends naturally to an action on $\Psi^e(G)$, namely

$$\check{g} \cdot (\phi, \rho) = (\check{g}\phi\check{g}^{-1}, \rho \circ \text{Ad}(\check{g})^{-1}).$$

We denote the set of equivalence classes by $\Phi^e(G)$.

Let X be a real or complex algebraic variety. We say that a family $\{(\phi_x, \rho_x) : x \in X\}$ of enhanced Langlands parameters is an algebraic family if $\phi_x|_{\mathbf{I}_F \times \text{SL}_2(\mathbb{C})}$ is independent of x , $\phi_x(\text{Frob})$ depends algebraically on x and all the ρ_x are (in some sense) equivalent.

Let Z be the centralizer in \check{G} of some element of $\Psi(G, \phi_f, Y)$, where Y is a torus as in (15). Write $\mathfrak{t}_Z = \text{Lie}(Y) \cap Z_{\text{Lie}(\check{G})}(Z)$ and put $T_Z = \exp(\mathfrak{t}_Z)$, a subtorus of Y . The elements $\phi \in \Psi(G, \phi_f, Y)$ with $Z_{\check{G}}(\text{im } \phi) \supset Z$ correspond bijectively to a set of the form

$$Y_Z = T_Z F_Z \subset Y,$$

where F_Z is finite. We remark that Y_Z need not contain the unit element. The subset of $\phi \in \Psi(G, \phi_f, Y)$ with $Z_{\check{G}}(\text{im } \phi) \supsetneq Z$ determines a finite union Y_Z^* of cosets of algebraic subtori of smaller dimension in T_Z . Of course Y_Z can be empty. Let $T \subset Y_Z$ be a coset of an algebraic subtorus of T_Z and write $T^* = Y_Z^* \cap T$. For $\rho \in \text{Irr}(\mathcal{S}_\phi)$ we have an algebraic family

$$(19) \quad \Psi(G, \phi_f, T \setminus T^*, \rho) = \{(\phi_{f,s}, \rho) : s \in T \setminus T^*\}.$$

Let $\Phi(G, \phi_f, T \setminus T^*, \rho)$ be its image in $\Phi^e(G)$. Conjugation by an element of \check{G} sends $\Psi(G, \phi_f, T \setminus T^*, \rho)$ to a family of the same form. It follows that

$$(20) \quad \Phi(G, \phi_{1f}, T_1 \setminus T_1^*, \rho) \cap \Phi(G, \phi_{2f}, T_2 \setminus T_2^*, \sigma) \text{ is empty or } \Phi(G, \phi_{1f}, T_1 \cap T_2', \rho)$$

for some subtorus $T_2' \subset Y$. Similarly the set of enhanced bounded Langlands parameters $\Psi_{\text{bdd}}^e(G)$ is a union of the algebraic families

$$(21) \quad \Psi(G, \phi_f, T_{\text{cpt}} \setminus T_{\text{cpt}}^*, \rho) := \{(\phi_{f,s}, \rho) : s \in T_{\text{cpt}} \setminus T_{\text{cpt}}^*\}.$$

Again we denote the image in $\Phi^e(G)$ by $\Phi(G, \phi_f, T_{\text{cpt}} \setminus T_{\text{cpt}}^*, \rho)$. By (18) the intersections of such families satisfy

$$\Phi(G, \phi_{1f}, T_{1\text{cpt}} \setminus T_{1\text{cpt}}^*, \rho) \cap \Phi(G, \phi_{2f}, T_{2\text{cpt}} \setminus T_{2\text{cpt}}^*, \sigma) \text{ is empty or } \Phi(G, \phi_{1f}, (T_1 \cap T_2')_{\text{cpt}} \setminus T_{\text{cpt}}^*, \rho),$$

where T_2' as in (20). We summarize the findings of this section in a proposition:

Proposition 3.2.

- (a) $\Psi^e(G)$ is in a natural way a union of algebraic families $\Psi(G, \phi_f, T \setminus T^*, \rho)$, each of which is parametrized a complex variety $T \setminus T^*$. Every T is a coset of a torus in \check{G} , and T^* is a (possibly empty) finite union of cosets of tori of smaller dimension.
- (b) $\Psi_{\text{bdd}}^e(G)$ is in a natural way a union of algebraic families $\Psi(G, \phi_f, T_{\text{cpt}} \setminus T_{\text{cpt}}^*, \rho)$, each of which is parametrized by the canonical real form $T_{\text{cpt}} \setminus T_{\text{cpt}}^*$ of the variety $T \setminus T^*$.
- (c) Via (a) and (b) $\Psi^e(G)$ can be regarded as the complexification of $\Psi_{\text{bdd}}^e(G)$.

- (d) The action of \check{G} on $\Psi^e(G)$ preserves these structures, and in that sense $\Phi^e(G)$ can be seen as the complexification of $\Phi_{\text{bdd}}^e(G)$.

Example 3.3. We will work out the above families for (enhanced) Langlands parameters for $G = \text{SL}_2(F)$, which are trivial on the inertia group \mathbf{I}_F . Put $\check{G} = \text{PGL}_2(\mathbb{C})$ and let \check{T} be torus of diagonal elements in \check{G} . The simply connected cover of \check{G} is $\check{G}_{sc} = \text{SL}_2(\mathbb{C})$ and we let \check{T}_{sc} be the torus of diagonal elements therein. We distinguish the families first by their restriction to $\text{SL}_2(\mathbb{C})$ and then by the possible tori.

- $\phi|_{\text{SL}_2(\mathbb{C})} = 1, \phi_{1f} = 1, T_1 = \check{T}$.

Then $T_1^* = \{(\begin{smallmatrix} 1 & 0 \\ 0 & 1 \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 0 & -1 \end{smallmatrix})\}$ and for all $\phi \in \Psi(G, 1, T_1 \setminus T_1^*)$ we have $C(\phi) = \check{T}_{sc}$ and $\mathcal{S}_\phi = 1$. Moreover

$$\Phi(G, 1, T_1 \setminus T_1^*) = \{\phi_{\text{nr}, s} : s \in T_1 \setminus T_1^*\} / W(\check{G}, \check{T}) \cong (\mathbb{C}^\times \setminus \{1, -1\}) / S_2,$$

$$\Phi(G, 1, T_{1\text{cpt}} \setminus T_{1\text{cpt}}^*) \cong \{z \in \mathbb{C}^\times : |z| = 1, z \neq 1, z \neq -1\} / S_2.$$

- $\phi|_{\text{SL}_2(\mathbb{C})} = 1, \phi_{2f} = 1, T_2 = 1$.

Now T_2^* is empty, $C(\phi) = \text{SL}_2(\mathbb{C})$ and $\mathcal{S}_\phi = 1$. Thus $\Phi(G, 1, T_2) = \{1\}$.

- $\phi|_{\text{SL}_2(\mathbb{C})} = 1, \phi_{3f} = \phi_{\text{nr}, (\begin{smallmatrix} 1 & 0 \\ 0 & -1 \end{smallmatrix})}, T_3 = 1$.

In this case T_3^* is empty, $C(\phi) = N_{\check{G}_{sc}}(\check{T}_{sc})$ and $\mathcal{S}_\phi = W(\check{G}_{sc}, \check{T}_{sc}) = S_2$. For every $\rho \in \text{Irr}(\mathcal{S}_\phi)$ we have $\Phi(\check{G}, \phi_{3f}, T_3, \rho) = \{(\phi_{3f}, \rho)\}$.

- $\phi|_{\text{SL}_2(\mathbb{C})}$ the projection $\text{SL}_2(\mathbb{C}) \rightarrow \text{PGL}_2(\mathbb{C})$, ϕ_{4f} trivial on \mathbf{W}_F and $T_4 = 1$.

Again $T_4^* = \emptyset$ and there is only one Langlands parameter $\phi = \phi_{4f}$ in this family, which satisfies $C(\phi) = Z(\text{SL}_2(\mathbb{C})) = \mathcal{S}_\phi$. For $\rho \in \text{Irr}(\mathcal{S}_\phi)$ we obtain $\Phi(G, \phi_{4f}, T_4, \rho) = \{(\phi_{4f}, \rho)\}$. We remark that for ρ nontrivial (ϕ_{4f}, ρ) does not parametrize a representation of G , but one of the essentially unique non-split inner form of G .

4. FROM A TEMPERED TO A GENERAL LOCAL LANGLANDS CORRESPONDENCE

In this section we will show how a local Langlands correspondence for $\text{Irr}^t(G)$ can be extended to $\text{Irr}(G)$. For this purpose we want the enhanced Langlands parameters, so that every L-packet $\Pi(\phi)$ is split into singletons by $\text{Irr}(\mathcal{S}_\phi)$. As not all irreducible representations of the S-group \mathcal{S}_ϕ need to appear here, we suppose that the LLC is an injective map from $\text{Irr}(G) \rightarrow \Phi^e(G)$. Of course we need to impose additional conditions on this LLC, which we discuss now.

Recall the algebraic families of irreducible G -representations and of enhanced Langlands parameters from Sections 2 and 3. We would like to say that via the local Langlands correspondence every algebraic family on one side is in bijection with an algebraic family on the other side. Unfortunately this is not true in general, because our algebraic families need not be maximal. In both $\Psi(G, \phi_f, T \setminus T^*, \rho)$ and $\text{Irr}_{M, X \setminus X^*, \rho}(G)$ it is possible that some points of T^* (resp. X^*) have a larger centralizer in \check{G} (resp. in $W(\mathcal{O})$), but the same

S-groups (resp. R-groups) as points of $T \setminus T^*$ (resp. $X \setminus X^*$). Then the algebraic family can be extended to a larger subvariety. This behaviour is very common, it occurs for most reductive p -adic groups. We could overcome this problem by adjusting the definitions of $\Psi(G, \phi_f, T \setminus T^*, \rho)$ and $\text{Irr}_{M, X \setminus X^*, \rho}(G)$ so that they include such points of T or X .

However, that would still not imply that our algebraic families are maximal. One reason is that $C(\phi_{f,t})$ could be larger than $C(\phi)$ for $\phi \in \Psi(G, \phi_f, T \setminus T^*)$, but that the subsets of $\text{Irr}(\mathcal{S}_{\phi_{f,t}})$ and $\text{Irr}(\mathcal{S}_\phi)$ that are relevant for the LLC could nevertheless be in natural bijection. Even more subtly, it is conceivable that $\mathfrak{R}_{\omega \otimes \chi}$ is strictly larger than the R-groups associated to $M, X \setminus X^*$, but still there exists a $\rho_{\omega \otimes \chi} \in \text{Irr}(\mathfrak{R}_{\omega \otimes \chi}, \kappa_{\omega \otimes \chi})$ such that $L(M, \omega \otimes \chi, \rho_{\omega \otimes \chi})$ fits in a natural way in $\text{Irr}_{M, X \setminus X^*, \rho}(G)$. Maybe such situations could be excluded with more precise conventions and some additional work.

We prefer to deal with this by proving two versions of our extension theorem: one that covers all situations which can theoretically arise inside the framework of the previous sections, and a more elegant version which works under slightly stronger conditions.

For the first version we assume only that every algebraic family, of the form described in Sections 2 and 3, is in correspondence with finitely many algebraic families, also as in Sections 2 and 3, on the other side (possibly minus some subfamilies of smaller dimension).

Theorem 4.1. *Let a tempered local Langlands correspondence for G be given as an injective map*

$$LL_G^t : \text{Irr}^t(G) \rightarrow \Phi_{\text{bdd}}^e(G).$$

Suppose that for every algebraic family of irreducible tempered G -representations $\text{Irr}_{M, X_{\text{cpt}} \setminus X_{\text{cpt}}^, \rho}(G)$ as in (13), there exist*

- (1) *finitely many algebraic families of enhanced bounded Langlands parameters $\Psi(G, \phi_f, T_{i,\text{cpt}} \setminus T_{i,\text{cpt}}^*, \rho_i)$ as in (21);*
- (2) *for every i , a coset $X_{i,\text{cpt}}$ of a compact subtorus of X_{cpt} and an isomorphism of real algebraic varieties $\psi_i : X_{i,\text{cpt}} \rightarrow T_{i,\text{cpt}}$;*
- (3) *an injection $\psi : X_{\text{cpt}} \setminus X_{\text{cpt}}^* \rightarrow \bigsqcup_i T_{i,\text{cpt}} \setminus T_{i,\text{cpt}}^*$;*

such that $\psi(\omega \otimes \chi) = \psi_i(\omega \otimes \chi)$ for $\omega \otimes \chi \in \psi_i^{-1}(T_{i,\text{cpt}} \setminus T_{i,\text{cpt}}^) \cap X_{\text{cpt}} \setminus X_{\text{cpt}}^*$, and*

$$(\phi_{f, \psi(\omega \otimes \chi)}, \rho_i) \in \Psi_{\text{bdd}}^e(G) \quad \text{represents} \quad LL_G^t(\pi(M, \omega \otimes \chi, \rho)).$$

Then LL_G^t can be extended in a unique way to an injective map

$$LL_G : \text{Irr}(G) \rightarrow \Phi^e(G)$$

such that

- (1) *the image of LL_G is the complexification of $LL_G^t(\text{Irr}^t(G))$ in the sense of Proposition 3.2;*
- (2) *the above conditions hold without the subscripts cpt.*

Proof. By complexification ψ_i extends to an isomorphism of complex algebraic varieties $\psi_i : X_i \rightarrow T_i$. Hence we can extend ψ to an injection

$$\begin{aligned} \psi : X \setminus X^* &\rightarrow \bigsqcup_i T_i \setminus T_i^*, \\ \psi(\omega \otimes \chi) &:= \psi_i(\omega \otimes \chi) \quad \text{for } \omega \otimes \chi \in \psi_i^{-1}(T_i \setminus T_i^*) \cap X \setminus X^*. \end{aligned}$$

Using this we put

$$LL'_G(M, \omega \otimes \chi, \rho) = (\phi_{f, \psi(\omega \otimes \chi)}, \rho_i) \in \Psi^e(G) \quad \text{for } \omega \otimes \chi \in \psi^{-1}(T_i \setminus T_i^*).$$

Notice that the argument is not a G -representation, but a parameter for that. By assumption $LL'_G(M, \omega \otimes \chi, \rho)$ represents $LL_G^t(\pi(M, \omega \otimes \chi, \rho))$ for $\chi \in X_{\text{unr}}(M)$. We want LL'_G to descend to a map $\text{Irr}(G) \rightarrow \Phi^e(G)$ via Theorem 1.6.b. Let us agree to use only one M from every conjugacy class of Levi subgroups of G . In view of Theorem 1.2 it suffices to check that

$$LL'_G(L(M, \omega \otimes \chi, \rho)) \text{ is } \check{G}\text{-conjugate to } LL'_G(L(M, w(\omega \otimes \chi), w\rho))$$

for all $w \in W(M)$. By construction this holds if $\chi \in X_{\text{unr}}(M)$. Otherwise $|\chi| \in X_{\text{nr}}(M)$ is of infinite order and

$$L(M, \omega \otimes \chi |\chi|^z, \rho) \cong L(M, w(\omega \otimes \chi |\chi|^z), w\rho) \quad \text{for all } z \in \mathbb{C}.$$

For $z \in i\mathbb{R} - 1$, $\chi |\chi|^z$ is unitary and

$$LL'_G(M, \omega \otimes \chi |\chi|^z, \rho) \text{ is } \check{G}\text{-conjugate to } LL'_G(M, w(\omega \otimes \chi |\chi|^z), w\rho)$$

because both represent $LL_G^t(\pi(M, \omega \otimes \chi |\chi|^z, \rho))$. These objects vary continuously with z , so we can find one element $\check{g} \in \check{G}$ which conjugates them for all $z \in i\mathbb{R} - 1$ simultaneously. Then \check{g} actually works for all $z \in \mathbb{C}$, and in particular for $\omega \otimes \chi$. We conclude that LL'_G induces a well-defined map $LL_G : \text{Irr}(G) \rightarrow \Phi^e(G)$.

By construction LL_G has all the properties described in the theorem, only the injectivity is not yet clear. Suppose that

$$\phi_1 = LL'_G(M, \omega_1 \otimes \chi_1, \rho_1) \quad \text{and} \quad \phi_2 = LL'_G(M, \omega_2 \otimes \chi_2, \rho_2)$$

are conjugate by some element $\check{g}' \in \check{G}$. Then

$$|\phi_1(\text{Frob})| = |\psi_1(\omega_1 \otimes \chi_1)| \text{ is } \check{G}\text{-conjugate to } |\phi_2(\text{Frob})| = |\psi_2(\omega_2 \otimes \chi_2)|,$$

by the same element \check{g}' . Hence

$$LL'_G(M, \omega_1 \otimes \chi_1 |\chi_1|^z, \rho_1) \text{ is } \check{G}\text{-conjugate to } LL'_G(M, \omega_2 \otimes \chi_2 |\chi_2|^z, \rho_2)$$

for all $z \in \mathbb{C}$. The injectivity of LL_G^t implies

$$L(M, \omega_1 \otimes \chi_1 |\chi_1|^z, \rho_1) \cong L(M, w(\omega_2 \otimes \chi_2 |\chi_2|^z), \rho_2) \quad \text{for all } z \in i\mathbb{R} - 1.$$

Proposition 2.1.b shows that this holds for all $z \in \mathbb{C}$, and in particular for $z = 0$. \square

For a cleaner version of this theorem we summarize the essence of algebraic families of irreducible representations in shorter terminology. Let $\pi \in \text{Irr}_{M,\omega}(G)$ with $\omega \in \text{Irr}(M)$ square-integrable modulo centre. For $\chi \in X_{\text{nr}}(M)$ we say that $\pi \otimes \chi$ is well-defined if there exists a path $t \mapsto \chi_t$ in $X_{\text{nr}}(M)$ with $\chi_0 = 1$ and $\chi_1 = \chi$, such that there is a canonical isomorphism $\mathfrak{R}_{\omega \otimes \chi_t} \cong \mathfrak{R}_\omega$ for all t . This definition makes sense by Lemma 1.5, while Proposition 2.1 shows how $\pi \otimes \chi$ can be constructed. In fact the $\pi \otimes \chi$ which are well-defined in this sense are precisely the members of a family of representations as in (13). Notice also that this convention generalizes the usual definition of $\pi \otimes \chi$ for $\chi \in X_{\text{nr}}(G)$.

In the above setting there is an inclusion $\check{M} \rightarrow \check{G}$, unique up to conjugation. We recall a desirable property of the local Langlands correspondence from [6, §10]: the Langlands parameter of π is that of ω , composed with the map $\check{M} \rightarrow \check{G}$. Equivalently, it is conjectured that π and ω have the same Langlands parameter up to conjugation by \check{G} .

The unramified character χ of M can be regarded as a character of the torus $Z(M)^\circ$. Via the LLC for tori it determines a smooth homomorphism $\hat{\chi} : \mathbf{W}_F \rightarrow Z(\check{M}) \subset \check{M}$.

Theorem 4.2. *Suppose that a tempered local Langlands correspondence for G is given as an injective map*

$$LL_G^t : \text{Irr}^t(G) \rightarrow \Phi_{\text{bdd}}^e(G).$$

Assume that for all $\pi \in \text{Irr}_{M,\omega}(G)$ we can find a representative $(\phi_\pi, \rho_\pi) \in \Psi_{\text{bdd}}^e(M)$ for $LL_G^t(\pi)$ such that, whenever $\chi \in X_{\text{nr}}(M)$ and $\pi \otimes \chi$ is well-defined (in the above sense):

- *there is a canonical isomorphism $\alpha_\chi : \mathcal{S}_{\phi_\pi} \rightarrow \mathcal{S}_{\phi_\pi \hat{\chi}}$;*
- *$(\phi_\pi \hat{\chi}, \alpha_\chi^* \rho_\pi) \in \Psi_{\text{bdd}}^e(M)$ represents $LL_G^t(\pi \otimes \chi)$.*

Then LL_G^t can be extended in a canonical way to an injection

$$LL_G : \text{Irr}(G) \rightarrow \Phi^e(G)$$

which fulfills the above conditions for all $\chi \in X_{\text{nr}}(M)$ such that $\pi \otimes \chi$ is well-defined.

Proof. It suffices to check that the conditions of Theorem 4.1 are fulfilled. Consider a family $\text{Irr}_{M, X_{\text{cpt}} \setminus X_{\text{cpt}}^*, \rho}(G)$ and an element $\omega \in X_{\text{cpt}} \setminus X_{\text{cpt}}^* \subset \text{Irr}^t(M)$. The assumptions enable us to find a family of Langlands parameters $\Psi(G, \phi_f, T_{\text{cpt}} \setminus T_{\text{cpt}}^*)$ which is in bijection with $\text{Irr}_{M, X_{\text{cpt}} \setminus X_{\text{cpt}}^*, \rho}(G)$ via $\pi \otimes \chi \rightarrow \phi_\pi \hat{\chi}$. Divide $\Psi(G, \phi_f, T_{\text{cpt}} \setminus T_{\text{cpt}}^*)$ into finitely many families of enhanced Langlands parameters $\Psi(G, \phi_f, T_{i,\text{cpt}} \setminus T_{i,\text{cpt}}^*, \rho_i)$ according to the different possibilities for $C(\phi_\pi \hat{\chi})$. Here the additional ingredient ρ_i is uniquely determined by the second assumption. Now we can apply Theorem 4.1. \square

The conditions of Theorem 4.2 hold in all cases which the authors checked, and it seems likely that they are valid for any p -adic group G (if a tempered local Langlands correspondence exists for G). For example they hold for all

inner forms of $\mathrm{GL}_n(F)$, because then the component groups and \mathfrak{R} -groups are trivial and compatibility with unramified twists is built in the LLC. In fact the usual LLC for $\mathrm{GL}_n(F)$, denoted $\mathrm{rec}_{F,n}$, fulfills the conditions of Theorem 4.2 for non-tempered representations as well. So if we start with $\mathrm{rec}_{F,n}|_{\mathrm{Irr}^t(\mathrm{GL}_n(F))}$, then Theorem 4.2 yields $\mathrm{rec}_{F,n}$.

The hypotheses are also fulfilled for inner forms of $\mathrm{SL}_n(F)$, as can be deduced from [17]. Furthermore both the work of Arthur [4] on quasi-split orthogonal and symplectic groups and the work on Mok on quasi-split unitary groups [23] should fit with Theorem 4.2. Indeed, the first condition in Theorem 4.2 will follow from the comparison of the analytic and geometric R-groups for tempered representations (see the next section), and the second condition should be a consequence of the functoriality of the twisted endoscopic transfers used in the construction of the representations of the classical and of the unitary groups.

5. GEOMETRIC R-GROUPS

In the next section we will explain why Theorem 4.2 applies to principal series representations of a split reductive p -adic group. To that end we first have to improve our understanding of the relations between \mathcal{S}_ϕ and the R-groups from Section 1, that is, between the analytic and the geometric R-groups. We discuss this for a general reductive p -adic group G .

Given $\phi \in \Psi(G)$, let M be a Levi subgroup of G such that the image of ϕ is contained in ${}^L M$, but not in any smaller Levi subgroup of ${}^L G$. We can regard \mathcal{S}_ϕ^M (that is, \mathcal{S}_ϕ for ϕ considered as a Langlands parameter for M) as a normal subgroup of \mathcal{S}_ϕ , so the conjugation action of \mathcal{S}_ϕ on \mathcal{S}_ϕ^M induces an action of the quotient $\mathcal{S}_\phi/\mathcal{S}_\phi^M$ on $\mathrm{Irr}(\mathcal{S}_\phi^M)$.

Definition 5.1. *Let $\mathfrak{R}_{\phi,\sigma}$ be the stabilizer of $\sigma \in \mathrm{Irr}(\mathcal{S}_\phi^M)$ in $\mathcal{S}_\phi/\mathcal{S}_\phi^M$. The group $\mathfrak{R}_{\phi,\sigma}$ is the geometric R-group attached to (ϕ, σ) .*

Assume that a local Langlands correspondence for essentially square-integrable representations of M is known, and that $(\phi, \sigma) \in \Psi^e(M)$ corresponds to $\omega \otimes \chi \in \mathrm{Irr}(M)$. The following conjecture extends to the non-tempered context a conjecture that was stated by Arthur in [1].

Conjecture 5.2. *$\mathfrak{R}_{\phi,\sigma}$ is isomorphic to $\mathfrak{R}_{\omega \otimes \chi}$.*

Conjecture 5.2 for tempered representations (Arthur's conjecture) is known to be true, in the case when F is of characteristic 0, when G is an inner form of $\mathrm{SL}_n(F)$ (see [12, 13]), and when G is a classical group, including the case of unitary groups, see [7] and the references therein. It was also studied, and proven in some other cases, in [24, §9].

In view of Propositions 2.1 and 3.2, the validity of Conjecture 5.2 for all bounded Langlands parameters $\phi \in \Phi_{\mathrm{bdd}}(G)$ would imply the validity for all $\phi \in \Phi(G)$.

On the other hand, the group $\mathfrak{R}_{\phi,\sigma}$ for $\phi \in \Psi_{\text{bdd}}^e(G)$ is a special case of the Arthur group $\mathfrak{R}_{\psi,\sigma}$ where $\psi: \mathbf{W}_F \times \text{SL}_2(\mathbb{C}) \times \text{SL}_2(\mathbb{C}) \rightarrow {}^L G$ is an Arthur parameter. Ban and Jantzen [8] provided an example in $G = \text{SO}_9(F)$, involving an Arthur parameter ψ that has non-trivial restriction to the second copy of $\text{SL}_2(\mathbb{C})$, for which the cardinality of $\mathfrak{R}_{\psi,\sigma}$ does not coincide with the number of components of the corresponding parabolically induced representation. However Conjecture 5.2 still holds in this case, as we will see.

Example 5.3. Ban and Jantzen considered the representation

$$\pi = \text{St}_{\text{GL}_2(F)} \times \text{triv}_{\text{GL}_2(F)} \rtimes 1,$$

which is parabolically induced from the representation $\text{St}_{\text{GL}_2(F)} \otimes \text{triv}_{\text{GL}_2(F)}$ of a Levi subgroup of the group $G = \text{SO}_9(F)$. The representation $\text{St}_{\text{GL}_2(F)} \otimes \text{triv}_{\text{GL}_2(F)}$ is not essentially square-integrable, so we want to compare π with the parabolically induced representation

$$\sigma = \nu^{1/2} \times \nu^{1/2} \times \text{St}_{\text{GL}_2(F)} \rtimes 1$$

where $\nu = |\det|_F$, to which our construction do apply. The representation π has three constituents [8, Theorem 2.5]:

$$\pi = Z(\nu^{-1/2}, \nu^{-1/2}; \tau_1) + Z(\nu^{-1/2}, \nu^{-1/2}; \tau_2) + Z(\nu^{-1/2}; \mathcal{S}),$$

where τ_1 , τ_2 , and \mathcal{S} are irreducible tempered representations of $\text{SO}_5(F)$ defined by

$$\tau_1 + \tau_2 = \text{St}_{\text{GL}_2} \rtimes 1 \quad \text{and} \quad \mathcal{S} = \text{St}_{\text{GL}_2} \rtimes \text{St}_{\text{SO}_3},$$

and $Z(\nu^{-1/2}, \nu^{-1/2}; \tau_i)$ is the unique subrepresentation of the parabolically induced representation $\nu^{-1/2} \times \nu^{-1/2} \rtimes \tau_i$, while $Z(\nu^{-1/2}; \mathcal{S})$ is the unique subrepresentation of the parabolically induced representation $\nu^{-1/2} \rtimes \mathcal{S} = \nu^{-1/2} \times \text{St}_{\text{GL}_2} \rtimes \text{St}_{\text{SO}_3}$. We have

$$Z(\nu^{-1/2}, \nu^{-1/2}; \tau_i) \in \text{Irr}_{M, \omega \otimes \chi}(G) \quad \text{and} \quad Z(\nu^{-1/2}; \mathcal{S}) \in \text{Irr}_{M', \omega' \otimes \chi'}(G),$$

where $M \simeq F^\times \times F^\times \times \text{GL}_2(F)$ and $\omega \otimes \chi = \nu^{-1/2} \otimes \nu^{-1/2} \otimes \text{St}_{\text{GL}_2}$, while $M' \simeq F^\times \times \text{GL}_2(F) \times \text{SO}_3(F)$ and $\omega' \otimes \chi' = \nu^{-1/2} \otimes \text{St}_{\text{GL}_2} \otimes \text{St}_{\text{SO}_3}$. Hence both the $Z(\nu^{-1/2}, \nu^{-1/2}; \tau_i)$ are Langlands constituents of σ and $Z(\nu^{-1/2}; \mathcal{S})$ falls into a different series, because it can be obtained via parabolic induction from a square-integrable representation of a larger parabolic subgroup.

In this example Arthur R -group $\mathfrak{R}_{\psi,\sigma}$ has four elements, clearly too many for the packet. If we do the same calculation as in [8, §2.3] with Langlands parameters instead of Arthur parameters, then we end up with a geometric R -group of order 2, which is also the analytic R -group of σ . Section 3.3 of [8] shows that its irreducible representations naturally parametrize the first two constituents of π discussed above.

We write $\text{Irr}(\mathcal{S}_\phi, \sigma) = \{\rho \in \text{Irr}(\mathcal{S}_\phi) : \text{Hom}_{\mathcal{S}_\phi^M}(\sigma, \rho) \neq 0\}$. Choose a minimal idempotent p_σ of $\mathbb{C}[\mathcal{S}_\phi^M]$ associated to σ . Then the algebra $\mathbb{C}[\mathcal{S}_\phi]p_\sigma\mathbb{C}[\mathcal{S}_\phi]$ is Morita equivalent to $p_\sigma\mathbb{C}[\mathcal{S}_\phi]p_\sigma$ and the map $V \mapsto p_\sigma V$ induces a bijection

$$(22) \quad \text{Irr}(\mathcal{S}_\phi, \sigma) = \text{Irr}(\mathbb{C}[\mathcal{S}_\phi]p_\sigma\mathbb{C}[\mathcal{S}_\phi]) \rightarrow \text{Irr}(p_\sigma\mathbb{C}[\mathcal{S}_\phi]p_\sigma).$$

On the other hand

$$p_\sigma \mathbb{C}[\mathcal{S}_\phi] p_\sigma \cong \text{End}_{\mathcal{S}_\phi}(\mathbb{C}[\mathcal{S}_\phi] p_\sigma) \cong \text{End}_{\mathcal{S}_\phi}(\mathbb{C}[\mathcal{S}_\phi] \otimes_{\mathbb{C}[\mathcal{S}_\phi^M]} \mathbb{C}[\mathcal{S}_\phi^M] p_\sigma) = \text{End}_{\mathcal{S}_\phi}(\text{ind}_{\mathcal{S}_\phi^M}^{\mathcal{S}_\phi} \sigma).$$

By [24, (9.1b)] a choice of intertwining operators $I_r \in \text{Hom}_{\mathcal{S}_\phi^M}(\sigma, r \cdot \sigma)$ for $r \in \mathfrak{R}_{\phi, \sigma}$ gives rise to a 2-cocycle $\kappa_{\phi, \sigma}$ such that

$$(23) \quad p_\sigma \mathbb{C}[\mathcal{S}_\phi] p_\sigma \cong \text{End}_{\mathcal{S}_\phi}(\text{ind}_{\mathcal{S}_\phi^M}^{\mathcal{S}_\phi} \sigma) \cong \mathbb{C}[\mathfrak{R}_{\phi, \sigma}, \kappa_{\phi, \sigma}].$$

Thus (22) and (23) provide a bijection between $\text{Irr}(\mathcal{S}_\phi, \sigma)$ and $\text{Irr}(\mathbb{C}[\mathfrak{R}_{\phi, \sigma}, \kappa_{\phi, \sigma}])$. We remark that in general this bijection is not natural, as it can depend on the choice of the intertwining operators I_r .

We call $\sigma \in \text{Irr}(\mathcal{S}_\phi^M)$ relevant for M if it corresponds to a representation of M (as opposed to a representation of an inner form of M), and we denote the set of such σ by $\text{Irr}_{\text{rel}M}(\mathcal{S}_\phi^M)$. The above action of \mathcal{S}_ϕ on $\text{Irr}(\mathcal{S}_\phi^M)$ permutes the different \mathcal{S}_ϕ^M -constituents of a representation of \mathcal{S}_ϕ , so on the p -adic side it should correspond to permuting the different $\omega \in \text{Irr}(M)$ for which $\text{Irr}_{M, \omega}(G)$ contains a fixed representation of G . Therefore $\text{Irr}_{\text{rel}M}(\mathcal{S}_\phi^M)$ should be stable under the action of \mathcal{S}_ϕ . The desirable properties of the local Langlands correspondence suggest that there are bijections

$$(24) \quad \begin{aligned} \Pi_\phi(G) = & \bigsqcup_{\omega \otimes \chi \in \Pi_\phi(M)/N_G(M, \phi)} \text{Irr}_{M, \omega \otimes \chi}(G) \longleftrightarrow \\ & \bigsqcup_{\sigma \in \text{Irr}_{\text{rel}M}(\mathcal{S}_\phi^M)/\mathcal{S}_\phi} \text{Irr}(\mathcal{S}_\phi, \sigma) \longleftrightarrow \bigsqcup_{\sigma \in \text{Irr}_{\text{rel}M}(\mathcal{S}_\phi^M)/\mathcal{S}_\phi} \text{Irr}(\mathbb{C}[\mathfrak{R}_{\phi, \sigma}, \kappa_{\phi, \sigma}]). \end{aligned}$$

Furthermore the comparison with (12) suggests that the cocycles $\kappa_{\phi, \sigma}$ and $\kappa_{\omega \otimes \chi}$ should be cohomologous via Conjecture (5.2). In that case (12) and (24) show how Langlands parameters for essentially square-integrable representations of Levi subgroups M can be used to produce a LLC for G . This fits well with the work of Heiermann [15], who proved that under certain conditions a parametrization of supercuspidal representations gives rise to one for essentially square-integrable representations.

6. THE PRINCIPAL SERIES OF A SPLIT GROUP

From now on we assume that G is F -split. The local Langlands correspondence for irreducible G -representations in the principal series was recently completed in [5]. It generalizes [19, 25] and relies among others on [26].

First we consider the unramified principal series. Let $(\phi, \sigma) \in \Psi_{\text{bdd}}^e(M)$ be elliptic for a Levi subgroup $M \subset G$ and let $\omega \in \text{Irr}(M)$ be the corresponding square-integrable (modulo centre) representation. The Kazhdan–Lusztig parametrization of irreducible Iwahori-spherical G -representations [19, 25] is compatible with parabolic induction in the sense that this operation does not change the first two ingredients of a Kazhdan–Lusztig parameter (s, u, ρ) . Since (s, u) determines a Langlands parameter, all elements of $\text{Irr}_{M, \omega}(G)$ have Langlands parameter $\phi \in \Phi(G)$.

The appropriate component groups for G -representations, at least for the principal series, are

$$Z_{\check{G}}(\phi)/Z_{\check{G}}(\phi)^\circ \cong \mathcal{S}_\phi/(\text{image of } Z(\check{G}_{sc})).$$

(In other words, the subtlety of replacing \check{G} by its simply connected cover is superfluous for split groups.) We denote the G -representation attached to ϕ in [19] by $\pi_G(\phi)$. By construction

$$\pi_G(\phi\hat{\chi}) = \pi_G(\phi) \otimes \chi \text{ for all } \chi \in X_{\text{nr}}(G).$$

In general $\pi_G(\phi)$ is reducible and endowed with a natural action of $Z_{\check{G}}(\phi)/Z_{\check{G}}(\phi)^\circ$. The third ingredient of a Kazhdan–Lusztig parameter is an irreducible representation ρ of the latter group. It is used to select an irreducible summand $\pi_G(\phi, \rho)$ of $\pi_G(\phi)$, by applying $\text{Hom}_{\mathcal{S}_\phi}(\rho, ?)$. Choose a $\sigma \in \text{Irr}(\mathcal{S}_\phi^M)$ which appears in the restriction of ρ to \mathcal{S}_ϕ^M . With (22) we obtain

$$(25) \quad \begin{aligned} \pi_G(\phi, \rho) &\cong \text{Hom}_{\mathcal{S}_\phi}(\rho, \pi_G(\phi)) = \text{Hom}_{\mathbb{C}[\mathcal{S}_\phi]_{p_\sigma\mathbb{C}[\mathcal{S}_\phi]}}(\rho, \pi_G(\phi)) \\ &\cong \text{Hom}_{p_\sigma\mathbb{C}[\mathcal{S}_\phi]_{p_\sigma}}(p_\sigma\rho, p_\sigma\pi_G(\phi)). \end{aligned}$$

By [19, Theorem 6.2] $\pi_G(\phi) \cong I_P^G(\pi_M(\phi))$ in an \mathcal{S}_ϕ^M -equivariant way. Let $\rho_\sigma \in \text{Irr}(\mathbb{C}[\mathfrak{R}_{\phi,\sigma}, \kappa_{\phi,\sigma}])$ correspond to $\rho \in \text{Irr}(\mathcal{S}_\phi, \sigma)$ via (22) and (23). It follows that the right hand side of (25) is isomorphic to

$$\text{Hom}_{\mathbb{C}[\mathfrak{R}_{\phi,\sigma}, \kappa_{\phi,\sigma}]}(\rho_\sigma, I_P^G(p_\sigma\pi_M(\phi))).$$

Because p_σ acts as a projection of rank one on the vector space underlying σ , it has essentially the same effect as applying $\text{Hom}_{\mathcal{S}_\phi^M}(\sigma, ?)$. We find

$$(26) \quad \begin{aligned} \pi_G(\phi, \rho) &\cong \text{Hom}_{\mathbb{C}[\mathfrak{R}_{\phi,\sigma}, \kappa_{\phi,\sigma}]}(\rho_\sigma, I_P^G(\text{Hom}_{\mathcal{S}_\phi^M}(\sigma, \pi_M(\phi)))) \\ &= \text{Hom}_{\mathbb{C}[\mathfrak{R}_{\phi,\sigma}, \kappa_{\phi,\sigma}]}(\rho_\sigma, I_P^G(\pi_M(\phi, \sigma))). \end{aligned}$$

Reeder [24, §9] proved that the analytic R-group \mathfrak{R}_ω is isomorphic to the subquotient $\mathfrak{R}_{\phi,\sigma}$ of $Z_{\check{G}}(\phi)/Z_{\check{G}}(\phi)^\circ$, and that the 2-cocycles $\kappa_{\phi,\sigma}$ and κ_ω are cohomologous. From this, (26) and Theorem 1.6.b it is clear that the way $\text{Irr}(Z_{\check{G}}(\phi)/Z_{\check{G}}(\phi)^\circ)$ is used here is equivalent to the method with R-groups in Section 1. We already knew that $\pi_M(\phi\hat{\chi}) = \pi_M(\phi) \otimes \chi$ for all $\chi \in X_{\text{nr}}(M)$, so the representations $\pi_G(\phi, \rho) \otimes \chi$ can just as well be constructed via (26). Consequently the Kazhdan–Lusztig parametrization satisfies $\pi_G(\phi\hat{\chi}, \rho) \cong \pi_G(\phi, \rho) \otimes \chi$ whenever this is well-defined for a $\chi \in X_{\text{unr}}(M)$.

Strictly speaking, [19] applies only if \check{G} has simply connected derived group. But Reeder’s generalization [25, Theorem 3.5.4] allows us to forget about this condition, as can be seen from equations (93) and (94) of [5]. Thus the assumptions of Theorem 4.2 are fulfilled for the unramified principal series of a split group.

In fact the Kazhdan–Lusztig–Reeder parametrization also fulfills the conclusion of Theorem 4.2. This can be shown by the above argument, combined with some Langlands quotients at the appropriate places. The latter do not

pose any additional problems, because they form an integral part of the constructions in [19].

Next we consider a Bernstein component $[T, \chi]_G$ in the principal series, such that the group $\tilde{H} = Z_{\tilde{G}}(\hat{\chi})$ is connected. Since the local Langlands correspondence for the irreducible representations in $[T, \chi]_G$ uses [26], we have to assume that the residual characteristic of F satisfies the mild conditions in [26, Remark 4.13]. According to [26, Theorem 9.14] the block of $\text{Rep}(G)$ determined by $[T, \chi]_G$ is equivalent with the block of $\text{Rep}(H)$ containing the unramified principal series. This equivalence comes from an isomorphism of Hecke algebras and it preserves all the important structure, like parabolic induction and R -groups. It was checked in [25, §4] that the component groups $Z_{\tilde{G}}(\phi)/Z_{\tilde{G}}(\phi)^\circ$ are also preserved in the process. Since the unramified principal series of H fit in the framework Theorem 4.2, as shown above, so does the Bernstein component $[T, \chi]_G$.

Finally, suppose that $Z_{\tilde{G}}(\hat{\chi})$ is disconnected, with identity component \tilde{H} and component group Γ . Then everything for the Bernstein component $[T, \chi]_G$ can be obtained from the setting for H , by taking the extended quotient (of the second kind) with respect to the action of Γ , see [5, §23]. This procedure is essentially the same for enhanced Langlands parameters as for G -representations, and therefore it does not disturb the properties of the local Langlands correspondence used in Theorem 4.2.

REFERENCES

- [1] J. Arthur, “Unipotent automorphic representations: conjectures”, *Astérisque*, **171-2** (1989), 13–71.
- [2] J. Arthur, “On elliptic tempered characters”, *Acta Math.* **171** (1993), 73–138.
- [3] J. Arthur, “A note on L-packets”, *Pure Appl. Math. Quarterly* **2.1** (2006), 199–217.
- [4] J. Arthur, *The endoscopic classification of representations: orthogonal and symplectic groups*, Colloquium Publication Series, American Mathematical Society (to appear), available at <http://www.claymath.org/cw/arthur/>
- [5] A.-M. Aubert, P. Baum, R.J. Plymen, M. Solleveld, “Geometric structure and the local Langlands conjecture”, *arXiv:1211.0180*.
- [6] A. Borel, “Automorphic L-functions”, *Proc. Symp. Pure Math* **33.2** (1979), 27–61.
- [7] D. Ban, D. Goldberg, “ R -groups and parameters”, *Pac. J. Math*, **255** (2012), 281–303.
- [8] D. Ban, C. Jantzen, “ R -groups and the action of intertwining operators in the nontempered case”, *Int. Math. Res. Not.* **20** (2007).
- [9] A. Borel, N.R. Wallach, *Continuous cohomology, discrete subgroups, and representations of reductive groups*, *Annals of Mathematics Studies* **94**, Princeton University Press, Princeton NJ, 1980.
- [10] J. Brodzki, R.J. Plymen, “Complex structure on the smooth dual of $\text{GL}(n)$ ”, *Doc. Math.* **7** (2002), 91–112.
- [11] W. Casselman, “Introduction to the theory of admissible representations of p -adic reductive groups”, preprint, 1995.
- [12] K.F. Chao, W. Li, “Dual R -groups of the inner forms of $\text{SL}(N)$ ”, *arXiv:1211.3039v2*.
- [13] K. Choï, D. Goldberg, “Transfer of R -groups between p -adic inner forms of SL_n ”, *arXiv:1211.5054v2*.
- [14] M. Harris, R. Taylor, “The geometry and cohomology of some simple Shimura varieties”, *Annals of Math. Studies* **151**, Princeton University Press, Princeton NJ, 2001.

- [15] V. Heiermann, “Orbites unipotentes et pôles d’ordre maximal de la fonction μ de Harish-Chandra”, *Canad. J. Math.* **58.6** (2006), 1203–1228.
- [16] G. Henniart, “Une preuve simple des conjectures de Langlands pour $GL(n)$ sur un corps p -adique”, *Inv. Math.* **139** (2000), 439–455.
- [17] K. Hiraga, H. Saito, “On L-packets for inner forms of SL_n ”, *Mem. Amer. Math. Soc.* 1013, Vol. **215** (2012).
- [18] T. Konno, “A note on the Langlands classification and irreducibility of induced representations of p -adic groups”, *Kyushu J. Math.* **57** (2003), 383–409.
- [19] D. Kazhdan, G. Lusztig, “Proof of the Deligne–Langlands conjecture for Hecke algebras”, *Invent. Math.* **87** (1987), 153–215.
- [20] G. Laumon, M. Rapoport, U. Stuhler, “ \mathcal{D} -elliptic sheaves and the Langlands correspondence”, *Invent. Math.* **113** (1993), 217–238.
- [21] G. Lusztig, “Some examples of square integrable representations”, *Trans. Amer. Math. Soc.* **277** (1983), 623–653.
- [22] C. Mœglin, M. Tadić, “Construction of discrete series for classical p -adic groups”, *J. Amer. Math. Soc.* **15** (2002), 715–786.
- [23] C.P. Mok, “Endoscopic classification of representations of quasi-split unitary groups I”, *arXiv:1206.0882*.
- [24] M. Reeder, “Euler–Poincaré pairings and elliptic representations of Weyl groups and p -adic groups”, *Compos. Math.* **129** (2001), 149–181.
- [25] M. Reeder, “Isogenies of Hecke algebras and a Langlands correspondence for ramified principal series representations”, *Representation Theory* **6** (2002), 101–126.
- [26] A. Roche, “Types and Hecke algebras for principal series representations of split reductive p -adic groups”, *Ann. Sci. École Norm. Sup.* **31.3** (1998), 361–413.
- [27] P. Schneider, E.-W. Zink, “K-types for the tempered components of a p -adic general linear group. With an appendix by Schneider and U. Stuhler”, *J. reine angew. Math.* **517** (1999), 161–208.
- [28] M. Solleveld, “Periodic cyclic homology of reductive p -adic groups”, *J. Noncommutative Geometry* **3.4** (2009), 501–558.
- [29] A.J. Silberger, “The Knapp–Stein dimension theorem for p -adic groups”, *Proc. Amer. Math. Soc.* **68.2** (1978), 243–246 and “The Knapp–Stein dimension theorem for p -adic groups. Correction”, *Proc. Amer. Math. Soc.* **76.1** (1979), 169–170.
- [30] A.J. Silberger, *Introduction to harmonic analysis on reductive p -adic groups*, *Mathematical Notes* **23**, Princeton University Press, Princeton NJ, 1979.
- [31] D. Vogan, “The local Langlands conjecture”, pp. 305–379 in: *Representation theory of groups and algebras*, *Contemp. Math.* **145**, American Mathematical Society, Providence RI, 1993.
- [32] J.-L. Waldspurger, “La formule de Plancherel pour les groupes p -adiques (d’après Harish-Chandra)”, *J. Inst. Math. Jussieu* **2.2** (2003), 235–333.

Anne-Marie Aubert

Institut de Mathématiques de Jussieu – Paris Rive-Gauche, U.M.R. 7586 du C.N.R.S.,
U.P.M.C., 4 place Jussieu 75005 Paris, France

E-mail: aubert@math.jussieu.fr

Pail Baum

Mathematics Department, Pennsylvania State University, University Park, PA 16802, USA

E-mail: baum@math.psu.edu

Roger Plymen

School of Mathematics, Southampton University, Southampton SO17 1BJ, England *and*
School of Mathematics, Manchester University, Manchester M13 9PL, England

E-mail: r.j.plymen@soton.ac.uk plymen@manchester.ac.uk

Maarten Solleveld

Radboud Universiteit Nijmegen, Heyendaalseweg 135, 6525AJ Nijmegen, the Netherlands

E-mail: m.solleveld@science.ru.nl