

***Invariant measures for the n -dimensional border
collision normal form***

Glendinning, Paul

2014

MIMS EPrint: **2015.10**

Manchester Institute for Mathematical Sciences
School of Mathematics

The University of Manchester

Reports available from: <http://eprints.maths.manchester.ac.uk/>

And by contacting: The MIMS Secretary
School of Mathematics
The University of Manchester
Manchester, M13 9PL, UK

ISSN 1749-9097

Bifurcation from stable fixed point to N -dimensional attractor in the border collision normal form

Paul Glendinning

School of Mathematics, University of Manchester, Manchester M13 9PL, U.K.

E-mail: `p.a.glendinning@manchester.ac.uk`

Abstract. The N -dimensional border collision normal form describes bifurcations of piecewise smooth systems. It is shown that there is an open set of parameters such that on one side of the bifurcation the map has a stable fixed point and on the other an attractor with Hausdorff dimension N . For generic parameters this attractor contains open sets and hence has topological dimension equal to N .

PACS numbers: 05.45.-a

Keywords: border collision bifurcation, attractor, piecewise smooth systems, piecewise affine systems, high dimensional attractors

1. Introduction

Piecewise smooth dynamical systems arise in many applications in control theory, the theory of electronic circuits and biological modelling [7]. They are characterized by regions of phase space in which the evolution is smooth separated by switching surfaces across which the evolution rule changes. Local analysis of special cases leads to the derivation of truncated systems called normal forms that capture much of the dynamics close to the situation being described. One such system is the border collision normal form which describes the local behaviour of a generic piecewise smooth map close to a bifurcation value of parameters at which a fixed point intersects a switching surface [18]. In the case of a two dimensional phase space the different changes of behaviour that can occur has been extensively studied [1, 2, 3, 7, 12, 16, 17, 18, 21]. Banerjee et al [3] show that stable periodic orbits or fixed points can be destroyed in the bifurcation, with either new periodic orbits, or chaotic sets with fractal structure being created after the bifurcation value of the parameter. Glendinning shows that a stable fixed point can be destroyed to create a fully two-dimensional attractor on open sets of parameters [14].

Higher dimensional phenomena are less well documented. The general normal form can be written down and it is possible to describe the existence and stability of fixed points [5, 8]. Glendinning and Jeffrey [15] show that these normal forms arise in the bifurcation of continuous time systems, and that after bifurcation there can be attractors

with the same dimension as the ambient phase space [13]. However, those results do not consider the dynamics before the bifurcation, and the examples are such that no stable periodic orbits can exist in the system. The aim of this paper is to show that the same jump of dimension possible in two dimensions described in [14] occurs in general: there are border collision normal forms in \mathbb{R}^N for which the dynamics on one side of the bifurcation is as simple as possible, a stable fixed point, whilst on the other side of the bifurcation there is an attractor with topological dimension equal to N . Moreover, this occurs generically in an open set of parameter values for the normal form.

Let $\mathbf{x} = (x_1, x_2, \dots, x_N)^T$ in \mathbb{R}^N , $N \geq 2$, then a border-collision system is the difference equation

$$\mathbf{x}(t+1) = F(\mathbf{x}(t)) = \begin{cases} A_0 \mathbf{x}(t) + \mathbf{m} & \text{if } x_1(t) \leq 0 \\ A_1 \mathbf{x}(t) + \mathbf{m} & \text{if } x_1(t) \geq 0 \end{cases} \quad (1)$$

where $t \in \mathbb{N}$, A_0 and A_1 are constant matrices, \mathbf{m} is a constant vector, and the system is continuous across the surface $x_1 = 0$. For generic choices of A_0 and A_1 an affine change of coordinates can be used to transform the matrices into observer canonical form [5, 18] and the vector \mathbf{m} to a vector that is zero apart from the first coordinate. In N dimensions this means that generically

$$A_j = \begin{pmatrix} -a_{j,1} & 1 & 0 & \cdots & 0 \\ -a_{j,2} & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & 0 \\ -a_{j,N-1} & 0 & 0 & \cdots & 1 \\ -a_{j,N} & 0 & 0 & \cdots & 0 \end{pmatrix}, \quad \text{and} \quad \mathbf{m} = \begin{pmatrix} \mu \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix} \quad (2)$$

for $j = 0, 1$. The difference equation (1) with constants given by (2) is called the border collision normal form. By a linear change of variables the constant μ can be rescaled to the sign of μ and so in the analysis of the dynamics we may take $\mu = -1$ if $\mu < 0$ and $\mu = 1$ if $\mu > 0$ without loss of generality. The special bifurcation value $\mu = 0$ will not be considered here.

The main result of this paper is the following theorem.

Theorem 1 *There exists an open set $\mathcal{U} \subset \mathbb{R}^{2N}$ such that if $(a_{0,1}, \dots, a_{0,N}, a_{1,1}, \dots, a_{1,N}) \in \mathcal{U}$ then the border collision normal form has a stable fixed point if $\mu < 0$ and an attractor with Hausdorff dimension equal to N if $\mu > 0$. This attractor has topological dimension equal to N generically in \mathcal{U} .*

(A property holds for a generic set of parameters in $\mathcal{U} \subset \mathbb{R}^{2N}$ if it holds for an open, dense subset of \mathcal{U} , thus generic sets are large in the topological sense.) No attempt will be made to optimize the set \mathcal{U} of the theorem, it will be enough to choose the set

$$\begin{aligned} |a_{j,k}| &< \epsilon, \quad j = 0, 1, \quad 1 \leq k < N-1; \\ 2 - \epsilon &< a_{1,N} < 2; \quad \frac{6}{11} < -a_{0,N} < \frac{10}{11}, \end{aligned} \quad (3)$$

for sufficiently small $\epsilon > 0$.




Figure 1. The three dimensional attractor of (1), (2) with $N = 3$ and the parameters of (4) and $\mu = 1$; 50000 iterates are shown from the initial condition $(-0.04, -0.3, -0.1)$.

To illustrate this result consider the normal form in three dimensions with

$$\begin{aligned} \epsilon = 1/400, \quad a_{0,1} = 0.2\epsilon, \quad a_{0,2} = -0.4\epsilon, \quad a_{0,3} = -9/11, \\ a_{1,1} = -0.7\epsilon, \quad a_{1,2} = 0.6\epsilon, \quad a_{1,3} = 2 - 0.5\epsilon, \end{aligned} \quad (4)$$

chosen without a great deal of thought so as to satisfy (3). Then the attractor is shown in Figure 1, and certainly appears three dimensional. The parameters have been chosen to satisfy the constraints of Lemma 1 for the existence of the invariant region of section 3, in particular $\epsilon < 1/(120N)$, and the expansion conditions of section 4 have been verified numerically. Thus, up to the numerical verification, ϵ does seem small enough for Theorem 1 to hold. Attractors that appear to be three dimensional can be found numerically at larger values of ϵ .

As in [14] the main theoretical results used are those of Buzzi [9, 11] and Tsujii [20]. However, in the two-dimensional case it is possible to provide a detailed description of carefully chosen regions and their images. In the higher dimensional cases there is a combinatoric proliferation of boundary points because a hypercube in \mathbb{R}^N has 2^N vertices, and this makes the approach taken in [14] infeasible here. Instead an alternative, and in many ways simpler, strategy is taken.

In section two the simple stable fixed point is shown to exist if $\mu < 0$ and the remaining parameter values are in \mathcal{U} . In section three an invariant region is defined for $\mu > 0$ and in section four a criterion for expansion of a suitable iterate of the border collision normal form is determined. Theorem 1 follows from this expansion result together with the results of Buzzi and Tsujii.

2. The stable fixed point, $\mu < 0$

If $\mu < 0$ then by rescaling variables we may take $\mu = -1$ without loss of generality. If there is a fixed point $(x_1^*, x_2^*, \dots, x_N^*)$ in $x_1 < 0$ with parameters satisfying (3) then it

satisfies

$$-a_{0,1}x_1^* + x_2^* - 1 = x_1^*, \quad x_{k+1}^* = x_k^* + O(\epsilon), \quad k = 2, \dots, N-1, \quad (5)$$

and

$$x_N^* = -a_{0,N}x_1^*. \quad (6)$$

Let $-a_{0,N} = \omega > 0$, then $x_2^* = \omega x_1^* + O(\epsilon)$ and so

$$x_1^* = -\frac{1}{1-\omega} + O(\epsilon).$$

Thus provided ω is less than one the fixed point exists for sufficiently small ϵ . The stability is determined from the eigenvalues of A_0 , i.e. the roots of the characteristic polynomial

$$s^N + \sum_{r=0}^{N-1} a_{0,N-r} s^r = 0.$$

If $\epsilon = 0$ the characteristic equation is $s^N = \omega$ and so there are N roots with modulus equal to $\omega^{\frac{1}{N}}$ which is less than one. Hence for sufficiently small $\epsilon > 0$ the roots are all within the unit circle and the fixed point is stable.

3. An invariant region, $\mu > 0$

The invariant region derived below is motivated by previous work [13, 15] and it is the existence of this region combined with the expansion of the next section that determined the choice of parameters defining \mathcal{U} in (refUeps). Consider the rectangular region in \mathbb{R}^N defined by

$$-(1+b_1) \leq x_1 \leq 1+a_1, \quad -(2+b_k) \leq x_k \leq a_k, \quad k = 2, \dots, N. \quad (7)$$

with constants (a_k) and (b_k) positive and less than one. Then if $\mu = 1$ (which may be chosen without loss of generality if $\mu > 0$) the images $(x'_1, x'_2, \dots, x'_N)$ of points in this region satisfy a set of inequalities that can be used to show that the constants can be chosen such that (7) defines an invariant region provided ϵ is sufficiently small in the definitions of parameters (3).

If $x_1 < 0$ then $x'_1 = -a_{0,1}x_1 + x_2 + 1$ and so

$$\begin{aligned} \min\{a_{0,1}(1+b_1), 0\} - (2+b_2) + 1 &\leq x'_1 \\ &\leq \max\{a_{0,1}(b_2+1), 0\} + a_2 + 1 \end{aligned}$$

and similarly if $x_1 > 0$ then

$$\begin{aligned} \min\{-a_{1,1}(1+a_1), 0\} - (2+b_2) + 1 &\leq x'_1 \\ &\leq \max\{-a_{1,1}(a_1+1), 0\} + a_2 + 1. \end{aligned}$$

Hence since $|a_{j,k}| < \epsilon$, $j = 0, 1$, $k = 1, \dots, N-1$, by (3)

$$\begin{aligned} \min\{-\epsilon(1+b_1), -\epsilon(1+a_1)\} - (1+b_2) &\leq x'_1 \\ &\leq \max\{\epsilon(1+b_1), \epsilon(1+a_1)\} + 1 + a_2. \end{aligned} \quad (8)$$

Similarly, if $k \in \{2, \dots, N-1\}$

$$\begin{aligned} \min\{-\epsilon(1+b_1), -\epsilon(1+a_1)\} - (2+b_{k+1}) &\leq x'_k \\ &\leq \max\{\epsilon(1+b_1), \epsilon(1+a_1), 0\} + a_{k+1}. \end{aligned} \quad (9)$$

Finally, consider x'_N . If $x_1 < 0$ and $\omega = -a_{0,N} > 0$ then

$$-\omega(1+b_1) \leq x'_N \leq 0 \quad (10)$$

whilst if $x_1 > 0$ then

$$-2(1+a_1) \leq x'_N \leq 0 \quad (11)$$

where we have used the assumption in (3) that $2 - \epsilon < a_{1,N} < 2$.

Comparison of (11) and (7) with $k = N$ and $\epsilon \rightarrow 0$ shows that the region defined by (7) is invariant for sufficiently small ϵ if $2a_1 < b_N$ and, using (8) and (9), $a_{k+1} < a_k$ and $b_{k+1} < b_k$. This motivates the following lemma. Again, no attempt has been made to optimize the sizes of the regions, the important feature is that the criteria are satisfied on an open set of parameter values.

Lemma 1 *Let*

$$a_k = 2(N+1-k)\epsilon \quad \text{and} \quad b_k = \frac{1}{20} - 2k\epsilon, \quad (12)$$

$k = 1, \dots, N$. If $\epsilon > 0$ is sufficiently small then (7) is an invariant region for the border collision normal form with $\mu = 1$ and coefficients satisfying (3).

Proof: The proof is by direct verification using equations (7) to (11). Choose $\epsilon > 0$ so that $b_1 > b_2 > \dots > b_N > 2a_1$, i.e.

$$0 < \epsilon < \frac{1}{120N}.$$

Then $b_1 > a_1$ and so the maximum modulus of $a_{j,k}(1+b_1)$ and $a_{j,k}(1+a_1)$ is $\epsilon(1+b_1)$ if $k \neq N$. Thus for x'_1 defined by (8) to be in the interval defined in (7)

$$1+b_1 > 1+b_2 + \epsilon(1+b_1), \quad 1+a_1 > 1+a_2 + \epsilon(1+b_1)$$

and since

$$b_1 - b_2 = a_1 - a_2 = 2\epsilon > (1+b_1)\epsilon$$

both these inequalities hold.

Similarly for the coordinates x'_k , $k = 2, \dots, N-1$, the interval of possible values for the image is contained in the original region provided

$$2+b_j > 2+b_{j+1} + \epsilon(1+b_1) < 2, \quad a_j > a_{j+1} + \epsilon(1+b_1)$$

which hold for the same reason. Since $b_1\omega < 1$ and $b_N > 0$, $2+b_N > \omega(1+b_1)$ so the iterate defined by (10) is in the defined region, and finally from (11), the region is invariant if

$$2+b_N > 2(1+a_1),$$

i.e. $b_N > 2a_1$ as noted earlier.

□

4. Expansion

Piecewise affine expanding maps are maps defined on polyhedral regions (regions bounded by hyperplanes) such that in each region the map is expanding in some metric. General results of Buzzi [9, 11] and Tsujii [20] show that if a piecewise affine map is defined on a bounded polyhedral region in \mathbb{R}^N then there is an attractor with an invariant measure which is absolutely continuous with respect to Lebesgue measure [11, 20] and hence has Hausdorff dimension equal to N . Moreover, for generic piecewise expanding maps the attractor contains open sets [9, 11]. The border collision normal form is not expanding in the Euclidean metric, although there are equivalent metrics for which induced maps are expanding. This creates some extra technical difficulties in [13, 14], and by using an appropriate induced map it is shown below that these difficulties can be avoided, making it possible to work with the Euclidean metric throughout the analysis.

The structure of the normal form matrices A_0 and A_1 of (2) is such that for small ϵ in (3) and high enough iterates of the map, expansion in the Euclidean metric is relatively straightforward. Let (α_r) be any countable set of real numbers, $r \in \mathbb{N}$, and let $M(r)$ denote the $N \times N$ matrix

$$M(r)_{ij} = \begin{cases} 1 & \text{if } j = i + 1, i = 1, \dots, N - 1, \\ \alpha_r & \text{if } i = N, j = 1 \\ 0 & \text{otherwise} \end{cases}$$

then an elementary manipulation shows that the product

$$M(N) \dots M(1) = \text{diag} (\alpha_1, \dots, \alpha_N)$$

and hence

$$M(2N)M(2N-1) \dots M(2)M(1) = \text{diag} (\alpha_1\alpha_{N+1}, \dots, \alpha_N\alpha_{2N}). \quad (13)$$

If $\epsilon = 0$ then A_0 and A_1 are matrices of the form $M(r)$ with $\alpha_r = -a_{0,N}$ or $\alpha_r = -a_{1,N}$. Since $a_{1,N}^2 > (2 - \epsilon)^2$ is close to 4 and $|a_{0,N}a_{1,N}|$ is greater than $\frac{12}{11}$ to order ϵ for parameters satisfying (3), the only products of $2N$ matrices A_0 and A_1 which can produce contracting diagonal terms are those with A_0 as both the i^{th} term and the $(N + i)^{\text{th}}$ term of the product. The next lemma shows that this cannot happen for the parameters chosen here.

Lemma 2 *Consider the border collision normal form in \mathbb{R}^N with $\mu = 1$ and parameters satisfying (3) with $\epsilon > 0$ sufficiently small. If \mathbf{x} lies in the invariant region of Lemma 1 and $x_1 < 0$ then $F^N(\mathbf{x})$ lies in $x_1 > 0$.*

Proof: Let $\pi_k \mathbf{x}$ denote the k^{th} coordinate of \mathbf{x} . Then $x_1 = \pi_1(\mathbf{x}) < 0$ implies that $\pi_N(F(\mathbf{x})) = \omega x_1 < 0$, with $\omega = -a_{0,N} > 0$. Then $\pi_{N-1}(F^2(\mathbf{x})) = \omega x_1 + O(\epsilon)$ and continuing up the orbit $\pi_1(F^N(\mathbf{x})) = \omega x_1 + 1 + O(\epsilon)$. Since $x_1 \in (-\frac{21}{20}, 0)$ by Lemma 1, $|\omega x_1| < \frac{10}{11} \frac{21}{20} = \frac{21}{22}$, and so if ϵ is sufficiently small so that the order ϵ terms are less than $\frac{1}{20}$ in magnitude, $\pi_1(F^N(\mathbf{x})) > 0$.

□

Now consider the iterated map $G = F^{2N}$. G is itself a piecewise smooth map on domains that are polyhedral (as they are the preimages of polyhedral regions). The linear part of each component of F^{2N} is a product of matrices of the form $M(r) + O(\epsilon)$ with $\alpha_r \in \{\omega, -2\}$ and hence the linear parts are all ϵ -close to diagonal matrices with coefficients in $\{4, -2\omega\}$ as the third possibility, ω^2 cannot occur at parameters defined in (3) by Lemma 2. Hence each linear part is expanding in the Euclidean metric for sufficiently small $\epsilon > 0$ and so the results of Buzzi and Tsujii hold for G , so G has an absolutely continuous invariant measure for all parameter values in \mathcal{U} and the support of this measure contains open sets generically. de Melo and van Strien [4] (see also [14]) show that this implies that the original map F also has an absolutely continuous invariant measure for all parameter values in \mathcal{U} and it follows from Buzzi's result that these contain open sets generically [11, 14]. This completes the proof of the main theorem, Theorem 1, of this paper.

5. Conclusion

Theorem 1 shows that the two-dimensional result of [14] holds in higher dimensions. In particular, there is a transition from a stable fixed point to an attractor with the highest possible dimension in the ambient phase space for a generic subset of an open set of parameters. This poses questions about how complicated a bifurcation theorem for piecewise smooth system might be. For example, is it possible to find parameters for which there is a transition from a stable fixed point to an attractor with Hausdorff dimension d for every $d \in [0, N]$? There is also a danger that such bifurcation theorems will become little more than (long) lists of possibilities.

One thing is clear: the fact that this bifurcation occurs on open sets of parameter values shows that it is not exceptional, and that dimension reduction techniques will have limited application to certain piecewise smooth systems even if there are stable fixed points at some nearby parameter values.

The border collision normal form can be derived close to grazing-sliding bifurcations of periodic orbits, though in this case it is a lowest order truncation of the dynamical system [6, 15], so an important question is whether the existence of N -dimensional attractors in the truncated, piecewise linear, map implies the existence of N -dimensional attractors in the full map. There are results [10, 19] for piecewise real analytic maps, but the derivation of the border collision normal form from grazing-sliding bifurcations involves non-analytic terms and hence these results cannot be applied. Resolving this problem is important in assessing the implications of the results shown here for applications.

Finally the proof of Buzzi [9, 10] that only generic expanding maps have open sets (and hence full topological dimension) does not show that at the remaining set of parameters the attractor does not contain open sets. Thus the restriction of full topological dimension to a generic set of parameters may be an artefact of the method of proof, and attractors may have topological dimension N for all parameters in \mathcal{U} .

References

- [1] Banerjee S and Grebogi C 1999 Border Collision Bifurcations in Two-Dimensional Piecewise Smooth Maps, *Phys. Rev. E*, **59** 4052–4061
- [2] Banerjee S, Ranjan P and Grebogi C 2000 Bifurcations in two-dimensional piecewise smooth maps — theory and applications in switching circuits, *IEEE Trans. Circ. Syst., I: Fundam. Theory Appl.*, **47** 633–643.
- [3] Banerjee S, Yorke JA and Grebogi C 1998 Robust Chaos *Phys. Rev. Lett.* **80** 3049–3052
- [4] de Melo W. and van Strien S 2003 *One-Dimensional Dynamics*, Springer, Berlin.
- [5] di Bernardo M 2003 Normal forms of border collision in high dimensional non-smooth maps, *Proceedings IEEE ISCAS 2003*, **3**, 76–79
- [6] di Bernardo M, Budd CJ and Champneys AR 2001 Grazing and Border-Collision in Piecewise-Smooth Systems: A Unifed Analytical Framework *Phys. Rev. Lett.* **86** 2554–2556
- [7] di Bernardo M, Budd CJ, Champneys AR and Kowalczyk P 2008 *Piecewise-smooth Dynamical Systems: Theory and Applications* Springer, London
- [8] di Bernardo M, Montanaro U, and Santini S 2011 Canonical Forms of Generic Piecewise Linear Continuous Systems, *IEEE Trans. Automatic Control* **56** 1911–1915
- [9] Buzzi J 1999 Absolutely continuous invariant measures for generic multi-dimensional piecewise affine expanding maps, *Int. J. Bifn. & Chaos* **9** 1743–1750
- [10] Buzzi J 2000 A.C.I.M.s for arbitrary expanding piecewise \mathbb{R} -analytic mappings of the plane, *Ergod. Thy. & Dyn. Syst.* **20** 697–708
- [11] Buzzi J 2001 Thermodynamic formalism for piecewise invertible maps: Absolutely continuous invariant measures as equilibrium states, in *Smooth Ergodic Theory and Its Applications*, eds. A Katok, R de la Llave, Y Pesin and H Weiss, AMS Proc. Symp. Pure Math **69**, 749–784
- [12] Gardini L 1992 Some global bifurcations of two-dimensional endomorphisms by use of critical lines *Nonl. Anal.* **18** 361–399
- [13] Glendinning P 2014 Attractors with dimension n for open sets of parameter space in the n -dimensional border collision normal form, *Int. J. Bif. & Chaos* **24** 1450164
- [14] Glendinning P 2015 Bifurcation from stable fixed point to two-dimensional attractor in the border collision normal form, preprint.
- [15] Glendinning P and Jeffrey M 2015 Grazing-sliding bifurcations, border collision maps and the curse of dimensionality for piecewise smooth bifurcation theory *Nonlinearity* **28** 263–283
- [16] Mira C, Gardini L, Barugola A and Cathala JC 1996 *Chaotic Dynamics in Two-Dimensional Noninvertible Maps*, World Scientific, Singapore.
- [17] Nusse HE, Ott E and Yorke JA 1994 Border-collision bifurcations: An explanation for observed bifurcation phenomena, *Phys. Rev. E*, **49** 1073–1076
- [18] Nusse HE and Yorke JA 1992 Border-collision bifurcation including ‘period two to period three’ for piecewise smooth systems, *Physica D*, **57** 39–57
- [19] Tsujii M 2000 Absolutely continuous invariant measures for piecewise real-analytic expanding maps on the plane, *Commun. Math. Phys.* **208** 605–622
- [20] Tsujii M 2001 Absolutely continuous invariant measures for expanding piecewise linear maps, *Invent. Math.* **143** 349–373
- [21] Young LS 1985 Bowen-Ruelle measures for certain piecewise hyperbolic maps, *Trans. Amer. Math. Soc.* **287** 41–48