

*Geometric structure in the representation theory
of p -adic groups II*

Aubert, Anne-Marie and Baum, Paul and Plymen, Roger

2010

MIMS EPrint: **2010.68**

Manchester Institute for Mathematical Sciences
School of Mathematics

The University of Manchester

Reports available from: <http://eprints.maths.manchester.ac.uk/>

And by contacting: The MIMS Secretary
School of Mathematics
The University of Manchester
Manchester, M13 9PL, UK

ISSN 1749-9097

Geometric structure in the representation theory of reductive p -adic groups II

Anne-Marie Aubert, Paul Baum, and Roger Plymen

1. Introduction

This expository note will state the ABP (Aubert-Baum-Plymen) conjecture [2, 3, 4]. The conjecture can be stated at four levels:

- K -theory of C^* algebras
- Periodic cyclic homology of finite type algebras
- Geometric equivalence of finite type algebras
- Representation theory

The emphasis in this note will be on representation theory. The first two items in the above list are topological, and the third item is algebraic. Validity for the two topological items is quite plausible, and thus gives some credibility to the representation theory version of the conjecture. A recent result of M. Solleveld [48], when combined with results of [13, 14, 18, 19, 20, 21, 23, 29, 30, 38, 40, 41, 42, 43, 44, 46] proves a very substantial part of the conjecture for many examples.

2. Bernstein Components

Let G be a reductive p -adic group. Examples are $GL(n, F)$, $SL(n, F)$ etc. where F is a local nonarchimedean field, that is, F is a finite extension of the p -adic numbers \mathbb{Q}_p , or the local function field $\mathbb{F}_q((x))$.

Definition. A *representation* of G is a group homomorphism

$$\phi : G \rightarrow \text{Aut}_{\mathbb{C}}(V)$$

where V is a vector space over the complex numbers \mathbb{C} .

The local field F in its natural topology is a locally compact and totally disconnected topological field. Hence G (in its p -adic topology) is a locally compact and totally disconnected topological group.

2010 *Mathematics Subject Classification.* 20G05, 22E50.

Key words and phrases. reductive p -adic group, representation theory, geometric structure, affine Hecke algebras.

The second author was partially supported by NSF grant DMS-0701184.

Definition. A representation

$$\phi : G \rightarrow \text{Aut}_{\mathbb{C}}(V)$$

of G is *smooth* if for every $v \in V$,

$$G_v = \{g \in G \mid \phi(g)v = v\}$$

is an open subgroup of G .

The smooth (or admissible) dual of G , denoted \widehat{G} , is the set of equivalence classes of smooth irreducible representations of G .

$$\widehat{G} = \{\text{Smooth irreducible representations of } G\} / \sim$$

Problem. Describe \widehat{G} .

Since G is locally compact we may fix a (left-invariant) Haar measure dg for G . The Hecke algebra of G , denoted $\mathcal{H}G$, is then the convolution algebra of all locally-constant compactly-supported complex-valued functions $f : G \rightarrow \mathbb{C}$.

$$\begin{aligned} (f + h)(g) &= f(g) + h(g) \\ (f * h)(g_0) &= \int_G f(g)h(g^{-1}g_0)dg \end{aligned} \quad \left\{ \begin{array}{l} g \in G \\ g_0 \in G \\ f \in \mathcal{H}G \\ h \in \mathcal{H}G \end{array} \right.$$

Definition. A *representation* of the Hecke algebra $\mathcal{H}G$ is a homomorphism of \mathbb{C} algebras

$$\psi : \mathcal{H}G \rightarrow \text{End}_{\mathbb{C}}(V)$$

where V is a vector space over the complex numbers \mathbb{C} .

Definition. A representation

$$\psi : \mathcal{H}G \rightarrow \text{End}_{\mathbb{C}}(V)$$

of the Hecke algebra $\mathcal{H}G$ is *irreducible* if ψ is not the zero map and \nexists a vector subspace W of V such that W is preserved by the action of $\mathcal{H}G$ and $\{0\} \neq W \neq V$.

Definition. A *primitive ideal* I in $\mathcal{H}G$ is the null space of an irreducible representation of $\mathcal{H}G$. Note that $\mathcal{H}G$ itself is not a primitive ideal.

Thus

$$0 \longrightarrow I \longrightarrow \mathcal{H}G \xrightarrow{\psi} \text{End}_{\mathbb{C}}(V)$$

is exact where ψ is an irreducible representation of $\mathcal{H}G$.

There is a (canonical) bijection of sets

$$\widehat{G} \longleftrightarrow \text{Prim}(\mathcal{H}G)$$

where $\text{Prim}(\mathcal{H}G)$ is the set of primitive ideals in $\mathcal{H}G$.

What has been gained from this bijection?

On $\text{Prim}(\mathcal{H}G)$ there is a topology—the Jacobson topology.

If S is a subset of $\text{Prim}(\mathcal{H}G)$ then the closure \overline{S} (in the Jacobson topology) of S is

$$\overline{S} = \{J \in \text{Prim}(\mathcal{H}G) \mid J \supset \bigcap_{I \in S} I\}$$

$\text{Prim}(\mathcal{H}G)$ (with the Jacobson topology) is the disjoint union of its connected components.

$\pi_o\text{Prim}(\mathcal{H}G)$ denotes the set of connected components of $\text{Prim}(\mathcal{H}G)$.

$\pi_o\text{Prim}(\mathcal{H}G)$ is a countable set and has no further structure.

$\pi_o\text{Prim}(\mathcal{H}G)$ is also known as the *Bernstein spectrum* of G .

$\pi_o\text{Prim}(\mathcal{H}G) = \{(M, \sigma)\} / \sim$ where (M, σ) is a cuspidal pair i.e. M is a Levi factor of a parabolic subgroup P of G and σ is an irreducible super-cuspidal representation of M . \sim is the conjugation action of G , combined with tensoring by unramified characters of M . Thus $(M, \sigma) \sim (M', \sigma')$ iff \exists an unramified character $\psi: M \rightarrow \mathbb{C} - \{0\}$ of M and an element g of G , $g \in G$, with

$$g(M, \psi \otimes \sigma) = (M', \sigma')$$

The meaning of this equality is:

- $gMg^{-1} = M'$
- $g_*(\psi \otimes \sigma)$ and σ' are equivalent smooth irreducible representations of M' .

For each $\mathfrak{s} \in \pi_o\text{Prim}(\mathcal{H}G)$, $\widehat{G}_{\mathfrak{s}}$ denotes the connected component of $\text{Prim}(\mathcal{H}G)$.

Using the canonical bijections

$$\widehat{G}_{\mathfrak{s}} \longleftrightarrow \text{Prim}(\mathcal{H}G) \quad \pi_o\text{Prim}(\mathcal{H}G) \longleftrightarrow \{(M, \sigma)\} / \sim$$

$\widehat{G}_{\mathfrak{s}}$ is obtained by fixing (M, σ) and then taking the irreducible constituents of $\text{Ind}_M^G(\psi \otimes \sigma)$ where Ind_M^G is (smooth) parabolic induction and ψ can be any unramified character of M .

The connected components of $\text{Prim}(\mathcal{H}G)$ are also known as the Bernstein components. The problem of describing \widehat{G} now breaks up into two problems.

Problem 1: Describe $\pi_o\text{Prim}(\mathcal{H}G)$.

Problem 2: For each $\mathfrak{s} \in \pi_o\text{Prim}(\mathcal{H}G)$ describe the Bernstein component $\widehat{G}_{\mathfrak{s}}$.

In this note we shall be concerned with Problem 2. Problem 1 involves describing the irreducible super-cuspidal representations of Levi subgroups of G . The basic conjecture on this issue is that if M is a reductive p -adic group (e.g. M is a Levi factor of a parabolic subgroup of G) then any irreducible super-cuspidal representation of M is obtained by smooth induction from an irreducible representation of a subgroup of M which is compact modulo the center of M . This basic conjecture is now known to be true to a very great extent [25] [49]. For Problem 2 the ABP conjecture proposes that each Bernstein component $\widehat{G}_{\mathfrak{s}}$ has a very simple geometric structure.

3. Infinitesimal Character

Notation. \mathbb{C}^\times denotes the (complex) affine variety $\mathbb{C} - \{0\}$.

Definition. A *complex torus* is a (complex) affine variety T such that there exists an isomorphism of affine varieties

$$T \cong \mathbb{C}^\times \times \mathbb{C}^\times \times \cdots \times \mathbb{C}^\times.$$

Bernstein [11] assigns to each $\mathfrak{s} \in \pi_o \text{Prim}(\mathcal{H}G)$ a complex torus $T_{\mathfrak{s}}$ and a finite group $W_{\mathfrak{s}}$ acting on $T_{\mathfrak{s}}$. Bernstein's construction can be recalled as follows. First, fix (M, σ) . M^0 denotes the subgroup of M consisting of all $g \in M$ such that:

$$\text{whenever } \varphi: M \rightarrow F^{\times} \text{ is an algebraic character, } \varphi(g) \in \mathcal{O}_F^{\times}.$$

Here F is the p-adic field over which G is defined, \mathcal{O}_F is the integers in F and \mathcal{O}_F^{\times} is the invertible elements of the ring \mathcal{O}_F . Equivalently, M^0 is the (closed) normal subgroup of M generated by all the compact subgroups of M . The quotient group M/M^0 is discrete and is a free abelian group of finite rank. Therefore $\text{Hom}_{\mathbb{Z}}(M/M^0, \mathbb{C}^{\times})$ is a complex torus. The points of this complex torus are (by definition) the unramified characters of M . Within this complex torus consider all the unramified characters φ of M such that:

$$\varphi \otimes \sigma \text{ is equivalent (as an irreducible smooth representation of } M) \text{ to } \sigma.$$

Denote this set of characters by I_{σ} . I_{σ} is a finite subgroup of $\text{Hom}_{\mathbb{Z}}(M/M^0, \mathbb{C}^{\times})$ so the quotient group $\text{Hom}_{\mathbb{Z}}(M/M^0, \mathbb{C}^{\times})/I_{\sigma}$ is a complex torus, and this is Bernstein's torus $T_{\mathfrak{s}}$.

$$T_{\mathfrak{s}} = \text{Hom}_{\mathbb{Z}}(M/M^0, \mathbb{C}^{\times})/I_{\sigma}$$

Denote $\text{Hom}_{\mathbb{Z}}(M/M^0, \mathbb{C}^{\times})$ by $T'_{\mathfrak{s}}$. Denote the quotient map $T'_{\mathfrak{s}} \rightarrow T_{\mathfrak{s}}$ by

$$\eta: T'_{\mathfrak{s}} \longrightarrow T_{\mathfrak{s}}$$

The Weyl group W_M of M is $N_G(M)/M$, where $N_G(M)$ is the normalizer of M in G . $W_{\mathfrak{s}}$ is the subgroup of W_M consisting of all $w \in W_M$ such that:

$$\text{given } w, \exists \varphi_w \in T'_{\mathfrak{s}} = \text{Hom}_{\mathbb{Z}}(M/M^0, \mathbb{C}^{\times}) \text{ with } \varphi_w \otimes \sigma \text{ equivalent to } w_*(\sigma)$$

($\varphi_w \otimes \sigma$ is equivalent as an irreducible smooth representation of M to $w_*(\sigma)$). If I_{σ} is not the trivial one-element group, then $w \mapsto \varphi_w$ is not well-defined as a map from $W_{\mathfrak{s}}$ to $T'_{\mathfrak{s}}$. However, when composed with the quotient map $T'_{\mathfrak{s}} \rightarrow T_{\mathfrak{s}}$, a well-defined group homomorphism $W_{\mathfrak{s}} \rightarrow T_{\mathfrak{s}}$ is obtained:

$$\begin{aligned} W_{\mathfrak{s}} &\longrightarrow T_{\mathfrak{s}} \\ w &\longmapsto \eta(\varphi_w) \end{aligned}$$

The evident conjugation action of W_M on M gives an action of W_M on $T'_{\mathfrak{s}} = \text{Hom}_{\mathbb{Z}}(M/M^0, \mathbb{C}^{\times})$. When restricted to $W_{\mathfrak{s}}$ this gives an action of $W_{\mathfrak{s}}$ on $T'_{\mathfrak{s}}$ which preserves I_{σ} and hence $W_{\mathfrak{s}}$ acts on the quotient torus $T_{\mathfrak{s}}$. Note that at this point $W_{\mathfrak{s}}$ is acting as automorphisms of the algebraic group $T_{\mathfrak{s}}$. Denote this action by:

$$wx = w_*(x) \quad w \in W_{\mathfrak{s}} \quad x \in T_{\mathfrak{s}}$$

In Bernstein's action of $W_{\mathfrak{s}}$ on $T_{\mathfrak{s}}$, $w \in W_{\mathfrak{s}}$ acts by

$$x \longmapsto w_*(x)[\eta(\varphi_w)] \quad x \in T_{\mathfrak{s}}$$

i.e. when $w \in W_{\mathfrak{s}}$ is applied to $x \in T_{\mathfrak{s}}$, the result is the product (using the group structure of $T_{\mathfrak{s}}$) of $w_*(x)$ and $\eta(\varphi_w)$. Hence $W_{\mathfrak{s}}$ is acting as automorphisms of the affine variety $T_{\mathfrak{s}}$ — but not necessarily as automorphisms of the algebraic group $T_{\mathfrak{s}}$.

Consider the quotient variety $T_{\mathfrak{s}}/W_{\mathfrak{s}}$. Define a surjective map $\pi_{\mathfrak{s}}$ mapping $\hat{G}_{\mathfrak{s}}$ onto $T_{\mathfrak{s}}/W_{\mathfrak{s}}$

$$\begin{array}{c} \widehat{G}_{\mathfrak{s}} \\ \downarrow \pi_{\mathfrak{s}} \\ T_{\mathfrak{s}}/W_{\mathfrak{s}} \end{array}$$

by:

given $\zeta \in \widehat{G}_{\mathfrak{s}}$ select $\varphi \in T'_{\mathfrak{s}}$ such that ζ is an irreducible constituent of $\text{Ind}_M^G(\varphi \otimes \sigma)$.

Then set $\pi_{\mathfrak{s}}(\zeta) = \eta(\varphi)$.

This map $\pi_{\mathfrak{s}}$ is referred to as the *infinitesimal character* or the *central character*. In Bernstein's work $\widehat{G}_{\mathfrak{s}}$ is a set (i.e. is only a set) so $\pi_{\mathfrak{s}}$

$$\begin{array}{c} \widehat{G}_{\mathfrak{s}} \\ \downarrow \pi_{\mathfrak{s}} \\ T_{\mathfrak{s}}/W_{\mathfrak{s}} \end{array}$$

is a map of sets.

$\pi_{\mathfrak{s}}$ is surjective, finite-to-one and generically one-to-one; *generically one-to-one* means that there is a sub-variety $\mathfrak{R}_{\mathfrak{s}}$ of $T_{\mathfrak{s}}/W_{\mathfrak{s}}$ such that the pre-image of each point in $T_{\mathfrak{s}}/W_{\mathfrak{s}} - \mathfrak{R}_{\mathfrak{s}}$ consists of just one point. $\mathfrak{R}_{\mathfrak{s}}$ is the subvariety of reducibility, i.e. $\mathfrak{R}_{\mathfrak{s}}$ is given by those unramified twists of σ such that there is reducibility when parabolically induced to G .

Remark. Let $\Delta_{\mathfrak{s}}$ be the maximal compact subgroup of $T_{\mathfrak{s}}$. As above, the Bernstein action of $w \in W_{\mathfrak{s}}$ on $T_{\mathfrak{s}}$ is given by:

$$x \mapsto w_*(x)[\theta(\varphi_w)] \quad x \in T_{\mathfrak{s}}.$$

This implies that $\Delta_{\mathfrak{s}}$ is preserved by the action.

4. Extended Quotient

Let Γ be a finite group acting on an affine variety X as automorphisms of the affine variety

$$\Gamma \times X \rightarrow X.$$

The quotient variety X/Γ is obtained by collapsing each orbit to a point. For $x \in X$, Γ_x denotes the stabilizer group of x :

$$\Gamma_x = \{\gamma \in \Gamma \mid \gamma x = x\}.$$

$c(\Gamma_x)$ denotes the set of conjugacy classes of Γ_x . The extended quotient is obtained by replacing the orbit of x by $c(\Gamma_x)$. This is done as follows:

Set $\widetilde{X} = \{(\gamma, x) \in \Gamma \times X \mid \gamma x = x\} \subset \Gamma \times X$. \widetilde{X} is an affine variety and is a sub-variety of $\Gamma \times X$. Γ acts on \widetilde{X} :

$$\Gamma \times \widetilde{X} \rightarrow \widetilde{X} \quad \alpha(\gamma, x) = (\alpha\gamma\alpha^{-1}, \alpha x), \quad \alpha \in \Gamma, \quad (\gamma, x) \in \widetilde{X}.$$

The extended quotient, denoted $X//\Gamma$, is \widetilde{X}/Γ . Thus the extended quotient $X//\Gamma$ is the ordinary quotient for the action of Γ on \widetilde{X} . In a quite straightforward way $X//\Gamma$ is an affine variety.

The projection $\tilde{X} \rightarrow X, (\gamma, x) \mapsto x$ passes to quotient spaces to give a morphism of affine varieties

$$\rho: X//\Gamma \rightarrow X/\Gamma.$$

This map will be referred to as the projection of the extended quotient onto the ordinary quotient.

The inclusion

$$\begin{aligned} X &\hookrightarrow \tilde{X} \\ x &\mapsto (e, x) \quad e = \text{identity element of } \Gamma \end{aligned}$$

passes to quotient spaces to give an inclusion $X/\Gamma \hookrightarrow X//\Gamma$. This will be referred to as the inclusion of the ordinary quotient in the extended quotient. Using this inclusion, $X//\Gamma - X/\Gamma$ denotes $X//\Gamma$ with X/Γ removed.

5. ABP Conjecture Part 1

As above, G is a reductive p-adic group and \mathfrak{s} is a point in the Bernstein spectrum of G .

Consider the two maps indicated by vertical arrows:

$$\begin{array}{ccc} T_{\mathfrak{s}}//W_{\mathfrak{s}} & & \widehat{G}_{\mathfrak{s}} \\ \downarrow \rho_{\mathfrak{s}} & \text{and} & \downarrow \pi_{\mathfrak{s}} \\ T_{\mathfrak{s}}/W_{\mathfrak{s}} & & T_{\mathfrak{s}}/W_{\mathfrak{s}} \end{array}$$

$\pi_{\mathfrak{s}}$ is the infinitesimal character and $\rho_{\mathfrak{s}}$ is the projection of the extended quotient on the ordinary quotient. In practice, $T_{\mathfrak{s}}//W_{\mathfrak{s}}$ and $\rho_{\mathfrak{s}}$ are much easier to calculate than $\widehat{G}_{\mathfrak{s}}$ and $\pi_{\mathfrak{s}}$.

The maps $\rho_{\mathfrak{s}}$ and $\pi_{\mathfrak{s}}$ are conceptually quite different; nevertheless, we conjecture that one can pass from one to the other, via a simple algebraic correction, and, in so doing, predict the number of inequivalent irreducible constituents in a given parabolically induced representation of G . The precise conjecture (ABP Part 1) consists of two statements.

Conjecture.

- (1) *The infinitesimal character*

$$\pi_{\mathfrak{s}}: \widehat{G}_{\mathfrak{s}} \rightarrow T_{\mathfrak{s}}/W_{\mathfrak{s}}$$

is one-to-one if and only if the action of $W_{\mathfrak{s}}$ on $T_{\mathfrak{s}}$ is free.

- (2) *There exists a bijection*

$$\mu_{\mathfrak{s}}: T_{\mathfrak{s}}//W_{\mathfrak{s}} \longleftrightarrow \widehat{G}_{\mathfrak{s}}$$

with the following five properties:

Notation for Property 1:

$$\mathfrak{s} \in \pi_0 \text{Prim}(\mathcal{H}G)$$

Within the admissible dual \widehat{G} have the tempered dual $\widehat{G}_{\text{tempered}}$.

$$\widehat{G}_{\text{tempered}} = \{\text{smooth tempered irreducible representations of } G\} / \sim$$

$\widehat{G}_{tempered} = \text{Support of the Plancherel measure}$
 $\Delta_{\mathfrak{s}} = \text{maximal compact subgroup of } T_{\mathfrak{s}}.$
 $\Delta_{\mathfrak{s}}$ is a compact torus. The action of $W_{\mathfrak{s}}$ on $T_{\mathfrak{s}}$ preserves the maximal compact subgroup $\Delta_{\mathfrak{s}}$, so can form the compact orbifold $\Delta_{\mathfrak{s}}//W_{\mathfrak{s}}$.

Property 1 of the bijection $\mu_{\mathfrak{s}}$

- The bijection $\mu_{\mathfrak{s}} : T_{\mathfrak{s}}//W_{\mathfrak{s}} \longleftrightarrow \widehat{G}_{\mathfrak{s}}$ maps $\Delta_{\mathfrak{s}}//W_{\mathfrak{s}}$ onto $\widehat{G}_{\mathfrak{s}} \cap \widehat{G}_{tempered}$
 $\Delta_{\mathfrak{s}}//W_{\mathfrak{s}} \longleftrightarrow \widehat{G}_{\mathfrak{s}} \cap \widehat{G}_{tempered}$

Property 2 of the bijection $\mu_{\mathfrak{s}}$

- For many \mathfrak{s} the diagram

$$\begin{array}{ccc}
 T_{\mathfrak{s}}//W_{\mathfrak{s}} & \xrightarrow{\mu_{\mathfrak{s}}} & \widehat{G}_{\mathfrak{s}} \\
 \rho_{\mathfrak{s}} \downarrow & & \downarrow \pi_{\mathfrak{s}} \\
 T_{\mathfrak{s}}/W_{\mathfrak{s}} & \xrightarrow{I} & T_{\mathfrak{s}}/W_{\mathfrak{s}}
 \end{array}$$

does not commute. ($I =$ the identity map of $T_{\mathfrak{s}}/W_{\mathfrak{s}}$.)

Property 3 of the bijection $\mu_{\mathfrak{s}}$

- In the possibly non-commutative diagram

$$\begin{array}{ccc}
 T_{\mathfrak{s}}//W_{\mathfrak{s}} & \xrightarrow{\mu_{\mathfrak{s}}} & \widehat{G}_{\mathfrak{s}} \\
 \rho_{\mathfrak{s}} \downarrow & & \downarrow \pi_{\mathfrak{s}} \\
 T_{\mathfrak{s}}/W_{\mathfrak{s}} & \xrightarrow{I} & T_{\mathfrak{s}}/W_{\mathfrak{s}}
 \end{array}$$

the bijection $\mu_{\mathfrak{s}} : T_{\mathfrak{s}}//W_{\mathfrak{s}} \longrightarrow \widehat{G}_{\mathfrak{s}}$ is continuous where $T_{\mathfrak{s}}//W_{\mathfrak{s}}$ has the Zariski topology and $\widehat{G}_{\mathfrak{s}}$ has the Jacobson topology

AND the composition

$$\pi_{\mathfrak{s}} \circ \mu_{\mathfrak{s}} : T_{\mathfrak{s}}//W_{\mathfrak{s}} \longrightarrow T_{\mathfrak{s}}/W_{\mathfrak{s}}$$

is a morphism of algebraic varieties.

Property 4 of the bijection $\mu_{\mathfrak{s}}$

- For each $\mathfrak{s} \in \pi_o \text{Prim}(\mathcal{HG})$ there is an algebraic family

$$\theta_{\tau} : T_{\mathfrak{s}}//W_{\mathfrak{s}} \longrightarrow T_{\mathfrak{s}}/W_{\mathfrak{s}}$$

of finite morphisms of algebraic varieties, with $\tau \in \mathbb{C}^{\times}$, such that

$$\theta_1 = \rho_{\mathfrak{s}}, \quad \theta_{\sqrt{q}} = \pi_{\mathfrak{s}} \circ \mu_{\mathfrak{s}}, \quad \text{and} \quad \theta_{\sqrt{q}}(T_{\mathfrak{s}}//W_{\mathfrak{s}} - T_{\mathfrak{s}}/W_{\mathfrak{s}}) = \mathfrak{R}_{\mathfrak{s}}.$$

Here q is the order of the residue field of the p -adic field F over which G is defined and $\mathfrak{R}_{\mathfrak{s}} \subset T_{\mathfrak{s}}/W_{\mathfrak{s}}$ is the sub-variety of reducibility.

Property 5 of the bijection $\mu_{\mathfrak{s}}$ (Correcting cocharacters)

- Fix $\mathfrak{s} \in \pi_o \text{Prim}(\mathcal{HG})$. For each irreducible component \mathbf{c} of the affine variety $T_{\mathfrak{s}}//W_{\mathfrak{s}}$ there is a cocharacter

$$h_{\mathbf{c}} : \mathbb{C}^{\times} \longrightarrow T_{\mathfrak{s}}$$

such that

$$\theta_{\tau}(w, t) = \lambda(h_{\mathbf{c}}(\tau) \cdot t)$$

for all $(w, t) \in \mathbf{c}$. In this context a cocharacter is a homomorphism of algebraic groups $\mathbb{C}^{\times} \longrightarrow T_{\mathfrak{s}}$. $\lambda : T_{\mathfrak{s}} \longrightarrow T_{\mathfrak{s}}/W_{\mathfrak{s}}$ is the quotient map from $T_{\mathfrak{s}}$ to $T_{\mathfrak{s}}/W_{\mathfrak{s}}$.

Remark. The equality

$$\theta_{\tau}(w, t) = \lambda(h_{\mathbf{c}}(\tau) \cdot t)$$

is to be interpreted thus:

Let Z_1, Z_2, \dots, Z_r be the irreducible components of the affine variety $T_{\mathfrak{s}}//W_{\mathfrak{s}}$ and let h_1, h_2, \dots, h_r be the cocharacters as in the statement of Property 5. Let

$$\nu_{\mathfrak{s}} : \widetilde{T}_{\mathfrak{s}} \longrightarrow T_{\mathfrak{s}}//W_{\mathfrak{s}}$$

be the quotient map.

Then irreducible components X_1, X_2, \dots, X_r of the affine variety $\widetilde{T}_{\mathfrak{s}}$ can be chosen with

- $\nu_{\mathfrak{s}}(X_j) = Z_j$ for $j = 1, 2, \dots, r$
- For each $\tau \in \mathbb{C}^{\times}$, the map $m_{\tau} : X_j \rightarrow T_{\mathfrak{s}}/W_{\mathfrak{s}}$ which is the composition

$$\begin{aligned} X_j &\longrightarrow T_{\mathfrak{s}} \longrightarrow T_{\mathfrak{s}}/W_{\mathfrak{s}} \\ (w, t) &\longmapsto h_j(\tau)t \longmapsto \lambda(h_j(\tau)t) \end{aligned}$$

makes the diagram

$$\begin{array}{ccc} X_j & \xrightarrow{\nu_{\mathfrak{s}}} & Z_j \\ \downarrow m_{\tau} & & \downarrow \theta_{\tau} \\ T_{\mathfrak{s}}/W_{\mathfrak{s}} & \xrightarrow{I} & T_{\mathfrak{s}}/W_{\mathfrak{s}} \end{array}$$

commutative.

Note that $h_j(\tau)t$ is the product in the algebraic group $T_{\mathfrak{s}}$ of $h_j(\tau)$ and t .

Remark. The conjecture asserts that to calculate

$$\pi_{\mathfrak{s}} : \widehat{G}_{\mathfrak{s}} \longrightarrow T_{\mathfrak{s}}/W_{\mathfrak{s}}$$

two steps suffice:

- Step 1: Calculate $\rho_{\mathfrak{s}} : T_{\mathfrak{s}}//W_{\mathfrak{s}} \longrightarrow T_{\mathfrak{s}}/W_{\mathfrak{s}}$.
- Step 2: Determine the correcting cocharacters.

6. Where are the correcting cocharacters coming from?

In this section, G is a split reductive group defined over F . By a “Langlands parameter” for G we mean a homomorphism of topological groups

$$W_F \times \mathrm{SL}(2, \mathbb{C}) \longrightarrow G^\vee.$$

Here G^\vee is the Langlands dual group of G (a complex Lie group with root datum dual to that of G , for instance $\mathrm{GL}(n, F)^\vee = \mathrm{GL}(n, \mathbb{C})$, $\mathrm{SL}(n, F)^\vee = \mathrm{PGL}(n, \mathbb{C})$, $\mathrm{PGL}(n, F)^\vee = \mathrm{SL}(n, F)$) and W_F is the Weil group attached to F . When restricted to $\mathrm{SL}(2, \mathbb{C})$ the homomorphism of topological groups is required to be algebraic, and when restricted to W_F , is required to map W_F to semisimple elements of G^\vee (see [31; p. 278], [45]).

Remark. For an earlier definition of Langlands parameter based on the Weil-Deligne group see [15; § 8.2] and [12; Ch. 11]. The $W_F \times \mathrm{SL}(2, \mathbb{C})$ definition is better suited to connecting ABP and local Langlands.

The correcting co-characters of Property 5 above seem to be produced by the $\mathrm{SL}(2, \mathbb{C})$ part of the Langlands parameters— i.e. the standard maximal torus of $\mathrm{SL}(2, \mathbb{C})$ identifies with \mathbb{C}^\times :

$$\zeta \longleftrightarrow \begin{bmatrix} \zeta & 0 \\ 0 & \zeta^{-1} \end{bmatrix}, \quad \zeta \in \mathbb{C}^\times$$

Using this identification, when a Langlands parameter is restricted to the maximal torus of $\mathrm{SL}(2, \mathbb{C})$, a cocharacter of G^\vee

$$\mathbb{C}^\times \longrightarrow G^\vee$$

is obtained, and in examples all the correcting cocharacters arise this way.

Example (The Iwahori-spherical component of $\mathrm{GL}(2, F)$). Let $G = \mathrm{GL}(2, F)$.

- $\widehat{G}_{\mathfrak{s}} = \{\text{Smooth irreducible representations of } \mathrm{GL}(2, F) \text{ having a non-zero Iwahori fixed vector}\}.$
- $T_{\mathfrak{s}} = \{\text{unramified characters of the maximal torus of } \mathrm{GL}(2, F) = \mathbb{C}^\times \times \mathbb{C}^\times\}.$
- $W_{\mathfrak{s}} = \text{the Weyl group of } \mathrm{GL}(2, F) = \mathbb{Z}/2\mathbb{Z}.$
- $0 \neq \gamma \in \mathbb{Z}/2\mathbb{Z} \quad \gamma(\zeta_1, \zeta_2) = (\zeta_2, \zeta_1) \quad (\zeta_1, \zeta_2) \in \mathbb{C}^\times \times \mathbb{C}^\times$

The extended quotient $(\mathbb{C}^\times \times \mathbb{C}^\times)/(\mathbb{Z}/2\mathbb{Z})$ is the disjoint union of the ordinary quotient $(\mathbb{C}^\times \times \mathbb{C}^\times)/(\mathbb{Z}/2\mathbb{Z})$ and \mathbb{C}^\times . The ordinary quotient $(\mathbb{C}^\times \times \mathbb{C}^\times)/(\mathbb{Z}/2\mathbb{Z})$ consists of unordered pairs of non-zero complex numbers. Such an unordered pair will be denoted $\{\zeta_1, \zeta_2\}$. The projection of the extended quotient onto the ordinary quotient is the identity map when restricted to the copy of the ordinary quotient contained in the extended quotient— and when restricted to \mathbb{C}^\times maps ζ to $\{\zeta, \zeta\}$. Hence the picture for the projection of the extended quotient onto the ordinary quotient is:

$$\begin{array}{c}
 (\mathbb{C}^\times \times \mathbb{C}^\times)/(\mathbb{Z}/2\mathbb{Z}) \\
 \diagup \\
 \{\zeta_1, \zeta_2\} \text{ such that} \\
 \zeta_1 = \zeta_2
 \end{array}$$

In this picture the ambient variety is the ordinary quotient $(\mathbb{C}^\times \times \mathbb{C}^\times)/(\mathbb{Z}/2\mathbb{Z})$. The locus of points in the ordinary quotient whose pre-image consists of more than one point is $\{\zeta_1, \zeta_2\}$ such that $\zeta_1 = \zeta_2$. This locus is a sub-variety indicated by the slanted line.


For the bijection μ_s composed with the infinitesimal character, the picture is:

$$\begin{array}{c}
 (\mathbb{C}^\times \times \mathbb{C}^\times)//(\mathbb{Z}/2\mathbb{Z}) = (\mathbb{C}^\times \times \mathbb{C}^\times)/(\mathbb{Z}/2\mathbb{Z}) \sqcup \mathbb{C}^\times \\
 (\mathbb{C}^\times \times \mathbb{C}^\times)/(\mathbb{Z}/2\mathbb{Z}) \\
 \diagup \\
 \{\zeta_1, \zeta_2\} \text{ such that} \\
 \{\zeta_1 \zeta_2^{-1}, \zeta_2 \zeta_1^{-1}\} = \{q, q^{-1}\}
 \end{array}$$

In this picture the ambient variety is the ordinary quotient $(\mathbb{C}^\times \times \mathbb{C}^\times)/(\mathbb{Z}/2\mathbb{Z})$. The locus of points in the ordinary quotient whose pre-image consists of more than one point is $\{\zeta_1, \zeta_2\}$ such that $\{\zeta_1 \zeta_2^{-1}, \zeta_2 \zeta_1^{-1}\} = \{q, q^{-1}\}$, where q is the order of the residue field of F . This locus is a sub-variety indicated by the slanted line. On the copy of the ordinary quotient contained within the extended quotient, the map is the identity map. On \mathbb{C}^\times the map is

$$\zeta \mapsto \{q^{1/2}\zeta, q^{-1/2}\zeta\}$$


The correcting cocharacter $\mathbb{C}^\times \rightarrow \mathbb{C}^\times \times \mathbb{C}^\times$ is $\tau \mapsto (\tau, \tau^{-1})$



The picture for $\rho_{\mathfrak{s}}$ is obtained by taking the picture for $\pi_{\mathfrak{s}}$ and setting $q = 1$.

7. Interaction with Baum-Connes and Local Langlands

ABP can be viewed as providing a link between LL (local Langlands conjecture) and BC (Baum-Connes conjecture).



As indicated above, the correcting cocharacters of ABP appear to be coming from the $SL(2, \mathbb{C})$ part of Langlands parameters. This is the interaction of ABP with LL. An intriguing question is "In the ABP view of \widehat{G} , what are the L -packets?".

For a reductive p -adic group G , BC [7] asserts that the Baum-Connes assembly map

$$K_j^G(\beta G) \longrightarrow K_j(C_r^*G) \quad j = 0, 1$$

is an isomorphism of abelian groups. $K_j^G(\beta G)$ is the Kasparov equivariant K -homology of the (extended) affine Bruhat-Tits building βG of G .

$$K_j^G(\beta G) := KK_G^j(C_0(\beta G), \mathbb{C}) \quad j = 0, 1$$

C_r^*G is the reduced C^* algebra of G . This is the C^* algebra obtained by completing $\mathcal{H}G$ via the (left) regular representation of G . $K_*(C_r^*G)$ is the K -theory—in the sense of C^* algebra K -theory—of C_r^*G . The Hecke algebra $\mathcal{H}G$ of G decomposes (canonically) into a direct sum of two-sided ideals:

$$\mathcal{H}(G) = \bigoplus_{\mathfrak{s} \in \pi_o \text{Prim}(\mathcal{H}G)} \mathcal{I}_{\mathfrak{s}}$$

Passing to the C^* algebra completion yields a direct sum decomposition—in the sense of C^* algebras—

$$C_r^*G = \bigoplus_{\mathfrak{s} \in \pi_o \text{Prim}(\mathcal{H}G)} \overline{\mathcal{I}}_{\mathfrak{s}}$$

and this gives a direct sum decomposition of $K_*C_r^*G$

$$K_jC_r^*G = \bigoplus_{\mathfrak{s} \in \pi_o \text{Prim}(\mathcal{H}G)} K_j(\overline{\mathcal{I}}_{\mathfrak{s}}) \quad j = 0, 1$$

ABP at the level of C^* algebra K -theory is:

Conjecture. *Let G be a reductive p -adic group. Then for each $\mathfrak{s} \in \pi_o \text{Prim}(\mathcal{H}G)$*

$$K_j(\overline{\mathcal{I}}_{\mathfrak{s}}) = K_{W_{\mathfrak{s}}}^j(\Delta_{\mathfrak{s}}) \quad j = 0, 1$$

Here $K_{W_{\mathfrak{s}}}^j(\Delta_{\mathfrak{s}})$ is Atiyah-Hirzebruch-Segal topological equivariant K -theory [1] for the finite group $W_{\mathfrak{s}}$ acting on the compact Hausdorff space $\Delta_{\mathfrak{s}}$. Note that the group structure of $\Delta_{\mathfrak{s}}$ is not being used. Applying the Chern character [8] gives a map

$$K_{W_{\mathfrak{s}}}^j(\Delta_{\mathfrak{s}}) \longrightarrow \bigoplus_l H^{j+2l}(\Delta_{\mathfrak{s}}//W_{\mathfrak{s}}; \mathbb{C})$$

which becomes an isomorphism when $K_{W_{\mathfrak{s}}}^j(\Delta_{\mathfrak{s}})$ is tensored with \mathbb{C} . Hence ABP at the level of C^* algebra K -theory gives a much finer and more precise formula for $K_*C_r^*G$ than BC alone provides. For an explicitly computed example see [26].

Theorem 1 (V. Lafforgue [32]). *Baum-Connes is valid for any reductive p -adic group G .*

Theorem 2 ([24, 22, 33]). *Local Langlands is valid for $\text{GL}(n, F)$.*

Theorem 3 ([2, 16, 17]). *ABP is valid for $\text{GL}(n, F)$.*

8. Where is the bijection $\mu_{\mathfrak{s}}: T_{\mathfrak{s}}//W_{\mathfrak{s}} \longleftrightarrow \widehat{G}_{\mathfrak{s}}$ coming from? (ABP Part 2)

Notation. If X is a (complex) affine variety, $\mathcal{O}(X)$ denotes the co-ordinate algebra of X .

As above, the Hecke algebra of G decomposes (canonically) into a direct sum of two-sided ideals:

$$\mathcal{H}(G) = \bigoplus_{\mathfrak{s} \in \pi_o \text{Prim}(\mathcal{H}G)} \mathcal{I}_{\mathfrak{s}}$$

Each ideal $\mathcal{I}_{\mathfrak{s}}$ is canonically Morita equivalent to a unital finite-type $k_{\mathfrak{s}}$ -algebra where

$$k_{\mathfrak{s}} = \mathcal{O}(T_{\mathfrak{s}}/W_{\mathfrak{s}})$$

This unital finite-type $k_{\mathfrak{s}}$ -algebra will be denoted $\mathcal{H}(G)_{\mathfrak{s}}$. The set of (isomorphism classes of) simple $\mathcal{H}(G)_{\mathfrak{s}}$ modules (equivalently, the set of primitive ideals in $\mathcal{H}(G)_{\mathfrak{s}}$) is canonically in bijection with $\widehat{G}_{\mathfrak{s}}$.

$$\text{Prim}(\mathcal{H}(G)_{\mathfrak{s}}) \longleftrightarrow \widehat{G}_{\mathfrak{s}}$$

Fix an affine variety X and consider the category of all finite-type k -algebras, where $k = \mathcal{O}(X)$. These k -algebras are required to be of finite-type (i.e. are required to be finitely generated as k -modules), but are not required to be unital. In [3] the authors of this note introduced an equivalence relation called *geometric equivalence* for such algebras. This equivalence relation is a weakening of Morita equivalence— if two unital finite-type k -algebras are Morita equivalent, then they are geometrically equivalent. A detailed exposition of geometric equivalence will be given in [6]. If

A_1 and A_2 are two finite-type k -algebras which are geometrically equivalent, then there is an isomorphism of periodic cyclic homology [9, 10]

$$\mathrm{HP}_*(A_1) \cong \mathrm{HP}_*(A_2)$$

and a bijection of sets

$$\mathrm{Prim}(A_1) \longleftrightarrow \mathrm{Prim}(A_2).$$

Part 2 of the ABP conjecture is the assertion that:

Conjecture.

- *The finite-type $\mathcal{O}(T_{\mathfrak{s}}/W_{\mathfrak{s}})$ -algebras $\mathcal{H}(G)_{\mathfrak{s}}$ and $\mathcal{O}(T_{\mathfrak{s}}//W_{\mathfrak{s}})$ are geometrically equivalent.*
- $\mathrm{HP}_j(\mathcal{H}(G)_{\mathfrak{s}}) = \bigoplus_l H^{j+2l}(T_{\mathfrak{s}}//W_{\mathfrak{s}}; \mathbb{C})$
- *A bijection $\mu_{\mathfrak{s}} : T_{\mathfrak{s}}//W_{\mathfrak{s}} \longleftrightarrow \widehat{G}_{\mathfrak{s}}$ having properties 1–5 listed above can be constructed by choosing a suitable geometric equivalence between $\mathcal{H}(G)_{\mathfrak{s}}$ and $\mathcal{O}(T_{\mathfrak{s}}//W_{\mathfrak{s}})$.*

9. Recent Developments

Maarten Solleveld [48] has recently proved a result which implies validity for Part 1 of the ABP Conjecture as stated in section 5 whenever $\mathcal{H}(G)_{\mathfrak{s}}$ is an (extended) affine Hecke algebra, perhaps with unequal parameters. For Part 2 of ABP Solleveld does prove the isomorphism of periodic cyclic homology

$$\mathrm{HP}_j(\mathcal{H}(G)_{\mathfrak{s}}) = \bigoplus_l H^{j+2l}(T_{\mathfrak{s}}//W_{\mathfrak{s}}; \mathbb{C}).$$

Although he does not prove that $\mathcal{H}(G)_{\mathfrak{s}}$ is geometrically equivalent to $\mathcal{O}(T_{\mathfrak{s}}//W_{\mathfrak{s}})$, he does obtain a result of this kind for appropriate topological completions of $\mathcal{H}(G)_{\mathfrak{s}}$ and $\mathcal{O}(T_{\mathfrak{s}}//W_{\mathfrak{s}})$.

The unital finite-type $k_{\mathfrak{s}}$ -algebra $\mathcal{H}(G)_{\mathfrak{s}}$ has been proved to be an (extended) affine Hecke algebra in the following cases:

- G split, \mathfrak{s} the Iwahori-spherical component [14];
- G split (some restriction on the residual characteristic of F), \mathfrak{s} in the principal series [46];
- G arbitrary, σ of level 0 [40, 41, 44, 38].
- $G = \mathrm{GL}_n(F)$, \mathfrak{s} arbitrary, and $\mathrm{SL}_n(F)$, many \mathfrak{s} - from the work of Bushnell and Kutzko on types [18, 19, 20] and Goldberg and Roche [21].
- $G = \mathrm{SO}_n(F)$, $G = \mathrm{Sp}_{2n}(F)$ or G an inner form of GL_n , \mathfrak{s} arbitrary, see [23], see also Kim [29, 30].
- $G = \mathrm{GSp}_4(F)$ or $G = \mathrm{U}(2, 1)$, \mathfrak{s} arbitrary [42, 43, 13];

One of these cases is the Iwahori component. This raises the question of reconciling ABP with the Kazhdan-Lusztig parametrization of the Iwahori component. This reconciliation will be given in [5]. More generally, the reconciliation with Reeder's result [45] on the principal series will also be given in [5]. For Reeder, G is assumed to be split and to have connected center (in the Zariski topology).

Detailed calculations which verify ABP completely for the principal series of the p -adic group G_2 ($p \neq 2, 3, 5$) will appear in [4].

10. Appendix

By *extended Coxeter group* we will mean below a semidirect product of a Coxeter group by a finite abelian group. The finite abelian group is assumed to act as automorphisms of the Coxeter system. We give below a construction which in many examples assigns to a Bernstein component an extended (infinite) Coxeter group. In this setting (in examples) a significant simplification is achieved in the correcting cocharacters because the cocharacters are indexed by the cells of the associated extended Coxeter group.

The group $W_{\mathfrak{s}}$ admits the structure of a finite extended Coxeter group.

Let A_M denote the identity component of the center of M and let Φ_M denote the set of roots of A_M . For each $\alpha \in \Phi_M$, we write U_{α} for the corresponding root group and set $G_{\alpha} := \langle M, U_{\alpha}, U_{-\alpha} \rangle$. Then $G_{\alpha} = G_{-\alpha}$ is a reductive group. It has two parabolic subgroups with Levi component M , $P_{\alpha} = MU_{\alpha}$ and $P_{-\alpha} = MU_{-\alpha}$. Let ψ be an unramified character of M . By normalized parabolic induction from P_{α} and $P_{-\alpha}$, we obtain the representations $\text{Ind}_{P_{\alpha}}^{G_{\alpha}}(\psi \otimes \sigma)$ and $\text{Ind}_{P_{-\alpha}}^{G_{\alpha}}(\psi \otimes \sigma)$. These are related by standard intertwining operators

$$J_{-\alpha, \alpha}: \text{Ind}_{P_{\alpha}}^{G_{\alpha}}(\psi \otimes \sigma) \rightarrow \text{Ind}_{P_{-\alpha}}^{G_{\alpha}}(\psi \otimes \sigma) \quad \text{and} \quad J_{\alpha, -\alpha}: \text{Ind}_{P_{-\alpha}}^{G_{\alpha}}(\psi \otimes \sigma) \rightarrow \text{Ind}_{P_{\alpha}}^{G_{\alpha}}(\psi \otimes \sigma).$$

There is a rational function $j_{\alpha}(\psi, \sigma)$ on the complex torus $\widetilde{T}_{\mathfrak{s}}$ such that

$$J_{\alpha, -\alpha} \circ J_{-\alpha, \alpha} = j_{\alpha}(\psi, \sigma) \text{id}$$

(and $j_{\alpha}(\psi, \sigma) = j_{-\alpha}(\psi, \sigma)$).

Let

$$\Phi_{\mathfrak{s}} := \{\alpha \in \Phi_M : j(\psi, \sigma) \text{ has a pole}\}.$$

The set $\Phi_{\mathfrak{s}}$ is a root system. It is preserved by $W_{\mathfrak{s}}$. Let $W_{\mathfrak{s}}^0$ denote the finite Weyl group associated to $\Phi_{\mathfrak{s}}$. Fix a positive system $\Phi_{\mathfrak{s}}^+$ in $\Phi_{\mathfrak{s}}$ and set

$$C_{\mathfrak{s}} := \{w \in W_{\mathfrak{s}} : w(\Phi_{\mathfrak{s}}^+) \subset \Phi_{\mathfrak{s}}^+\}.$$

Then the group $W_{\mathfrak{s}}$ is the semi-direct product

$$W_{\mathfrak{s}} = W_{\mathfrak{s}}^0 \rtimes C_{\mathfrak{s}}.$$

Hence $W_{\mathfrak{s}}$ occurs to be a finite extended Coxeter group.

An extended (infinite) Coxeter group attached to \mathfrak{s} .

Set

$$M_{\mathfrak{s}} := \bigcap_{\varphi \in I_{\sigma}} \text{Ker}(\varphi) \quad \text{and} \quad \Lambda_{\mathfrak{s}} := M_{\mathfrak{s}}/M^0.$$

The group $\Lambda_{\mathfrak{s}}$ is free abelian of the same rank as M/M^0 . Conjugation by $W_{\mathfrak{s}}$ preserves $M_{\mathfrak{s}}$. There is therefore an induced action of $W_{\mathfrak{s}}$ on $\Lambda_{\mathfrak{s}}$. Now $\Lambda_{\mathfrak{s}}$ is isomorphic to the group of characters $X(T_{\mathfrak{s}})$ of the complex torus $T_{\mathfrak{s}}$.

Then we set

$$\widetilde{W}_{\mathfrak{s}} := X(T_{\mathfrak{s}}) \rtimes W_{\mathfrak{s}}.$$

We have

$$\widetilde{W}_{\mathfrak{s}} \simeq (X(T_{\mathfrak{s}}) \rtimes W_{\mathfrak{s}}^0) \rtimes C_{\mathfrak{s}}.$$

In many examples the group $X(T_{\mathfrak{s}}) \rtimes W_{\mathfrak{s}}^0$ is an affine Weyl group. Hence $\widetilde{W}_{\mathfrak{s}}$ in many examples is an extended Coxeter group.

A weight function attached to \mathfrak{s}

Let M^\vee denote the identity component of the Langlands dual group of M , let $\alpha \in \Phi_M$, and let r_α denote the adjoint representation of M^\vee on the Lie algebra of U_α^\vee . We are assuming here that σ is generic so that the corresponding local L -functions $L(s, \sigma, r_\alpha)$ are defined by Shahidi. Then the definition of $\Phi_\mathfrak{s}$ can be rephrased as follows (thanks to a formula by Shahidi for $j_\alpha(\psi, \sigma)$ [47]):

$$\Phi_\mathfrak{s} = \{\alpha \in \Phi_M : L(s, \sigma, r_\alpha) \neq 1\}.$$

For each $\alpha \in \Phi_M$, we denote by q_α the degree of the L -function $L(s, \sigma, r_\alpha)$, that is, q_α is the degree of $P(T)$ where $P(T)$ is the polynomial such that $L(s, \sigma, r_\alpha) = P(q^{-s})^{-1}$. We have $q_{w\alpha} = q_\alpha$ for all $w \in W_\mathfrak{s}$. From this we define a *weight function* $\mathbf{q}_\mathfrak{s}: \Phi_M \rightarrow \mathbf{N}$ by

$$\mathbf{q}_\mathfrak{s}(\alpha) := q_\alpha \quad \alpha \in \Phi_M.$$

The collection of two-sided cells attached to \mathfrak{s}

In the case when the function $\mathbf{q}_\mathfrak{s}$ is constant, we will attach to \mathfrak{s} the collection of all the *two-sided cells* of the group $\widetilde{W}_\mathfrak{s}$ as originally defined by Lusztig in [34], [35].

In the general case, Lusztig stated in [39] (and proved in several cases) a list of conjectures, and, assuming their validity, defined partitions of any extended Coxeter group equipped with a weight function as above into left cells, right cells and two-sided cells, which extend the theory previously developed by Kazhdan and him in the case of “equal parameters” (that is, when $\mathbf{q}_\mathfrak{s}$ is constant). Hence we can attach (at least conjecturally) a collection of two-sided cells to $(\widetilde{W}_\mathfrak{s}, \mathbf{q}_\mathfrak{s})$. We will denote this collection by $\text{Cell}(\widetilde{W}_\mathfrak{s}, \mathbf{q}_\mathfrak{s})$.

It is part of ABP Conjecture that each cocharacter should be attached to a two-sided cell of $(\widetilde{W}_\mathfrak{s}, \mathbf{q}_\mathfrak{s})$. This part of ABP Conjecture is proved for principal series of G_2 (when $p \neq 2, 3, 5$) in [4].

More precisely, for each Bernstein component \mathfrak{s} attached to a principal series of G_2 , the weight function $\mathbf{q}_\mathfrak{s}$ is the constant function, and there is a decomposition of the extended quotient

$$T_\mathfrak{s} // W_\mathfrak{s} = \bigsqcup_{\mathbf{c} \in \text{Cell}(\widetilde{W}_\mathfrak{s}, \mathbf{q}_\mathfrak{s})} (T_\mathfrak{s} // W_\mathfrak{s})_{\mathbf{c}},$$

such that each $(T_\mathfrak{s} // W_\mathfrak{s})_{\mathbf{c}}$ is a union of irreducible components c of the affine variety $T_\mathfrak{s} // W_\mathfrak{s}$ and

$$h_c = h_{c'} \quad \text{for all } c, c' \subset (T_\mathfrak{s} // W_\mathfrak{s})_{\mathbf{c}}.$$

Moreover, the partition can be chosen so that the following property holds:

$$T_\mathfrak{s} // W_\mathfrak{s} \subset (T_\mathfrak{s} // W_\mathfrak{s})_{\mathbf{c}_0},$$

where \mathbf{c}_0 denotes the *lowest* two-sided cell in $\widetilde{W}_\mathfrak{s}$ in the natural partial ordering on $\text{Cell}(\widetilde{W}_\mathfrak{s}, \mathbf{q}_\mathfrak{s})$. Note the inclusion above is not an equality in general.

Where is the geometric equivalence $\mathcal{H}(G)_\mathfrak{s} \simeq \mathcal{O}(T_\mathfrak{s} // W_\mathfrak{s})$ coming from?

Let $\mathcal{J}^\mathfrak{s}$ denote the *Lusztig asymptotic algebra* attached to the group $(\widetilde{W}_\mathfrak{s}, \mathbf{q}_\mathfrak{s})$ (in case the weight function $\mathbf{q}_\mathfrak{s}$ is constant, $\mathcal{J}^\mathfrak{s}$ is defined in [36; §1.3], in general, the definition of $\mathcal{J}^\mathfrak{s}$ is given in [39] up to a list of conjectural properties). The

algebra $\mathcal{J}^{\mathfrak{s}}$ admits a canonical decomposition into finitely many two-sided ideals

$$\mathcal{J}^{\mathfrak{s}} = \bigoplus_{\mathfrak{c} \in \text{Cell}(\widetilde{W}_{\mathfrak{s}}, \mathfrak{q}_{\mathfrak{s}})} \mathcal{J}_{\mathfrak{c}}^{\mathfrak{s}}.$$

One can provide $\mathcal{J}^{\mathfrak{s}}$ (and also each $\mathcal{J}_{\mathfrak{c}}^{\mathfrak{s}}$) with a structure of $k_{\mathfrak{s}}$ -module algebra (see [3; § 9]). Then $\mathcal{J}_{\mathfrak{s}}$ is a finite-type $k_{\mathfrak{s}}$ algebra.

It is also part of ABP Conjecture that the conjectural geometric equivalence $\mathcal{H}(G)_{\mathfrak{s}} \simeq \mathcal{O}(T_{\mathfrak{s}}//W_{\mathfrak{s}})$ comes from the combination of the following two geometric equivalences:

$$\mathcal{H}(G)_{\mathfrak{s}} \simeq \mathcal{J}^{\mathfrak{s}} \quad \text{and} \quad \mathcal{J}^{\mathfrak{s}} \simeq \mathcal{O}(T_{\mathfrak{s}}//W_{\mathfrak{s}}).$$

The first geometric equivalence is proved for G arbitrary, \mathfrak{s} the Iwahori component in [9, 10], and G split (with mild restriction on the residual characteristic) in the principal series case, by combining [9, 10] with [46].

The second geometric equivalence is proved for $G = \text{GL}(n, F)$, \mathfrak{s} arbitrary in [16] [17], for $G = \text{SL}(2, F)$, \mathfrak{s} arbitrary in [3; § 7], for $G = \text{G}_2$, principal series case in [4], for $G = \text{PGL}(n, F)$, \mathfrak{s} the Iwahori component in [3; § 12], for $G = \text{SO}(5, F)$, \mathfrak{s} the Iwahori component in [3; § 13], for $G = \text{SO}(4, F)$, \mathfrak{s} the Iwahori component in [4; § 8].

Geometric equivalence respects direct sums. For $G = \text{G}_2$, principal series case, and for $G = \text{SO}(4, F)$, \mathfrak{s} the Iwahori component, the second geometric equivalence comes from a finite collection of geometric equivalences:

$$\mathcal{J}_{\mathfrak{c}}^{\mathfrak{s}} \simeq \mathcal{O}(T_{\mathfrak{s}}//W_{\mathfrak{s}})_{\mathfrak{c}}, \quad \text{for any } \mathfrak{c} \in \text{Cell}(\widetilde{W}_{\mathfrak{s}}, \mathfrak{q}_{\mathfrak{s}}).$$

Beyond the p -adic world.

Thanks to the work of Solleveld [48], ABP Conjecture (at least Part 1) is still existing (and partly proved) even if there is no p -adic group in the picture. In the situation considered by Solleveld, no field F given, he is working with an (extended) Hecke algebra with parameters, which essentially means that the weight function $\mathfrak{q}_{\mathfrak{s}}$ is here replaced by a more general parameter function

$$\mathfrak{q}: \Phi_{\text{nr}}^{\vee} \rightarrow \mathbb{R}_{>0},$$

where $\mathcal{R} = (X, \Phi, X^{\vee}, \Phi^{\vee})$ is a root datum, Φ a reduced root system, and Φ_{nr}^{\vee} the non-reduced root system:

$$\Phi_{\text{nr}}^{\vee} := \Phi \cup \{2\alpha : \alpha^{\vee} \in X^{\vee}\}.$$

The function \mathfrak{q} is assumed to be W_0 -invariant, with W_0 the Weyl group of \mathcal{R} . Solleveld extends in [48] the previous notion of cocharacters to this setting, and states (and mainly proved) a version of ABP conjecture in the context of extended Hecke algebras with parameters. His work includes in particular the case of exotic Kazhdan-Lusztig parameters as defined in [27].

References

- [1] M.F. Atiyah, *K-theory*. Lecture notes by D. W. Anderson W. A. Benjamin, Inc., New York-Amsterdam 1967 v+166+xlix pp.
- [2] A.-M. Aubert, P. Baum, R.J. Plymen, Geometric structure in the representation theory of p -adic group, C.R. Acad. Sci. Paris, Ser. I, 345 (2007) 573–578.

- [3] A.-M. Aubert, P. Baum, R.J. Plymen, The Hecke algebra of a reductive p -adic group: a geometric conjecture, *Aspects of Math.* 37, Vieweg Verlag (2006) 1–34.
- [4] A.-M. Aubert, P. Baum, R.J. Plymen, Geometric structure in the principal series of the p -adic group G_2 , *Represent. Theory*, to appear.
- [5] A.-M. Aubert, P. Baum, R.J. Plymen, Geometric structure for L -parameters of Springer type, preprint 2010.
- [6] A.-M. Aubert, P. Baum, R.J. Plymen, Morita equivalence revisited, (to appear).
- [7] P. Baum, A. Connes, N. Higson, Classifying space for proper actions and K -theory of group C^* -algebras, In: *C^* -algebras:1943-1993*, Amer. Math. Soc., Providence, RI, 1994, pp.240-291.
- [8] P. Baum, A. Connes, Chern character for discrete groups, In: *A Fete of Topology*, Academic Press, New York, 1988, pp. 163–232.
- [9] P. Baum, V. Nistor, Periodic cyclic homology of Iwahori-Hecke algebras. *K-Theory* 27 (2002), no. 4, 329–357.
- [10] P. Baum, V. Nistor, Periodic cyclic homology of Iwahori-Hecke algebras. *C. R. Acad. Sci. Paris Sr. I Math.* 332 (2001), no. 9, 783–788.
- [11] J.N. Bernstein, Le “centre” de Bernstein. (French) [The Bernstein “center”] Edited by P. Deligne. *Travaux en Cours, Representations of reductive groups over a local field*, 1–32, Hermann, Paris, 1984.
- [12] J.N. Bernstein, S. Gelbart, *An introduction to the Langlands program*, Birkhäuser, 2003.
- [13] L. Blasco, C. Blondel, Algèbres de Hecke et séries principales généralisées de $\mathrm{Sp}_4(F)$, *Proc. London Math. Soc.* (3) 85 (2002), 659–685.
- [14] A. Borel, Admissible representations of a semisimple group over a local field with vectors fixed under an Iwahori subgroup, *Inv. Math.* 35 (1976), 233–259.
- [15] A. Borel, Automorphic L -functions, *PSPM* 33 (1979) 27–61.
- [16] J. Brodzki, R.J. Plymen, Geometry of the smooth dual of $\mathrm{GL}(n)$, *C.R. Acad. Sci. Paris, Ser. I*, 331 (2000) 213–218.
- [17] J. Brodzki, R.J. Plymen, Complex structure on the smooth dual of $\mathrm{GL}(n)$, *Documenta Math.* 7 (2002) 91–112.
- [18] C.J. Bushnell, P.C. Kutzko, The admissible dual of $\mathrm{GL}(n)$ via compact open subgroups, *Ann. Math. Study* 129, Princeton Univ. Press 1993.
- [19] C.J. Bushnell, P.C. Kutzko, Semisimple types in GL_n , *Comp. Math.* 119 (1999), 53–97.
- [20] C.J. Bushnell, P.C. Kutzko, Smooth representations of reductive p -adic groups: Structure theory via types, *Proc. London Math. Soc.* 77 (1998), 582–634.
- [21] D. Goldberg, A. Roche, Hecke algebras and SL_n -types, *Proc. London Math. Soc.* (3) 90 (2005), no. 1, 87–131.
- [22] M. Harris, R. Taylor, Geometry and cohomology of simple Shimura varieties, *Annals of Math. Studies* 151 (2001).
- [23] V. Heiermann, Opérateurs d’entrelacement et algèbres de Hecke avec paramètres d’un groupe réductif p -adique: le cas des groupes classiques, preprint 2009, arXiv:0804.4398v2.
- [24] G. Henniart, Une preuve simple des conjectures de Langlands pour $\mathrm{GL}(n)$ sur un corps p -adiques, *Invent. Math.* 139 (2000) 439–455.

- [25] Kim, J-L: Supercuspidal representations: An exhaustion theorem, *J. Amer. Math. Soc.* 20 (2007), 273-320.
- [26] J. Jawdat, R.J. Plymen, R -groups and geometric structure in the representation theory of $SL(N)$, *J. noncommut. Geom.* 4 (2010) 265–279.
- [27] Syu Kato, An exotic Deligne-Langlands correspondence for symplectic groups, *Duke Math. J.* 148.2 (2009) 305–371.
- [28] D. Kazhdan, G. Lusztig, Proof of the Deligne-Langlands conjecture for Hecke algebras, *Invent. Math.* 87 (1987) 153–215.
- [29] J.-L. Kim, Hecke algebras of classical groups over p -adic fields and supercuspidal representations, *Amer. J. Math.* 121 (1999), 967–1029.
- [30] J.-L. Kim, Hecke algebras of classical groups over p -adic fields II, *Compositio Math.* 127 (2001), 117–167.
- [31] A.W. Knap, Introduction to the Langlands program, *Proc. Symp. Pure Math.* 61 (1997) 245–302.
- [32] V. Lafforgue, KK -théorie bivariante pour les algèbres de Banach et conjecture de Baum-Connes, *Invent. Math.* 149 (2002) 1–95.
- [33] G. Laumon, M. Rapoport, U. Stuhler, \mathcal{D} -elliptic sheaves and the Langlands correspondence, *Invent. Math.* 113 (1993) 217–338.
- [34] G. Lusztig, Cells in affine Weyl groups, *Advanced Studies in Pure Math.* 6 (1985), in *Algebraic Groups and Related Topics*, 255–287.
- [35] G. Lusztig, Cells in affine Weyl groups, II, *J. Algebra* 109 (1987), 536–548.
- [36] G. Lusztig, Cells in affine Weyl groups, III, *J. Fac. Sci. Univ. Tokyo, Sect. IA, Math.* 34 (1987), 223–243.
- [37] G. Lusztig, Cells in affine Weyl groups, IV, *J. Fac. Sci. Univ. Tokyo, Sect. IA, Math.* 36 (1989), 297–328.
- [38] G. Lusztig, Classification of unipotent representations of simple p -adic groups, *Internat. Math. Res. Notices* 11 (1995), 517–589.
- [39] G. Lusztig, Hecke algebras with unequal parameters, *CRM Monograph Series*, 18, Amer. Math. Soc. 2003.
- [40] L. Morris, Tamely ramified intertwining algebras, *Invent. Math.* 114 (1993), 1–54.
- [41] L. Morris, Level zero G -types, *Compositio Math.* 118 (1999), 135–157.
- [42] A. Moy, Representations of $\mathrm{GSp}(4)$ over a p -adic field, I and II, *Compositio Math.* 66.3 (1988) 237–284 and 285–328.
- [43] A. Moy, Representations of $\mathrm{U}(2, 1)$ over a p -adic field, *J. Reine angew. Math.* 372 (1986) 178–208.
- [44] A. Moy, G. Prasad, Jacquet functors and unrefined minimal K -types, *Comment. Helvetici* 71 (1996) 98–121.
- [45] M. Reeder, Isogenies of Hecke algebras and a Langlands correspondence for ramified principal series representations, *Representation Theory* 6 (2002), 101–126.
- [46] A. Roche, Types and Hecke algebras for principal series representations of split reductive p -adic groups, *Ann. scient. Éc. Norm. Sup.* 31 (1998), 361–413.
- [47] F. Shahidi, Langlands’ conjecture on Plancherel measures for p -adic groups, in *Harmonic analysis on reductive groups* (Brunswick, ME, 1989), *Progr. Math.*, 101, Birkhäuser Boston, Boston, MA, 277–295, 1991.
- [48] M. Solleveld, On the classification of irreducible representations of affine Hecke algebras with unequal parameters, preprint 2010.

- [49] Stevens, S, The supercuspidal representations of p -adic classical groups, *Invent. Math.* 172(2) (2008) 289-352.

INSTITUT DE MATHÉMATIQUES DE JUSSIEU, U.M.R. 7586 DU C.N.R.S., PARIS, FRANCE
E-mail address: `aubert@math.jussieu.fr`

MATHEMATICS DEPARTMENT, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16802,
USA

E-mail address: `baum@math.psu.edu`

SCHOOL OF MATHEMATICS, ALAN TURING BUILDING, MANCHESTER UNIVERSITY, MANCHESTER
M13 9PL, ENGLAND

E-mail address: `plymen@manchester.ac.uk`