

***Affine Weyl groups and Langlands duality***

Niblo, Graham and Plymen, Roger and Wright, Nick

2014

MIMS EPrint: **2014.45**

Manchester Institute for Mathematical Sciences  
School of Mathematics

The University of Manchester

Reports available from: <http://eprints.maths.manchester.ac.uk/>

And by contacting: The MIMS Secretary  
School of Mathematics  
The University of Manchester  
Manchester, M13 9PL, UK

ISSN 1749-9097

# AFFINE WEYL GROUPS AND LANGLANDS DUALITY

GRAHAM NIBLO, ROGER PLYMEN AND NICK WRIGHT

ABSTRACT. Let  $G$  be a compact connected semisimple Lie group. We show that, as well as the duality between  $K$ -theory and  $K$ -homology, there is also a Langlands duality in the Baum-Connes correspondence for the (extended) affine Weyl group attached to  $G$ .

## CONTENTS

1. Introduction	1
2. Langlands duality	3
2.1. Complex reductive groups	3
2.2. Compact semisimple groups	4
2.3. Example: $SU_3(\mathbb{C})$	5
2.4. The nodal group	5
2.5. A table of Langlands dual groups	6
3. Affine Weyl groups	6
4. $C^*$ -algebras	7
5. “ $KK$ -Langlands”	9
5.1. Construction of the Dirac class	9
5.2. The Kasparov product	12
5.3. The $KK$ -equivalence	16
6. Notes and computations	16
6.1. The extended quotient	17
6.2. The quotient $T/W$	18
6.3. The extended affine Weyl group attached to $PSU_3$	18
6.4. The affine Weyl group attached to $PSU_3$	19
6.5. The affine Weyl group attached to $SU_3$	19
6.6. The affine Weyl group attached to $G_2$	19
References	20

## 1. INTRODUCTION

In this paper we examine the Baum-Connes correspondence in the context of (extended) affine Weyl groups associated with a compact connected semisimple Lie group  $G$ . The extended affine Weyl group  $W'_a$  of a Lie group  $G$  can be realised as a group of affine isometries of the Lie algebra of a maximal torus of  $G$ . We denote the maximal torus by  $T$  and its Lie algebra

by  $\mathfrak{t}$ . The action of  $W'_a$  on the Lie algebra  $\mathfrak{t}$  provides a universal example for proper actions of  $W'_a$  and hence the domain of the assembly map is the equivariant  $K$ -homology group  $K_*^{W'_a}(\mathfrak{t})$ . The group  $W'_a$  is the semidirect product of a lattice  $\Gamma$  of translations of  $\mathfrak{t}$  by the Weyl group  $W$  of  $G$ , which acts linearly on  $\mathfrak{t}$ . The quotient of  $\mathfrak{t}$  by the translation action of  $\Gamma$  recovers the torus  $T$  and hence the left hand side of the assembly map can be identified as

$$K_*^{W'_a}(\mathfrak{t}) \cong K_W^*(C_0(\mathfrak{t}) \rtimes \Gamma) \cong K_*^W(T)$$

see below.

On the other hand, the right hand side of the assembly map is the  $K$ -theory of the algebra

$$C_r^*W'_a \cong C_r^*\Gamma \rtimes W \cong C(\widehat{\Gamma}) \rtimes W.$$

Here  $\widehat{\Gamma}$  denotes the Pontryagin dual of the lattice  $\Gamma$ , which is a torus of the same dimension as  $T$ . One might therefore be tempted to think that the Baum-Connes correspondence in this case is an isomorphism between the  $W$ -equivariant  $K$ -homology and  $K$ -theory of the torus  $T$ . While such an isomorphism very often exists, this is not the Baum-Connes correspondence. Indeed although  $\widehat{\Gamma}$  is a torus of the same dimension as  $T$ , there is in general no  $W$ -equivariant identification of the two tori.

An example is given by the Lie group  $SU(3)$  whose extended affine Weyl group (which in this case is its affine Weyl group) is the  $(3, 3, 3)$ -triangle group acting on the plane. The maximal torus  $T$  can be realised as a hexagon  $X$  with opposing sides identified and the Weyl group  $W$  (which is the dihedral group  $D_3$ ) acts by reflecting in the three diagonals of  $X$ . By contrast we show that the dual torus  $\widehat{\Gamma}$  can be realised as a different hexagon  $X^\vee$  with opposing sides identified. The new hexagon should be viewed as the dual hexagon of  $X$  and the group  $W$  now acts by reflections in the bisectors of the edges of  $X^\vee$ . The corresponding action on the plane is by an index 3 extension of the triangle group, obtained by adjoining an order 3 rotation. Hence the dual picture has  $C_3$  isotropy (as well as  $C_2$  and  $D_3$  isotropy), while the undualised picture does not. The tori cannot therefore be  $W$ -equivariantly identified in this example.

Given that the left- and right-hand sides of the Baum-Connes correspondence look so different in this example the isomorphism might almost appear coincidental. This ‘coincidence’ however can be explained by a duality between the tori  $T$  and  $\widehat{\Gamma}$  which, as we will show, yields a duality in  $K$ -theory. This is in addition to the Poincaré duality from  $K$ -theory to  $K$ -homology and Fourier-Pontryagin duality from  $C_r^*(\Gamma)$  to  $C(\widehat{\Gamma})$ . The torus  $\widehat{\Gamma}$  is the  $T$ -dual of the torus  $T$ , which means at the level of Lie groups that  $\widehat{\Gamma}$  is the maximal torus  $T^\vee$  of the Langlands dual  $G^\vee$  of  $G$ . We show that the identification of  $\widehat{\Gamma}$  with  $T^\vee$  is  $W$ -equivariant and thus the action of  $W$  on  $\widehat{\Gamma}$  corresponds to the action of the extended affine Weyl group of  $G^\vee$  on the Lie algebra  $\mathfrak{t}^\vee$  (which is canonically identified with the dual space  $\mathfrak{t}^*$ ). We

show that the duality between  $T$  and  $T^\vee$  yields a natural isomorphism from the  $W$ -equivariant  $K$ -homology of  $T$  to the  $W$ -equivariant  $K$ -theory of  $T^\vee$ . We thus obtain the following commutative diagram,

$$\begin{array}{ccc} KK_{W'_a}^*(C_0(\mathfrak{t}), \mathbb{C}) & \xrightarrow{\mu} & KK(\mathbb{C}, C_r^* W'_a) \\ \downarrow \cong & & \uparrow \cong \\ KK_W^*(C(T), \mathbb{C}) & \xrightarrow{\cong} & KK_W^*(\mathbb{C}, C(T^\vee)) \end{array}$$

where  $\mu$  is the Baum-Connes assembly map.

We obtain the bottom isomorphism as the composition of the Poincaré duality isomorphism from  $KK_W^*(C(T), \mathbb{C})$  to  $KK_W^*(\mathbb{C}, C(T) \hat{\otimes} \mathcal{C}\ell(\mathfrak{t}))$  with a ‘Langlands’ isomorphism from  $KK_W^*(\mathbb{C}, C(T) \hat{\otimes} \mathcal{C}\ell(\mathfrak{t}))$  to  $KK_W^*(\mathbb{C}, C(T^\vee))$ , thereby giving an independent proof of the Baum-Connes correspondence in this context. Here  $\mathcal{C}\ell(\mathfrak{t})$  denotes the complex Clifford algebra of  $\mathfrak{t}$ .

The duality between  $G$  and  $G^\vee$  is further amplified by the following theorem.

**Theorem 1.1.** *Let  $G$  be a compact connected semisimple Lie group and  $G^\vee$  its Langlands dual. Let  $W'_a(G)$ ,  $W'_a(G^\vee)$  denote the extended affine Weyl groups of  $G$  and  $G^\vee$  respectively. Then there is a natural isomorphism*

$$K_*(C_r^*(W'_a(G))) \cong K_*(C_r^*(W'_a(G^\vee))).$$

Hence in particular, if  $G$  is of adjoint type then for  $W_a(G)$  the affine Weyl group of  $G$  we have

$$K_*(C_r^*(W'_a(G))) \cong K_*(C_r^*(W_a(G))).$$

In the above example the Langlands dual of  $SU(3)$  is  $PSU(3)$ , and the index 3 extension of the triangle group is the extended affine Weyl group of  $PSU(3)$ .

We note that the last isomorphism does not hold in general in the case of a group of non adjoint type. Such an example is provided by the group  $SU(4)/\{\pm I\}$ . This group is neither simply connected nor of adjoint type. It is, however, self-dual.

## 2. LANGLANDS DUALITY

**2.1. Complex reductive groups.** Let  $\mathbf{H}$  be a connected complex reductive group, with maximal torus  $\mathbf{S}$ . This determines a root datum

$$R(\mathbf{H}, \mathbf{S}) := (\mathbf{X}^*(\mathbf{S}), R, \mathbf{X}_*(\mathbf{S}), R^\vee)$$

Here  $R$  and  $R^\vee$  are the sets of roots and coroots of  $\mathbf{H}$ , while

$$(1) \quad \mathbf{X}^*(\mathbf{S}) := \text{Hom}(\mathbf{S}, \mathbb{C}^\times) \quad \text{and} \quad \mathbf{X}_*(\mathbf{S}) := \text{Hom}(\mathbb{C}^\times, \mathbf{S})$$

are its character and co-character lattices.

The root datum implicitly includes the pairing  $\mathbf{X}^*(\mathbf{S}) \times \mathbf{X}_*(\mathbf{S}) \rightarrow \mathbb{Z}$  and the bijection  $R \rightarrow R^\vee$ ,  $\alpha \mapsto h_\alpha$  between roots and coroots. Root data

classify complex reductive Lie groups, in the sense that two such groups are isomorphic if and only if their root data are isomorphic.

A root datum determines the reductive group  $\mathbf{H}$  up to isomorphism. Interchanging the roles of roots and coroots and of the character and co-character lattices results in a new root datum:

$$R(\mathbf{H}, \mathbf{S})^\vee := (\mathbf{X}_*(\mathbf{S}), R^\vee, \mathbf{X}^*(\mathbf{S}), R)$$

The Langlands dual group of  $\mathbf{H}$  is the complex reductive group  $\mathbf{H}^\vee$  (unique up to isomorphism) determined by the dual root datum  $R(\mathbf{H}, \mathbf{S})^\vee$ . A root datum also implies a choice of maximal torus  $\mathbf{S} \subset \mathbf{H}$  via the canonical isomorphism  $\mathbf{S} \simeq \text{Hom}(\mathbf{X}^*(\mathbf{S}), \mathbb{C}^\times)$ , and likewise a natural choice of maximal torus for the Langlands dual group  $\mathbf{H}^\vee : \mathbf{S}^\vee := \text{Hom}(\mathbf{X}_*(\mathbf{S}), \mathbb{C}^\times) \subset \mathbf{H}^\vee$ .

In particular, we have the equation

$$(2) \quad \mathbf{X}^*(\mathbf{S}^\vee) = \mathbf{X}_*(\mathbf{S})$$

**2.2. Compact semisimple groups.** Let now  $G$  be a compact connected semisimple Lie group, with maximal torus  $T$ . We recall that a compact connected Lie group is semisimple if and only if it has finite centre [B, p.285]. The classical examples are the compact real forms

$$\text{SU}_n, \text{SO}_{2n+1}, \text{Sp}_{2n}, \text{SO}_{2n}, E_6, E_7, E_8, F_4, G_2$$

The passage from  $G$  to its Langlands dual  $G^\vee$  is via the *complexification*  $G_\mathbb{C}$  of  $G$ . The correspondence

$$G \mapsto G_\mathbb{C}$$

is *bijective*, see [D, 27.17.11], and we have  $\mathfrak{g}_\mathbb{C} = \mathfrak{g} \oplus i\mathfrak{g}$ . Then pass to the Langlands dual  $(G_\mathbb{C})^\vee$  of the complex group  $G_\mathbb{C}$ . Finally, define  $G^\vee$  to be a maximal compact subgroup of  $(G_\mathbb{C})^\vee$ .

The dual group  $G^\vee$  is determined up to isomorphism by the condition

$$(G^\vee)_\mathbb{C} = (G_\mathbb{C})^\vee.$$

This determines  $G^\vee$  up to  $(G_\mathbb{C})^\vee$ -conjugacy.

Let  $T$  be a maximal torus in  $G$ . Then  $\mathbf{S} := T_\mathbb{C}$  is a maximal torus in  $\mathbf{H} := G_\mathbb{C}$ , and so the dual torus  $\mathbf{S}^\vee$  is well-defined in the dual group  $\mathbf{H}^\vee$ . Then  $T^\vee$  is determined by the condition

$$(T^\vee)_\mathbb{C} = \mathbf{S}^\vee.$$

By definition, the torus  $T^\vee$  is the  $T$ -dual of  $T$ . Corresponding to (2), we have the  $T$ -duality equation

$$(3) \quad X^*(T^\vee) = X_*(T)$$

where we agree that  $X^*(T)$  shall mean the group of morphisms from the Lie group  $T$  to the Lie group  $\mathbf{U} = \{z \in \mathbb{C} : |z| = 1\}$ , and  $X_*(T)$  shall mean the group of morphisms from the Lie group  $\mathbf{U}$  to the Lie group  $T$ .

**2.3. Example:**  $\mathrm{SU}_3(\mathbb{C})$ . Let  $G = \mathrm{SU}_3(\mathbb{C})$ . Then  $G^\vee = \mathrm{PSU}_3(\mathbb{C})$  and we have

$$\begin{aligned} T &= \{(z_1, z_2, z_3) : z_j \in \mathbf{U}, z_1 z_2 z_3 = 1\} \\ T^\vee &= \{(z_1 : z_2 : z_3) : z_j \in \mathbf{U}, z_1 z_2 z_3 = 1\} \end{aligned}$$

the latter being in homogeneous coordinates. The map

$$T \rightarrow T^\vee, \quad (z_1, z_2, z_3) \mapsto (z_1 : z_2 : z_3)$$

is a 3-fold cover: the pre-image of  $(z_1 : z_2 : z_3)$  is the set

$$\{\eta z_1, \eta z_2, \eta z_3) : \eta \in \mathbf{U}, \eta^3 = 1\}.$$

The Lie group  $G$  and its dual  $G^\vee$  admit a common Weyl group

$$W = N(T)/T = N(T^\vee)/T^\vee$$

in this case the symmetric group  $\mathfrak{S}_3$ . Note that, in general,  $T$  and  $T^\vee$  are *not* isomorphic as  $W$ -spaces. With  $G = \mathrm{SU}_3(\mathbb{C})$ ,  $T$  admits three  $W$ -fixed points, namely

$$\{(1, 1, 1), (\omega, \omega, \omega), (\omega^2, \omega^2, \omega^2) : \omega = \exp(2\pi i/3)\}$$

whereas the unique  $W$ -fixed point in  $T^\vee$  is the identity  $I \in T^\vee$ .

**2.4. The nodal group.** Let  $G$  be as above,  $T$  a maximal torus in  $G$  and define the *nodal group*

$$\Gamma(T) := \ker(\exp : \mathfrak{t} \rightarrow T)$$

**Lemma 2.1.** *We have a  $W$ -equivariant isomorphism*

$$X_*(T) \simeq \Gamma(T)$$

*Proof.* The group  $X_*(T)$  is the group of morphisms from the Lie group  $\mathbf{U}$  to the Lie group  $T$ . Given  $f \in X_*(T)$ , we have

$$\Gamma(f) : \Gamma(\mathbf{U}) \rightarrow \Gamma(T).$$

We identify  $\Gamma(\mathbf{U})$  with the subgroup  $2\pi i\mathbb{Z}$  of  $L(\mathbf{U}) = i\mathbb{R}$ . We then have the isomorphism

$$X_*(T) \simeq \Gamma(T), \quad f \mapsto \Gamma(f)(2\pi i)$$

as in [B, p.307]. □

**Lemma 2.2.** *If  $A$  is a locally compact abelian topological group, let  $\hat{A}$  denote its Pontryagin dual. Then we have a  $W$ -equivariant isomorphism*

$$\widehat{\Gamma(T)} \simeq T^\vee.$$

*Proof.* We have, by Lemma (2.1) and the  $T$ -duality equation (3),

$$\Gamma(T) \simeq X_*(T) = X^*(T^\vee) = \widehat{T^\vee}.$$

Now apply Pontryagin duality. □

The groups  $\Gamma(T)$  and  $T^\vee$  are *in duality* in the sense of locally compact abelian topological groups.

**2.5. A table of Langlands dual groups.** The *connection index* is a numerical invariant denoted  $f$  in [B, VI, p.240]. The connection indices are listed in [B, VI, Plates I–X, p.265–292]. The connection index is a useful invariant, thanks to the following property:

$$|\pi_1(G)| \cdot |\mathcal{Z}(G)| = f$$

see [B, IX, p.320]. For example, we have

$$\pi_1 \mathrm{SO}_{2n+1} = \mathbb{Z}/2\mathbb{Z}, \quad \mathcal{Z}(\mathrm{SO}_{2n+1}) = 1, \quad f = 2$$

Here is a table of Langlands duals and connection indices for compact connected semisimple groups:

$G$	$G^\vee$	$f$
$A_n = \mathrm{SU}_{n+1}$	$\mathrm{PSU}_{n+1}$	$n + 1$
$B_n = \mathrm{SO}_{2n+1}$	$\mathrm{Sp}_{2n}$	2
$C_n = \mathrm{Sp}_{2n}$	$\mathrm{SO}_{2n+1}$	2
$D_n = \mathrm{SO}_{2n}$	$\mathrm{SO}_{2n}$	4
$E_6$	$E_6$	3
$E_7$	$E_7$	2
$E_8$	$E_8$	1
$F_4$	$F_4$	1
$G_2$	$G_2$	1

In this table, the simply-connected form of  $E_6$  (resp.  $E_7$ ) corresponds to the adjoint form of  $E_6$  (resp.  $E_7$ ).

### 3. AFFINE WEYL GROUPS

There is a vital distinction between the affine Weyl group  $W_a$  and the extended affine Weyl group  $W'_a$ . The quotient  $W'_a/W_a$  is a finite abelian group which dominates the discussion.

Our reference at this point is [B, IX, p.309–327]. Let  $\mathfrak{t}$  denote the Lie algebra of  $T$ , and let  $\exp : \mathfrak{t} \rightarrow T$  denote the exponential map. The map  $\exp : \mathfrak{t} \rightarrow T$  is a morphism of Lie groups, surjective with discrete kernel [B, p.282]. The kernel of  $\exp$  is the *nodal group*  $\Gamma(T)$ .

The inclusion  $\iota : T \rightarrow G$  induces the homomorphism  $\pi_1(\iota) : \pi_1(T) \rightarrow \pi_1(G)$ . Now  $f(G, T)$  will denote the composite of the canonical isomorphism from  $\Gamma(T)$  to  $\pi_1(T)$  and the homomorphism  $\pi_1(\iota)$ :

$$f(G, T) : \Gamma(T) \simeq \pi_1(T) \rightarrow \pi_1(G).$$

Denote by  $N(G, T)$  the kernel of  $f(G, T)$ . We have a short exact sequence

$$(4) \quad 0 \rightarrow N(G, T) \rightarrow \Gamma(T) \rightarrow \pi_1(G) \rightarrow 0$$

see [B, p.315].

Denote by  $N_G(T)$  the normalizer of  $T$  in  $G$ . Let  $W$  denote the Weyl group  $N_G(T)/T$ . The *affine Weyl group* is

$$W_a = N(G, T) \rtimes W$$

and the *extended affine Weyl group* is

$$W'_a = \Gamma(T) \rtimes W$$

The subgroup  $W_a$  of  $W'_a$  is normal.

If  $\mathfrak{t} - \mathfrak{t}_r$  denotes the union of the singular hyperplanes in  $\mathfrak{t}$ , then the *alcoves* of  $\mathfrak{t}$  are the connected component of  $\mathfrak{t}_r$ .

The group  $W_a$  operates simply-transitively on the set of alcoves. Let  $A$  be an alcove. Then  $\overline{A}$  is a fundamental domain for the operation of  $W_a$  on  $\mathfrak{t}$ .

Let  $H_A$  be the stabilizer of  $A$  in  $W'_a$ . Then  $H_A$  is a finite abelian group which can be identified naturally with  $\pi_1(G)$ , see [B, IX, p.326]. The extended affine Weyl group  $W'_a$  is the semi-direct product

$$W'_a = W_a \rtimes H_A.$$

EXAMPLE. In the special case of  $SU_3$ , the vector space  $\mathfrak{t}$  is the Euclidean plane  $\mathbb{R}^2$ . The singular hyperplanes tessellate  $\mathbb{R}^2$  into equilateral triangles. The interior of each equilateral triangle is an alcove. Barycentric subdivision refines this tessellation into isosceles triangles. The extended affine Weyl group  $W'_a$  acts simply transitively on the set of these isosceles triangles, but the closure  $\overline{\Delta}$  of one such triangle is not a fundamental domain (in the strict sense) for the action of  $W'_a$ . The corresponding quotient space is [B, IX. §5.2]:

$$\mathfrak{t}/W'_a \simeq \overline{A}/H_A.$$

The abelian group  $H_A$  is the cyclic group  $\mathbb{Z}/3\mathbb{Z}$  which acts on  $\overline{A}$  by rotation about the barycentre of  $\overline{A}$  through  $2\pi/3$ .

#### 4. $C^*$ -ALGEBRAS

**Theorem 4.1.** *We have*

$$C^*(W'_a) \simeq C(T^\vee) \rtimes W$$

*Proof.* By Lemma (2.2), the spectrum of the commutative  $C^*$ -algebra  $C^*(\Gamma(T))$  is homeomorphic to the compact Hausdorff space  $T^\vee$ , and we have the *Gelfand* isomorphism [Sp, p.67]:

$$C^*(\Gamma(T)) \simeq C(T^\vee)$$

Related to this, we have, by the Mackey machine:

$$(5) \quad C^*(W'_a) = C^*(\Gamma(T) \rtimes W)$$

$$(6) \quad \simeq C(T^\vee) \rtimes W$$

by Lemma (2.2). □

**Lemma 4.2.** *Let  $\tilde{G}$  denote the universal cover (simply connected covering group) of  $G$  and let  $\tilde{T}$  denote a maximal torus in  $\tilde{G}$ . We have*

$$N(T) = \Gamma(\tilde{T}) = N(\tilde{T})$$

*Proof.* Let  $\pi_1 = \pi_1(G)$ . According to [B, p.291 ], we have

$$G = \tilde{G}/\pi_1, \quad T = \tilde{T}/\pi_1$$

and  $\pi_1$  is central in  $G$ .

Consider the adjoint representation

$$\text{Ad}_{\tilde{G}} : \tilde{G} \rightarrow \text{Aut}(\mathfrak{g})$$

Since  $\pi_1$  is a central subgroup, this representation descends to the adjoint representation of  $G$ :

$$\text{Ad}_G : G \rightarrow \text{Aut}(\mathfrak{g})$$

Since the roots are the nonzero weights in the adjoint representation, it follows that

$$R(\tilde{G}, \tilde{T}) = R(G, T).$$

Now  $N(\tilde{T})$  is the subgroup of  $\mathfrak{t}$  generated by the nodal vectors

$$\{K_\alpha : \alpha \in R(\tilde{G}, \tilde{T})\}$$

and  $N(T)$  is the subgroup of  $\mathfrak{t}$  generated by the nodal vectors

$$\{K_\alpha : \alpha \in R(G, T)\}$$

see [B, p.314]. It follows that

$$N(T) = N(\tilde{T}).$$

By (4) we infer that

$$N(\tilde{T}) = \Gamma(\tilde{T}).$$

□

**Theorem 4.3.** *We have*

$$C^*(W_a) \simeq C(T_{adj}^\vee) \rtimes W$$

where  $T_{adj}^\vee$  is dual to  $\tilde{T}$ , i.e.  $T_{adj}^\vee$  is a maximal torus in the adjoint form of  $G$ .

*Proof.* By Lemma (4.2), we have

$$\begin{aligned} W_a(G) &= N(T) \rtimes W \\ &= \Gamma(\tilde{T}) \rtimes W \\ &= N(\tilde{T}) \rtimes W \end{aligned}$$

so that we have

$$W_a(G) = W'_a(\tilde{G}) = W_a(\tilde{G}).$$

Then we infer that

$$\begin{aligned} C^*(W_a(G)) &= C^*(W'_a(\tilde{G})) \\ &= C(T_{adj}^\vee) \rtimes W \end{aligned}$$

□

5. “ $KK$ -LANGLANDS”

In this section we will establish the isomorphism from  $KK_W^*(C(T), \mathbb{C})$  to  $KK_W^*(\mathbb{C}, C(T^\vee))$ . Poincaré duality (see Kasparov [K] section 4) yields an isomorphism from  $KK_W^*(C(T), \mathbb{C})$  to  $KK_W^*(\mathbb{C}, C(T) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t}))$ . We will show that  $C(T) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t})$  is  $KK$ -equivalent to  $C(T^\vee)$  hence obtaining the required isomorphism from  $KK_W^*(\mathbb{C}, C(T) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t}))$  to  $KK_W^*(\mathbb{C}, C(T^\vee))$ .

**5.1. Construction of the Dirac class.** We begin with the construction of a  $W$ -equivariant Dirac class  $[D]$  in the  $KK$ -group  $KK_W(C(T), C(T^\vee) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t}^*))$ . Let  $L_c^2(\mathfrak{t})$  denote the space of compactly supported square-integrable functions on  $\mathfrak{t}$ , where  $\mathfrak{t}$  has the Haar measure normalised so that  $T = \Gamma \backslash \mathfrak{t}$  has mass 1. This is a  $\Gamma$ -space in the obvious way. We equip the space  $L_c^2(\mathfrak{t})$  with the  $\mathbb{C}[\Gamma]$ -valued inner product

$$\langle u, v \rangle = \sum_{\gamma \in \Gamma} \int_{\mathfrak{t}} \overline{u(x)} (\gamma \cdot v)(x) dx [\gamma]$$

and right  $\mathbb{C}[\Gamma]$ -module structure defined by

$$v[\gamma] = \gamma^{-1} \cdot v.$$

Completing  $L_c^2(\mathfrak{t})$  we obtain a  $C_r^*(\Gamma)$ -Hilbert module  $\overline{L_c^2(\mathfrak{t})}$ , which we view as a  $C(T^\vee)$ -Hilbert module via the isomorphism  $C_r^*(\Gamma) \cong C(T^\vee)$ . In this form the inner product is given explicitly by

$$\langle u, v \rangle(\exp(\eta)) = \sum_{\gamma \in \Gamma} \int_{\mathfrak{t}} \overline{u(x)} (\gamma \cdot v)(x) dx e^{2\pi i \langle \eta, \gamma \rangle}$$

where  $\eta \in \mathfrak{t}^*$ . We note that the inner product can be expressed in a more symmetrical form, at the cost of selecting a fundamental domain  $X$  for the action of  $\Gamma$  on  $\mathfrak{t}$ . The integral over  $\mathfrak{t}$  can be expressed as the sum over  $\delta \in \Gamma$  of the integrals over translates of  $X$ . This gives

$$\int_{\mathfrak{t}} \overline{u(x)} (\gamma \cdot v)(x) dx = \sum_{\delta \in \Gamma} \int_X \overline{\delta \cdot u(x)} ((\delta + \gamma) \cdot v)(x) dx$$

and changing variables to  $\gamma' = \delta + \gamma$  we obtain the formula

$$(7) \quad \langle u, v \rangle(\exp(\eta)) = \sum_{\gamma', \delta \in \Gamma} \int_X \overline{(\delta \cdot u)(x)} e^{2\pi i \langle \eta, \delta \rangle} (\gamma' \cdot v)(x) e^{2\pi i \langle \eta, \gamma' \rangle} dx.$$

Now, having constructed  $\overline{L_c^2(\mathfrak{t})}$ , we form the tensor product Hilbert module  $\mathcal{E} = \overline{L_c^2(\mathfrak{t})} \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t}^*)$  over  $C(T^\vee) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t}^*)$ .

The Weyl group  $W$  acts on  $\mathfrak{t}$  and hence on  $L_c^2(\mathfrak{t})$ , giving rise to an action of  $W$  on the completion  $\overline{L_c^2(\mathfrak{t})}$ . Dually  $W$  acts on  $\mathfrak{t}^*$  and hence on the Clifford algebra  $\mathcal{C}\ell(\mathfrak{t}^*)$ . We equip  $\mathcal{E}$  with the diagonal action of  $W$  on the two factors. It is easy to verify that this makes  $\mathcal{E}$  into a  $W$ -equivariant Hilbert module.

Next we define a representation  $\rho$  of  $C(T)$  as adjointable operators on  $\mathcal{E}$ . Viewing a function  $f$  in  $C(T)$  as a  $\Gamma$ -periodic function  $\tilde{f}$  on  $\mathfrak{t}$ , we simply define  $(\rho(f)(v \otimes a))(x) = \tilde{f}(x)v(x) \otimes a$ . The action of  $W$  on  $\mathfrak{t}$  is (tautologically) compatible with the action on  $T$ , hence this representation is  $W$ -equivariant.

Finally we define the operator  $D$  on  $\mathcal{E}$ . This is an unbounded operator defined (using Einstein summation convention) by

$$D(v \otimes a) = \frac{\partial}{\partial x^j} v \otimes \varepsilon^j a$$

where  $\{\varepsilon^j\}$  denotes the dual basis of  $\mathfrak{t}^*$  corresponding to the basis  $\{\mathbf{e}_j = \frac{\partial}{\partial x^j}\}$  of  $\mathfrak{t}$ .

Self-adjointness of  $D$  follows easily by the usual Stokes' Theorem argument, along with the observation that  $\frac{\partial}{\partial x^j}(\gamma \cdot v) = \gamma \cdot \frac{\partial}{\partial x^j} v$ . For  $f$  in  $C(T)$  differentiable, it is immediate that  $[D, \rho(f)]$  extends to a bounded operator on  $\mathcal{E}$ .

**Proposition 5.1.** *The operator  $D$  on  $\mathcal{E}$  is given by a field over  $T^\vee$  of operators on the Hilbert module  $L^2(X) \hat{\otimes} \mathcal{C}\ell(\mathfrak{t}^*)$ , which have discrete spectrum (with finite multiplicities), namely  $\{\pm 2\pi|\chi + \eta| : \chi \in \Gamma^\vee\}$  at the point  $\exp(\eta)$  in  $T^\vee$ . Hence  $D$  is regular and has compact resolvent.*

*Proof.* Evaluation at a point  $\exp(\eta) \in T^\vee$  gives a map  $C(T^\vee) \hat{\otimes} \mathcal{C}\ell(\mathfrak{t}^*) \rightarrow \mathcal{C}\ell(\mathfrak{t}^*)$ , and we let  $\mathcal{E}_\eta = \mathcal{E} \hat{\otimes}_{C(T^\vee) \hat{\otimes} \mathcal{C}\ell(\mathfrak{t}^*)} \mathcal{C}\ell(\mathfrak{t}^*)$  be the corresponding Hilbert module. Using Equation (7), the map  $\phi_\eta : \mathcal{E}_\eta \rightarrow L^2(X) \otimes \mathcal{C}\ell(\mathfrak{t}^*)$  defined by

$$\phi_\eta((v \otimes a) \otimes 1) = \sum_{\gamma \in \Gamma} v(-\gamma + x) e^{2\pi i \langle \eta, \gamma \rangle} \otimes a$$

preserves the inner product and hence is an isomorphism. We are interested in the localisation of the operator  $D$  given by transferring  $D \otimes 1$  on  $\mathcal{E}_\eta$  onto  $L^2(X) \otimes \mathcal{C}\ell(\mathfrak{t}^*)$ . We will denote the operator on  $L^2(X) \otimes \mathcal{C}\ell(\mathfrak{t}^*)$  by  $D_\eta$ . Like  $D$ , the operator  $D_\eta$  is a differential operator satisfying the local formula  $D_\eta = \frac{\partial}{\partial x^j} \otimes \varepsilon^j$ , however we must also determine the boundary conditions. These are given by considering the image of the smooth functions on  $\mathfrak{t}$  under the map  $\phi_\eta$ : for  $v$  a smooth function we have

$$\begin{aligned} \phi_\eta((v \otimes a) \otimes 1)(\delta + x) &= \sum_{\gamma \in \Gamma} v(-\gamma + \delta + x) e^{2\pi i \langle \eta, \gamma \rangle} \otimes a \\ &= \sum_{\gamma' \in \Gamma} v(-\gamma' + x) e^{2\pi i \langle \eta, \delta + \gamma' \rangle} \otimes a \end{aligned}$$

so  $\phi_\eta((v \otimes a) \otimes 1)(\delta + x) = e^{2\pi i \langle \eta, \delta \rangle} \phi_\eta((v \otimes a) \otimes 1)(x)$ . We note that the boundary conditions vary with the point  $\eta$ , however this variation is  $\Gamma^\vee$ -periodic, so in fact they depend only on  $\exp(\eta)$ .

Fixing  $\eta$ , the space  $L^2(X)$  has an orthonormal basis consisting of functions of the form  $e^{2\pi i \langle \chi + \eta, x \rangle}$  where  $\chi$  ranges over the dual lattice  $\widehat{T} = \Gamma^\vee$ . These

functions satisfy the boundary conditions. Applying the operator  $D_\eta$  and then pulling the coordinates  $\chi_j + \eta_j$  through the tensor we obtain

$$D_\eta(e^{2\pi i \langle \chi + \eta, x \rangle} \otimes b) = \frac{\partial}{\partial x^j} e^{2\pi i \langle \chi + \eta, x \rangle} \otimes \varepsilon^j b = 2\pi i e^{2\pi i \langle \chi + \eta, x \rangle} \otimes (\chi + \eta) b$$

whence the operator  $D_\eta$  is  $2\pi i \otimes (\chi + \eta)$  on the corresponding subspace. The submodules

$$E_{\chi, \pm} = \{e^{2\pi i \langle \chi + \eta, x \rangle} \otimes (i(\chi + \eta) \pm |\chi + \eta|)a : a \in \mathcal{C}\ell(\mathfrak{t}^*)\}$$

are eigenspaces with eigenvalue  $\pm 2\pi |\chi + \eta|$ . For each  $\eta$  the set of  $\chi$  such that  $|\chi + \eta|$  takes a given value is finite, hence the eigenvalues have finite multiplicity.

It now follows easily that  $D$  is regular and has compact resolvent.  $\square$

To show that the triple  $(\mathcal{E}, \rho, D)$  is an unbounded  $W$ -equivariant Kasparov triple, thereby defining an element of  $KK_W(C(T), C(T^\vee) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t}^*))$ , it remains to show that the Dirac operator  $D$  is  $W$ -equivariant. This follows from a more general statement, Proposition 5.2 below, which shows in particular that  $\frac{\partial}{\partial x^j} \otimes \varepsilon^j$  in  $\mathfrak{t} \otimes \mathfrak{t}^*$  is invariant under the natural action of  $\mathrm{GL}(\mathfrak{t})$  and hence that it is  $W$ -invariant. The Proposition moreover shows an invariance result for elements of  $\mathcal{C}\ell(\mathfrak{t}) \otimes \mathcal{C}\ell(\mathfrak{t}^*)$  which we will make use of later to carry out a  $W$ -equivariant restriction from the Clifford algebras to spinors.

Consider the abstract setup of a finite dimensional vector space  $V$ . The natural action of  $\mathrm{GL}(V)$  on  $V$ , induces a diagonal action on  $V \otimes V^*$ .

If  $V$  is equipped with a non-degenerate symmetric bilinear form  $g$  then we can form the Clifford algebra  $\mathcal{C}\ell(V)$ . The subgroup  $\mathrm{O}(g)$  of  $\mathrm{GL}(V)$ , consisting of those elements preserving  $g$ , acts naturally on  $\mathcal{C}\ell(V)$ . The bilinear form additionally gives an isomorphism from  $V$  to  $V^*$  and hence induces a bilinear form  $g^*$  on  $V^*$ , allowing us to form the Clifford algebra  $\mathcal{C}\ell(V^*)$ . Clearly the dual action of  $\mathrm{O}(g)$  on  $V^*$  preserves  $g^*$  hence there is a diagonal action of  $\mathrm{O}(g)$  on  $\mathcal{C}\ell(V) \widehat{\otimes} \mathcal{C}\ell(V^*)$  which we identify with  $\mathcal{C}\ell(V \times V^*)$ .

We say that an element  $a$  of  $\mathcal{C}\ell(V \times V^*)$  is *symmetric* if there exists a  $g$ -orthonormal<sup>1</sup> basis  $\{\mathbf{e}_j : j = 1, \dots, n\}$  with dual basis  $\{\varepsilon^j : j = 1, \dots, n\}$  such that  $a$  can be written as  $p(\mathbf{e}_1 \varepsilon^1, \dots, \mathbf{e}_n \varepsilon^n)$  where  $p(x_1, \dots, x_n)$  is a symmetric polynomial.

**Proposition 5.2.** *For any basis  $\{\mathbf{e}_j\}$  of  $V$  with dual basis  $\{\varepsilon^j\}$  for  $V^*$ , the Einstein sum  $\mathbf{e}_j \otimes \varepsilon^j$  in  $V \otimes V^*$  is  $\mathrm{GL}(V)$ -invariant.*

*Suppose moreover that  $V$  is equipped with a non-degenerate symmetric bilinear form  $g$  and that the underlying field has characteristic zero. Then every symmetric element of  $\mathcal{C}\ell(V) \widehat{\otimes} \mathcal{C}\ell(V^*) \cong \mathcal{C}\ell(V \times V^*)$  is  $\mathrm{O}(g)$ -invariant.*

<sup>1</sup>We say that  $\{\mathbf{e}_j\}$  is  $g$ -orthonormal if  $g_{jk} = \pm \delta_{jk}$  for each  $j, k$ .

*Proof.* Identifying  $V \otimes V^*$  with endomorphisms of  $V$  in the natural way, the action of  $\mathrm{GL}(V)$  is the action by conjugation and  $\mathbf{e}_j \otimes \varepsilon^j$  is the identity which is invariant under conjugation.

For the second part, over a field of characteristic zero the symmetric polynomials are generated by power sum symmetric polynomials  $p(x_1, \dots, x_n) = x_1^k + \dots + x_n^k$ , so it suffices to consider

$$\begin{aligned} p(\mathbf{e}_1 \varepsilon^1, \dots, \mathbf{e}_n \varepsilon^n) &= (\mathbf{e}_1 \varepsilon^1)^k + \dots + (\mathbf{e}_n \varepsilon^n)^k \\ &= (-1)^{k(k-1)/2} \left( (\mathbf{e}_1)^k (\varepsilon^1)^k + \dots + (\mathbf{e}_n)^k (\varepsilon^n)^k \right). \end{aligned}$$

When  $k$  is even, writing  $(\mathbf{e}_j)^k = (\mathbf{e}_j^2)^{k/2} = (g_{jj})^{k/2}$  and similarly  $(\varepsilon^j)^k = (g^{jj})^{k/2}$ , we see that each term  $(\mathbf{e}_j)^k (\varepsilon^j)^k$  is 1 since  $g_{jj} = g^{jj} = \pm 1$  for an orthonormal basis. Thus  $p(\mathbf{e}_1 \varepsilon^1, \dots, \mathbf{e}_n \varepsilon^n) = n(-1)^{k(k-1)/2}$  which is invariant.

Similarly when  $k$  is odd we get  $(\mathbf{e}_j)^k (\varepsilon^j)^k = \mathbf{e}_j \varepsilon^j$  so

$$p(\mathbf{e}_1 \varepsilon^1, \dots, \mathbf{e}_n \varepsilon^n) = (-1)^{k(k-1)/2} (\mathbf{e}_1 \varepsilon^1 + \dots + \mathbf{e}_n \varepsilon^n).$$

As the sum  $\mathbf{e}_j \otimes \varepsilon^j$  in  $V \otimes V^*$  is invariant under  $\mathrm{GL}(V)$ , it is in particular invariant under  $\mathrm{O}(g)$ , and hence the sum  $\mathbf{e}_j \varepsilon^j$  is  $\mathrm{O}(g)$ -invariant in the Clifford algebra.  $\square$

**5.2. The Kasparov product.** In the previous section we constructed a class  $(\mathcal{E}, \rho, D)$  in  $KK_W(C(T), C(T^\vee) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t}^*))$ . Replacing the group  $G$  with its Langlands dual this construction produces an element  $(\mathcal{E}^\vee, \rho^\vee, D^\vee)$  in  $KK_W(C(T^\vee), C(T) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t}))$ . Taking the Kasparov product with  $(\mathcal{E}^\vee, \rho^\vee, D^\vee)$  gives a map

$$KK_W(\mathbb{C}, C(T^\vee)) \rightarrow KK_W(\mathbb{C}, C(T) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t}))$$

which we will prove is an isomorphism. To construct the inverse, first we take the Kasparov product with  $(\mathcal{E}, \rho, D)$  over  $C(T)$  which gives a map

$$KK_W(\mathbb{C}, C(T) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t})) \rightarrow KK_W(\mathbb{C}, C(T^\vee) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t}^*) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t}))$$

and then use a  $W$ -equivariant Morita equivalence to get a map

$$KK_W(\mathbb{C}, C(T^\vee) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t}^*) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t})) \rightarrow KK_W(\mathbb{C}, C(T^\vee)).$$

The composition of these three maps gives a map

$$(2) \quad KK_W(\mathbb{C}, C(T^\vee)) \rightarrow KK_W(\mathbb{C}, C(T^\vee))$$

which we will show is the identity.

Consider the projection  $P = \prod_j \frac{1}{2}(1 + i\mathbf{e}_j \varepsilon^j)$  in the Clifford algebra  $\mathcal{C}\ell(\mathfrak{t}^*) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t}) \cong \mathcal{C}\ell(\mathfrak{t}^* \times \mathfrak{t})$ . By Proposition 5.2 this is  $W$ -invariant. The corner algebra  $P\mathcal{C}\ell(\mathfrak{t}^* \times \mathfrak{t})P$  is  $\mathbb{C}P$ , and we will identify this with  $\mathbb{C}$ . Now let  $\mathcal{S} = \mathcal{C}\ell(\mathfrak{t}^* \times \mathfrak{t})P$ , which is a finite dimensional Hilbert space, with inner product given by  $\langle aP, bP \rangle = Pa^*bP$ . This is naturally equipped with a representation of  $\mathcal{C}\ell(\mathfrak{t}^* \times \mathfrak{t})$  as adjointable operators, indeed  $\mathcal{C}\ell(\mathfrak{t}^* \times \mathfrak{t})$  is identified

with the algebra of compact operators on  $\mathcal{S}$ . Thus  $\mathcal{S}$  gives a  $W$ -equivariant graded Morita equivalence from  $\mathcal{C}\ell(\mathfrak{t}^* \times \mathfrak{t})$  to  $\mathbb{C}$ .

Our inverse to the element  $(\mathcal{E}^\vee, \rho^\vee, D^\vee)$  is thus given by the  $\mathcal{S}$ -spinor reduction of  $(\mathcal{E}, \rho, D)$ , viz.

$$((\mathcal{E} \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t})) \widehat{\otimes}_{\mathcal{C}\ell(\mathfrak{t}^*) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t})} \mathcal{S}, \rho \otimes 1 \otimes 1, D \otimes 1 \otimes 1).$$

The module  $(\mathcal{E} \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t})) \widehat{\otimes}_{\mathcal{C}\ell(\mathfrak{t}^*) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t})} \mathcal{S}$  simplifies as  $\overline{L_c^2(\mathfrak{t})} \widehat{\otimes} \mathcal{S}$ , with the representation of  $C(T)$  acting on  $\overline{L_c^2(\mathfrak{t})}$ , and with the Dirac operator now acting on spinor fields. By a slight abuse of notation we will denote the representation and operator on  $\overline{L_c^2(\mathfrak{t})} \widehat{\otimes} \mathcal{S}$  by  $\rho \otimes 1$  and  $D \otimes 1$  respectively.

We will now proceed to compute the Kasparov product

$$(\mathcal{E}^\vee, \rho^\vee, D^\vee) \otimes_{C(T) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t})} (\overline{L_c^2(\mathfrak{t})} \widehat{\otimes} \mathcal{S}, \rho \otimes 1, D \otimes 1)$$

corresponding to the map (8).

The first step is to consider the Hilbert module for this Kasparov product which is the following  $C(T^\vee)$ -Hilbert module:

$$\mathcal{E}^\vee \widehat{\otimes}_{C(T) \widehat{\otimes} \mathcal{C}\ell(\mathfrak{t})} (\overline{L_c^2(\mathfrak{t})} \widehat{\otimes} \mathcal{S}) \cong \overline{L_c^2(\mathfrak{t}^*)} \widehat{\otimes}_{C(T)} \overline{L_c^2(\mathfrak{t})} \widehat{\otimes} \mathcal{S}.$$

We remark that the group  $W$  acts diagonally on all three factors of this tensor product, and indeed that the action on  $\mathcal{S}$  is diagonal in terms of the  $\mathfrak{t}^*$  and  $\mathfrak{t}$  parts.

We define a map  $\phi$  from  $\overline{L_c^2(\mathfrak{t}^*)} \widehat{\otimes}_{C(T)} \overline{L_c^2(\mathfrak{t})}$  to the Hilbert module  $L^2(\mathfrak{t}^*) \widehat{\otimes} C(T^\vee)$  by

$$\phi(u \otimes v)(\eta, \exp(\zeta)) = \int_T \sum_{\chi \in \Gamma^\vee} \sum_{\gamma \in \Gamma} u(-\chi + \eta + \zeta) v(-\gamma + x) e^{2\pi i \langle \chi - \zeta, -\gamma + x \rangle} dx.$$

We note that the summation over  $\chi$  ensures that this function is  $\Gamma^\vee$ -periodic in the  $\zeta$  variable and thus depends only on the point  $\exp(\zeta) \in T^\vee$ , not the chosen representative  $\zeta$  in  $\mathfrak{t}^*$ .

**Lemma 5.3.** *The map  $\phi$  preserves the inner product and gives an isomorphism of Hilbert modules*

$$\phi : \overline{L_c^2(\mathfrak{t}^*)} \widehat{\otimes}_{C(T)} \overline{L_c^2(\mathfrak{t})} \rightarrow L^2(\mathfrak{t}^*) \widehat{\otimes} C(T^\vee).$$

*Proof.* We need to compute

$$\langle u \otimes v, u' \otimes v' \rangle = \langle v, \langle u, u' \rangle v' \rangle.$$

The function  $\langle u, u' \rangle$  is given by

$$\langle u, u' \rangle(x) = \sum_{\chi \in \Gamma^\vee} \int_{\mathfrak{t}^*} \overline{u(\eta)} u'(-\chi + \eta) d\eta e^{2\pi i \langle \chi, x \rangle}$$

which is  $\Gamma$ -periodic. Now choose a fundamental domain  $X$  for the action of  $\Gamma$  on  $\mathfrak{t}$ , and for  $x \in X$  define  $w(x, \zeta) = \sum_{\gamma \in \Gamma} v(-\gamma + x) e^{2\pi i \langle \zeta, \gamma \rangle}$ . We similarly

define  $w'(x, \zeta)$  in terms of  $v'(x)$ . We have (using Equation 7),

$$\begin{aligned} \langle v, \langle u, u' \rangle v' \rangle(\exp(\zeta)) &= \int_X \overline{w(x, \zeta)} (\langle u, u' \rangle(x) w'(x, \zeta)) dx \\ &= \int_{\mathfrak{t}^*} \sum_{\chi \in \Gamma^\vee} \int_X \overline{w(x, \zeta) u(\eta)} u'(-\chi + \eta) e^{2\pi i \langle \chi, x \rangle} w'(x, \zeta) dx d\eta \end{aligned}$$

changing the orders of integration and summation. The inner integral is an inner product in  $L^2(X)$ . For each  $\zeta$  this space has orthonormal basis consisting of functions of the form  $e^{2\pi i \langle -\psi + \zeta, x \rangle}$ , where  $\psi$  ranges over the dual lattice  $\widehat{T} = \Gamma^\vee$ , and hence the integral can be expressed as an inner product in  $\ell^2(\Gamma^\vee)$  by Parseval's identity:

$$\begin{aligned} &\int_X \overline{w(x, \zeta) u(\eta)} u'(-\chi + \eta) e^{2\pi i \langle \chi, x \rangle} w'(x, \zeta) dx \\ &= \sum_{\psi \in \Gamma^\vee} \int_X \overline{w(x, \zeta) u(\eta) e^{2\pi i \langle \psi - \zeta, x \rangle}} dx \int_X w'(x', \zeta) u(-\chi + \eta) e^{2\pi i \langle \chi, x' \rangle} e^{2\pi i \langle \psi - \zeta, x' \rangle} dx' \end{aligned}$$

We now substitute  $\chi' = \psi + \chi$  for  $\chi$  and  $\eta' = \psi + \eta$  for  $\eta$  to get

$$\begin{aligned} &\langle v, \langle u, u' \rangle v' \rangle(\exp(\zeta)) \\ &= \int_{\mathfrak{t}^*} \sum_{\psi \in \Gamma^\vee} \int_X \overline{w(x, \zeta) u(-\psi + \eta') e^{2\pi i \langle \psi - \zeta, x \rangle}} dx \sum_{\chi' \in \Gamma^\vee} \int_X w'(x', \zeta) u(-\chi' + \eta') e^{2\pi i \langle \chi' - \zeta, x' \rangle} dx' d\eta' \end{aligned}$$

Finally, substituting in the sums defining  $w(x, \zeta)$  and  $w'(x', \zeta)$ , and noting that  $e^{2\pi i \langle \psi, -\gamma \rangle} = e^{2\pi i \langle \chi', -\gamma \rangle} = 1$  we get

$$\langle v, \langle u, u' \rangle v' \rangle(\exp(\zeta)) = \int_{\mathfrak{t}^*} \overline{\phi(u \otimes v)(\eta', \exp(\zeta))} \phi(u' \otimes v')(\eta', \exp(\zeta)) d\eta'$$

as required.

To show surjectivity of  $\phi_\zeta$ , let  $u \in L_c^2(\mathfrak{t}^*)$  and for  $\delta \in \Gamma$  let  $v \in L_c^2(\mathfrak{t})$  be defined by

$$v(x) = \begin{cases} e^{2\pi i \langle \zeta, x \rangle} & \text{for } x \in -\delta + X \\ 0 & \text{otherwise.} \end{cases}$$

Then

$$\begin{aligned} \int_X \sum_{\gamma \in \Gamma} v(-\gamma + x) e^{2\pi i \langle \chi - \zeta, -\gamma + x \rangle} dx &= \int_X e^{2\pi i \langle \zeta, x \rangle} e^{2\pi i \langle \chi - \zeta, -\delta + x \rangle} dx \\ &= \int_X e^{2\pi i \langle \chi, x \rangle} e^{2\pi i \langle \zeta, \delta \rangle} dx \end{aligned}$$

since  $e^{2\pi i \langle \chi, -\delta \rangle} = 1$ . This vanishes unless  $\chi = 0$  in which case it is  $e^{2\pi i \langle \zeta, \delta \rangle}$ , hence  $\phi_\zeta(u \otimes v) = u \otimes e^{2\pi i \langle \zeta, \delta \rangle}$ . The image of  $\phi$  is thus dense, hence  $\phi$  must be surjective.  $\square$

The algebra  $C(T^\vee)$  is represented on  $\overline{L_c^2(\mathfrak{t}^*)} \hat{\otimes}_{C(T)} \overline{L_c^2(\mathfrak{t})}$  by pointwise multiplication on the first factor, viewing elements  $f$  in  $C(T^\vee)$  as periodic functions  $\tilde{f}$  on  $\mathfrak{t}^*$ . Applying  $\phi$  we have

$$\begin{aligned} \phi(\tilde{f}u \otimes v)(\eta, \exp(\zeta)) &= \int_T \sum_{\chi \in \Gamma^\vee} \sum_{\gamma \in \Gamma} \tilde{f}(-\chi + \eta + \zeta) u(-\chi + \eta + \zeta) v(-\gamma + x) e^{2\pi i \langle \chi - \zeta, -\gamma + x \rangle} dx \\ &= \tilde{f}(\eta + \zeta) \phi(u \otimes v)(\eta, \exp(\zeta)). \end{aligned}$$

The representation of  $C(T^\vee)$  on  $\overline{L_c^2(\mathfrak{t}^*)} \hat{\otimes}_{C(T)} \overline{L_c^2(\mathfrak{t})}$  thus corresponds to the representation  $\sigma$  of  $C(T^\vee)$  on  $L^2(\mathfrak{t}^*) \hat{\otimes} C(T^\vee)$  defined by pointwise multiplication with  $\tilde{f}(\eta + \zeta)$ . Note that for  $\tilde{f}$  of the form  $\tilde{f}(\xi) = e^{2\pi i \langle \xi, \gamma \rangle}$ , for  $\gamma \in \Gamma$  we have  $\tilde{f}(\eta + \zeta) = \tilde{f}(\eta) \tilde{f}(\zeta)$  so the representation  $\sigma$  is ‘diagonal’ on such functions.

We will now define an unbounded operator on the Hilbert module of spinor-valued functions  $L^2(\mathfrak{t}^*) \hat{\otimes} C(T^\vee) \hat{\otimes} \mathcal{S}$ . We define the operator  $\hat{D}$  by the formula

$$\hat{D}(u \otimes g \otimes s)(\eta, \exp(\zeta)) = \eta_j u(\eta) \otimes g(\exp(\zeta)) \otimes \varepsilon^j s$$

where  $\eta_j$  denotes the  $j$ th coordinate of  $\eta$ , and we are using Einstein summation convention. We can alternatively write  $\eta_j = \langle \eta, \mathbf{e}_j \rangle$ , hence  $W$ -invariance of  $\hat{D}$  follows from the  $W$ -invariance of  $\mathbf{e}_j \otimes \varepsilon^j$  (Proposition 5.2).

Recall that  $D \otimes 1$  is an unbounded operator on  $\overline{L_c^2(\mathfrak{t})} \hat{\otimes} \mathcal{S}$ . Identifying the Hilbert module  $\mathcal{E}^\vee \hat{\otimes}_{C(T) \hat{\otimes} \mathcal{C}\ell(\mathfrak{t})} (\overline{L_c^2(\mathfrak{t})} \hat{\otimes} \mathcal{S})$  with  $L^2(\mathfrak{t}^*) \hat{\otimes} C(T^\vee) \hat{\otimes} \mathcal{S}$ , the operator  $\hat{D}$  is an unbounded connection for  $D \otimes 1$  in the following sense.

**Lemma 5.4.** *The bounded operator  $\hat{D}(1 + \hat{D}^2)^{-1/2}$  is a  $D(1 + D^2)^{-1/2} \otimes 1$ -connection.*  $\square$

It now follows that the Kasparov product is represented by the unbounded triple

$$(L^2(\mathfrak{t}^*) \hat{\otimes} C(T^\vee) \hat{\otimes} \mathcal{S}, \sigma \otimes 1, D^\vee \otimes 1 + \hat{D})$$

where the operator is given by

$$D^\vee \otimes 1 + \hat{D} = \frac{\partial}{\partial \eta_j} \otimes 1 \otimes \mathbf{e}_j + \eta_j \otimes 1 \otimes \varepsilon^j.$$

**Theorem 5.5.** *The Kasparov product*

$$(\mathcal{E}^\vee, \rho^\vee, D^\vee) \otimes_{C(T) \hat{\otimes} \mathcal{C}\ell(\mathfrak{t})} (\overline{L_c^2(\mathfrak{t})} \hat{\otimes} \mathcal{S}, \rho \otimes 1, D \otimes 1)$$

*is the identity in  $KK_W(C(T^\vee), C(T^\vee))$*

*Proof.* We have seen that the Kasparov product is represented by the Kasparov triple

$$(L^2(\mathfrak{t}^*) \hat{\otimes} C(T^\vee) \hat{\otimes} \mathcal{S}, \sigma \otimes 1, D^\vee \otimes 1 + \hat{D}).$$

We now define a family of representations  $\sigma_t$  of  $C(T^\vee)$  on  $L^2(\mathfrak{t}^*) \hat{\otimes} C(T^\vee)$  by

$$\sigma_t(f)(u \otimes g) = \tilde{f}(t\eta + \zeta)(u \otimes g)$$

where as usual  $\tilde{f}$  is the pull-back of  $f$  to  $\mathfrak{t}^*$ . We thus obtain defines a homotopy from the above triple (where  $\sigma = \sigma_1$ ) to

$$(L^2(\mathfrak{t}^*) \hat{\otimes} C(T^\vee) \hat{\otimes} \mathcal{S}, \sigma_0 \otimes 1, D^\vee \otimes 1 + \hat{D}).$$

This decouples the operator, which is constant in the  $T^\vee$ -variable, from the representation which is now the identity representation of  $C(T^\vee)$  on itself.

A standard ‘ladder operator’ calculation shows that the operator  $D^\vee \otimes 1 + \hat{D}$ , which we can view as an operator on the graded Hilbert space  $(L^2(\mathfrak{t}^*) \hat{\otimes} \mathcal{S})$ , is Fredholm with  $\mathbb{Z}/2$ -graded index 1. Specifically the operator has 1-dimensional kernel yielding a splitting of the Hilbert module  $L^2(\mathfrak{t}^*) \hat{\otimes} C(T^\vee) \hat{\otimes} \mathcal{S}$  into a copy of  $C(T^\vee)$  and a complement of this on which the triple is degenerate.

Since the representation  $\sigma_0$  acts by the identity on  $C(T^\vee)$  we see that the non-degenerate piece is given by the triple  $(C(T^\vee), \text{id}, 0)$  which is the identity in  $KK_W(C(T^\vee), C(T^\vee))$   $\square$

**5.3. The KK-equivalence.** To complete our proof of the isomorphism

$$KK_W(\mathbb{C}, C(T^\vee)) \cong KK_W(\mathbb{C}, C(T) \hat{\otimes} \mathcal{C}\ell(\mathfrak{t}))$$

we prove the following.

**Theorem 5.6.** *The class  $(\mathcal{E}^\vee, \rho^\vee, D^\vee)$  gives a KK-equivalence from  $C(T^\vee)$  to  $C(T) \hat{\otimes} \mathcal{C}\ell(\mathfrak{t})$ .*

*Proof.* For brevity of notation we will write  $[D]$  for the class of  $(\mathcal{E}, \rho, D)$  and  $[D^\vee]$  for the class of  $(\mathcal{E}^\vee, \rho^\vee, D^\vee)$ . We will write  $[\mathcal{S}]$  for the KK-class corresponding to the Morita equivalence  $\mathcal{S} = \mathcal{C}\ell(\mathfrak{t}^* \times \mathfrak{t})P$  from  $\mathcal{C}\ell(\mathfrak{t}^* \times \mathfrak{t})$  to  $\mathbb{C}$ , and  $[\mathcal{S}^*]$  for its inverse, which is given by the module  $\mathcal{S}^* = P\mathcal{C}\ell(\mathfrak{t}^* \times \mathfrak{t})$ .

We showed in the previous section (Theorem 5.5) that  $[D^\vee] \otimes [D] \otimes [\mathcal{S}] = 1_{C(T^\vee)}$ .

On the other hand, replacing the group  $G$  by its Langlands dual  $G^\vee$ , Theorem 5.5 shows that  $[D] \otimes [D^\vee] \otimes [\mathcal{S}] = 1_{C(T)}$ , hence  $[D] \otimes [D^\vee] = 1_{C(T)} \otimes [\mathcal{S}^*]$ . We note that  $\mathcal{S}$  and  $\mathcal{S}^*$  remain unchanged (up to isomorphism) by the Langlands duality.

We conclude that the product of  $1_{C(T)} \otimes [\mathcal{S}]$  with  $[D] \otimes [D^\vee]$  gives  $1_{C(T)}$  hence  $[D^\vee]$  has a left inverse  $(1_{C(T)} \otimes [\mathcal{S}]) \otimes [D]$  as well as right inverse  $[D] \otimes [\mathcal{S}]$ . Thus in fact these two elements of  $KK_W(C(T) \hat{\otimes} \mathcal{C}\ell(\mathfrak{t}), C(T^\vee))$  are equal and  $[D^\vee]$  is invertible as required.  $\square$

**Corollary 5.7.** *There is an isomorphism from  $KK_W(C(T), \mathbb{C})$  to  $KK_W(\mathbb{C}, C(T^\vee))$  given by the composition of Poincaré duality with the Langlands KK-equivalence.*

## 6. NOTES AND COMPUTATIONS

It may be helpful to outline the main ideas in §5, omitting the technicalities.

**Theorem 6.1.** *As well as the duality between  $K$ -theory and  $K$ -homology, there is also a Langlands duality in the Baum-Connes correspondence for the extended affine Weyl group  $W'_a$ , namely*

$$(9) \quad K_*^W(T) \simeq K_W^*(T^\vee)$$

where  $W$  is the finite Weyl group.

*Proof.* The real vector space  $\mathfrak{t}$  is a contractible space on which  $W_a$  and  $W'_a$  act properly, and serves as universal example for the (extended) affine Weyl group:

$$\mathfrak{t} = \underline{E}W_a = \underline{E}W'_a$$

The LHS of (9) relies on the fact that the lattice  $\Gamma(T)$  acts freely on the Lie algebra  $\mathfrak{t}$ . Then we have

$$\begin{aligned} K_*^{W'_a}(\mathfrak{t}) &\simeq K_*^W(\mathfrak{t}/\Gamma(T)) \\ &\simeq K_*^W(T) \end{aligned}$$

For the RHS of (9) we have

$$\begin{aligned} K_*(C^*(W'_a)) &\simeq C(T^\vee) \rtimes W \\ &\simeq K_W^*(T^\vee) \end{aligned}$$

by Theorem (4.1) and Eqn.(10).  $\square$

**6.1. The extended quotient.** Let  $K_W^j(T)$  denote the classical topological equivariant  $K$ -theory [A, 2.3] for the Weyl group  $W$  acting on the compact torus  $T$ .

By the Green-Julg theorem [Black, Theorem 11.7.1], we have

$$(10) \quad K_j(C(T^\vee) \rtimes W) \simeq K_W^j(T^\vee)$$

Applying the equivariant Chern character for discrete groups [BC] gives a map

$$(11) \quad \text{ch}_W : K_W^j(T^\vee) \rightarrow \bigoplus_l H^{j+2l}(T^\vee // W; \mathbb{C})$$

which becomes an isomorphism when  $K_W^j(T^\vee)$  is tensored with  $\mathbb{C}$ . In fact, it is enough to tensor by  $\mathbb{Q}(\zeta)$  where  $\zeta$  is a primitive  $d$ th root of unity, where  $d$  is the order of  $W$ .

In the formula (11) for the equivariant Chern character,  $T^\vee // W$  denotes the extended quotient of  $T^\vee$  by  $W$ . This, for any  $W$ -space  $X$ , is defined as follows. Set

$$\tilde{X} := \{(w, t) \in W \times X : wt = t\}.$$

Then  $\tilde{X} \subset W \times X$ . The group  $W$  acts on  $\tilde{X}$ :

$$W \times \tilde{X} \rightarrow \tilde{X}, \quad \alpha(w, t) = (\alpha w \alpha^{-1}, \alpha t)$$

with  $(w, t) \in \tilde{X}, \alpha \in W$ . The extended quotient is defined by

$$X//W := \tilde{X}/W.$$

The extended quotient  $X//W$  is the ordinary quotient for the action of  $W$  on  $\tilde{X}$ .

**6.2. The quotient  $T/W$ .** View  $\mathfrak{t}$  as an additive group, and form the Euclidean group  $\mathfrak{t} \rtimes O(\mathfrak{t})$ . We have  $W'_a \subset \mathfrak{t} \rtimes O(\mathfrak{t})$  and so  $W'_a$  acts as affine transformations of  $\mathfrak{t}$ . Now  $H_A$  leaves  $\bar{A}$  invariant, so  $H_A$  acts as affine transformations of  $\bar{A}$ . Let  $v_0, v_1, \dots, v_n$  be the vertices of the simplex  $\bar{A}$ . We will use barycentric coordinates, so that

$$x = \sum_{i=0}^n t_i v_i$$

with  $x \in \bar{A}$ . The barycentre  $x_0$  has coordinates  $t_j = 1, 0 \leq j \leq n$  and so is  $H_A$ -fixed. Then  $\bar{A}$  is equivariantly contractible to  $x_0$ :

$$(12) \quad r_t(x) := tx_0 + (1-t)x$$

with  $0 \leq t \leq 1$ . This is an affine  $H_A$ -equivariant retract from  $\bar{A}$  to  $x_0$ .

The exponential map  $\bar{A} \rightarrow T$  and the canonical injection  $T \rightarrow G$  induce a homeomorphism

$$\bar{A}/H_A \rightarrow T/W$$

see [B, p.326]. It follows that  $T/W$  is a contractible space.

**6.3. The extended affine Weyl group attached to  $\text{PSU}_3$ .** Let  $G = \text{PSU}_3$ , then  $G^\vee = \text{SU}_3$ . Then  $\pi_1(G) = \mathbb{Z}/3\mathbb{Z}$  and so  $W_a$  has index 3 in  $W'_a$ . The dual torus  $T^\vee$  is the standard maximal torus in  $\text{SU}_3$ . The finite Weyl group is the symmetric group  $\mathfrak{S}_3$ . Computing the extended quotient  $T//W$  we find

$$T^\vee//W = T^\vee/W \sqcup (T^\vee)^{s_1}/Z(s_1) \sqcup (T^\vee)^{s_1 s_2}/Z(s_1 s_2)$$

Now

- $T^\vee/W$  is contractible as in §6.2
  - the second term is a copy of  $\mathbf{U}$
  - the third term is the set  $\{I, \omega I, \omega^2 I\}$  with  $\omega$  a cube root of unity
- By Theorem (4.1), we have (modulo torsion)

$$K_0 C^*(W'_a) = \mathbb{Z}^5, \quad K_1 C^*(W'_a) = \mathbb{Z}$$

We obtain isomorphisms after tensoring over  $\mathbb{Z}$  with  $\mathbb{Q}(e^{2\pi i/6})$ .

**6.4. The affine Weyl group attached to  $\mathrm{PSU}_3$ .** We stay with  $G = \mathrm{PSU}_3$ ,  $G^\vee = \mathrm{SU}_3$ . For  $W_a$  we need the maximal torus in the adjoint form of  $\mathrm{SU}_3$ , namely  $\mathrm{PSU}_3$ . Denote by  $S$  the standard maximal torus in  $\mathrm{PSU}_3$ . In homogeneous coordinates, we have

$$S = \{(z_1 : z_2 : z_3) : z_j \in \mathbf{U}\}$$

We obtain

$$S//W = S/W \sqcup S^{s_1}/Z(s_1) \sqcup S^{s_1 s_2}/Z(s_1 s_2)$$

Now

- $S/W$  is contractible as in §6.2
- the second term is a copy of  $\mathbf{U}$
- the third term is the set  $(1 : 1 : 1) \sqcup \{1 : \omega : \omega^2\} \sqcup \{1 : \omega^2 : \omega\}$  with  $\omega$  a cube root of unity

By Theorem (4.3), we have (modulo torsion)

$$K_0 C^*(W_a) = \mathbb{Z}^5, \quad K_1 C^*(W_a) = \mathbb{Z}$$

**6.5. The affine Weyl group attached to  $\mathrm{SU}_3$ .** Let  $G = \mathrm{SU}_3$ . The group  $\mathrm{SU}_3$  is simply connected and  $W_a = W'_a$ . We have  $G^\vee = \mathrm{PSU}_3$  and  $T^\vee = S$  in the above notation. Modulo torsion, we have

$$K_0 C^*(W_a) = K_0 C^*(W'_a) = \mathbb{Z}^5, \quad K_1 C^*(W_a) = K_1 C^*(W'_a) = \mathbb{Z}$$

**6.6. The affine Weyl group attached to  $G_2$ .** The exceptional Lie group  $G_2$  has connection index 1, and so its compact real form is unique: it is simply connected and of adjoint type. The maximal torus has dimension 2, and the Weyl group  $W$  is the dihedral group of order 12.

The extended quotient  $T//W$  may be computed directly. The extended quotient has 8 connected components:

$$T//W = \mathrm{pt} \sqcup \mathrm{pt} \sqcup \mathrm{pt} \sqcup \mathrm{pt} \sqcup \mathrm{pt} \sqcup \mathbb{I} \sqcup \mathbb{I} \sqcup T/W$$

where  $\mathbb{I}$  is the closed unit interval. For the ordinary quotient, we have

$$T/W \simeq \overline{A}$$

which is a contractible space. The space  $\mathbb{I}$  is contractible. So we have a homotopy equivalence

$$T//W \sim 8 \text{ isolated points}$$

so we have (modulo torsion)

$$K_0 = \mathbb{Z}^8, \quad K_1 = 0$$

## REFERENCES

- [A] M.F. ATIYAH, *K-Theory* (Benjamin, New York 1967).
- [BC] P. BAUM and A. CONNES, ‘The Chern character for discrete groups’, *A Fête of Topology*, Academic Press, New York, 1988, 163 – 232.
- [Black] B. BLACKADAR, *K-Theory for operator algebras* (Cambridge University Press, 2002.)
- [B] N. BOURBAKI, *Lie groups and Lie algebras* (chapters 4-6, Springer 2002; chapters 7-9, Springer 2008).
- [Sp] N. BOURBAKI, *Théories Spectrales*, Springer, 2007.
- [D] J. DIEUDONNÉ, *Éléments d’analyse*, volume 5, Bordas, Paris, 1975.
- [K] G. KASPAROV, *Equivariant KK-theory and the Novikov conjecture* *Invent. math.* 91,147-201 (1988).

SCHOOL OF MATHEMATICS, SOUTHAMPTON UNIVERSITY, SOUTHAMPTON SO17 1BJ,  
ENGLAND *and* SCHOOL OF MATHEMATICS, MANCHESTER UNIVERSITY, MANCHESTER  
M13 9PL, ENGLAND

*E-mail address:* g.a.niblo@soton.ac.uk, wright@soton.ac.uk, r.j.plymen@soton.ac.uk,  
plymen@manchester.ac.uk