

A short review on Landsberg spaces

Dodson, CTJ

2006

MIMS EPrint: **2006.82**

Manchester Institute for Mathematical Sciences
School of Mathematics

The University of Manchester

Reports available from: <http://eprints.maths.manchester.ac.uk/>

And by contacting: The MIMS Secretary
School of Mathematics
The University of Manchester
Manchester, M13 9PL, UK

ISSN 1749-9097

A short review on Landsberg spaces*

C.T.J. Dodson

School of Mathematics, University of Manchester
Manchester M60 1QD, UK

Abstract

This short review is concerned with real finite-dimensional Finsler manifolds (M, F) with Finsler structures $F : TM \rightarrow [0, \infty)$ that satisfy the Landsberg conditions. In particular this includes the case of Berwald manifolds since their Chern connections on π^*TM are fibre-independent. The aim is to provide an annotated collection of references to geometric results that seem important in the study of Landsberg spaces and to suggest some areas for further work in this context.

AMS Subject Classification (2001): 53B40, 53C60, 58B20

Key words: Finsler manifold, Cartan tensor, Ehresmann connection, Chern connection, curvature, exponential map, Landsberg space, Berwald space, Randers space, conformal equivalence, Killing field, parallel isometry, completeness

1 Introduction

Finsler manifolds (or spaces) can be thought of as generalizations of Riemannian manifolds; tangent spaces carry Minkowski norms instead of inner products and geometric objects on tangent vectors depend not only on the base but also on the fibre component. Chern and Shen [21] have provided an authoritative treatment of the subject, for which treatise the book by Bao, Chern and Shen [20] is a helpful introduction with a wealth of detail and we mainly follow the notational conventions of these authors and, similarly, we restrict our attention only to the real case. Finsler manifolds have intrinsic geometrical significance and also they have been used to model a variety of problems from dynamics, optics, ecology and relativity, cf. eg. Antonelli et al. [4], Bao et al. [20] and Asanov [7].

Consider $\phi : U \rightarrow \mathbb{R}^n : x \mapsto (x^i)$ as a local coordinate system on an open set U of a \mathbb{C}^∞ manifold M , with (∂_{x^i}) as the induced coordinate basis for the tangent space T_xM at a point $x \in M$. Let

$$F : TM \rightarrow [0, \infty) : (x, y) \mapsto F(x, y)$$

be \mathbb{C}^∞ on $TM \setminus \{0\}$, positively homogeneous of degree 1 in the fibre coordinate, and satisfying for each $x \in M$ the Minkowski norm condition

$$\lim_{s \rightarrow 0, t \rightarrow 0} \frac{F^2(x, y + su + tv)}{2} = g_y(u, v)$$

where g_y is an inner product on T_xM . In this case, using local coordinates for $(x, y) \in TM \setminus \{0\}$, the Hessian

$$(1.1) \quad [g_{ij}] = [\partial_{y^i} \partial_{y^j} (\frac{F^2(x, y^i \partial_{y^i})}{2})]$$

is positive definite, so as a matrix it has everywhere rank $n - 1$. Then we call F a Finsler structure on TM and, at each $x \in M$, $F(x, -)$ is a Minkowski norm on T_xM . Given a manifold M and a Finsler structure F on TM , the pair (M, F) is called a Finsler manifold. Sometimes, an n -dimensional Finsler manifold (M, F) is referred to as F^n with base space M . In the circumstance that each $F(x, -)$ is a

*Invited paper, Workshop on Finsler and semi-Riemannian geometry 24-26 May 2006, San Luis Potosí, México

Euclidean norm, subordinate to an inner product on $T_x M$, we reduce to the Riemannian case (M, g) with

$$F(x^i, y^j) = g_{ij} y^i y^j.$$

On a Finsler manifold (M, F) , the family of Minkowski norms $\{F(x, -) | x \in M\}$ yields a length function on oriented piecewise smooth curves in M . By taking infima over all such curves between a given pair of points in M we obtain a function d_F that is positive definite and satisfies the triangle inequality but it is not necessarily symmetric. The topology induced by d_F coincides with the manifold topology of M .

The matrix (1.1) gives the components of the fundamental tensor g of (M, F) ; explicitly, g is a Riemannian metric tensor on the pullback bundle $\pi^* TM$ (over $TM \setminus \{0\}$) in natural (pullback) coordinates denoted also by (x^i) . This Riemannian metric on $\pi^* TM$ determines formal Christoffel symbols, γ_{jk}^i via the usual formula.

The Cartan tensor A is a trilinear symmetric form on $\pi^* TM$ (over $TM \setminus \{0\}$) and has components

$$(1.2) \quad [A_{ijk}] = \frac{F}{2} [\partial_{y^k} g_{ij}] = \frac{F}{4} [\partial_{y^i} \partial_{y^j} \partial_{y^k} F^2].$$

For Riemannian manifolds, $A = 0$.

The Hilbert form ω on $\pi^* TM$ (over $TM \setminus \{0\}$) and its dual ℓ are given in natural coordinates by

$$(1.3) \quad \omega = \partial_{y^i} F dx^i = \omega_i dx^i$$

$$(1.4) \quad \ell = \frac{y^i}{F} \partial_{x^i} = \ell^i \partial_{x^i}$$

and they satisfy $\omega(\ell) = 1 = g(\ell, \ell)$.

The nonlinear Ehresmann connection on $\pi^* TM$ has components N_j^i given by

$$(1.5) \quad \begin{aligned} N_j^i &= \gamma_{jk}^i y^k - \frac{1}{F} g^{im} A_{mjk} \gamma_{rs}^k y^r y^s \\ &= \gamma_{jk}^i y^k - C_{jk}^i \gamma_{rs}^k y^r y^s. \end{aligned}$$

The Chern (linear torsion free) connection ∇ on $\pi^* TM$ has components Γ_{jk}^i given by

$$(1.6) \quad \begin{aligned} \Gamma_{jk}^i &= \gamma_{jk}^i - \frac{g^{im}}{F} (A_{mjs} N_k^s - A_{jks} N_m^s + A_{kms} N_j^s) \\ &= \frac{g^{is}}{2} (\delta_{x^k} g_{sj} - \delta_{x^s} g_{jk} + \delta_{x^j} g_{ks}) \end{aligned}$$

with connection forms ω_j^i satisfying

$$(1.7) \quad \omega_j^i = \Gamma_{jk}^i dx^k$$

$$(1.8) \quad d(dx^i) - dx^j \wedge \omega_j^i = -dx^j \wedge \omega_j^i = 0$$

$$(1.9) \quad dg_{ij} - g_{kj} \omega_i^k - g_{ik} \omega_j^k = \frac{2}{F} A_{ijs} \delta_{y^s}.$$

If the Γ_{jk}^i are functions of x only, then (M, F) is called a Berwald space (or manifold) and F is called a Berwald metric; in this case the Γ_{jk}^i actually arise from a family of associated Riemannian metrics.

The Finsler manifold is called a Landsberg space (or manifold) and F is called a Landsberg metric if

$$(1.10) \quad \ell^s \nabla_{\delta_x^s} (A_{ijk}) = 0$$

$$(1.11) \quad \text{equivalently } 0 = (\delta_{x^s} A_{ijk} - A_{mjk} \Gamma_{is}^m - A_{imk} \Gamma_{js}^m - A_{ijm} \Gamma_{ks}^m) \ell^s$$

$$(1.12) \quad \text{equivalently } \Gamma_{kj}^i = \frac{1}{2} \partial_{y^j} \partial_{y^k} (\Gamma_{kj}^i y^k y^j).$$

Particular Finsler metric functions include:

(i) $F = \alpha + \beta$, the Randers metric,

(ii) $F = \alpha^2/\beta$, the Kropina metric,

where $\alpha^2 = a_{ij}(x)y^i y^j$ is a Riemannian metric, and $\beta = b_i(x)y^i$ is a non-zero differential 1-form on M . A Randers Landsberg space of dimension two is a Berwald space; a Kropina space of dimension two with $b^2 = 0$ is a Landsberg space, then it is a Berwald space (where $b^2 = a_{rs}(x)b^r b^s$ and a_{rs} is the associated Riemannian metric tensor) [33].

A Finsler manifold is called locally projectively flat if, for all $(x, y) \in TM \setminus \{0\}$,

$$(1.13) \quad y^k \partial_{x^k} \partial_{y^i} F(x, y) = \partial_{x^i} F(x, y)$$

then about every point there are local charts in which the geodesic segments are mapped to lines in coordinate space. A Berwald space with scalar curvature is projectively flat and it is Riemannian if of nonzero constant curvature.

Atkin [11] proved a very nice theorem that extends to the Finsler case the Hopf-Rinow completeness theorem of Riemannian manifolds. He proved that a connected C^1 manifold M of finite dimension, possibly with boundary, admits a bounded complete Finsler structure if and only if it is compact. Furthermore, if M is a C^1 Banach manifold admitting a complete Finsler structure, and N is a connected noncompact C^1 Banach manifold admitting a bounded complete Finsler structure, then $M \times N$ admits a bounded complete Finsler structure. It follows that any C^1 Banach manifold satisfying a certain stability condition admits a complete bounded Finsler structure. See also Bao et al. [20] Chapter VI for a detailed study of Finsler forward and backward completeness and reference there to the thesis work of Dazord in 1969.

Rademacher [78, 79] extended some classical comparison theorems in Riemannian geometry to the Finsler case. He introduced the reversibility

$$\lambda := \max\{F(x, -y) \mid F(x, y) = 1\},$$

and showed that if M is a simply-connected, compact Finsler manifold of dimension $n \geq 3$ with reversibility λ and the flag curvature satisfies $(1 - \frac{1}{1+\lambda})^2 < K \leq 1$, then the length of a geodesic loop is at least $\pi(1 + \frac{1}{\lambda})$. He then proved that a simply-connected and compact Finsler manifold of dimension $n \geq 3$ with reversibility λ and flag curvature $(1 - \frac{1}{1+\lambda})^2 < K \leq 1$ is homotopy equivalent to the n -sphere.

1.1 Examples

We collect some examples but see the following texts for details of many more: Bao, Chern and Shen [20], Matsumoto [63], Shen [84], Chern and Shen [21], Antonelli et al. [4, 6] and Asanov [7].

1. A Finsler manifold (M, F) is called a Finsler torus if the space is homeomorphic to a torus, and a two-dimensional Finsler torus is called flat if it is equipped with a Finsler structure that is obtained by passing to a quotient from a Finsler structure of \mathbb{R}^2 invariant under translations.
2. If a two-dimensional Finsler torus is a Landsberg space and has no conjugate point then the space is isometric to a flat Finsler torus [27].
3. 3-dimensional Landsberg spaces of constant curvature are either Riemannian spaces or spaces of vanishing curvature [67].
4. In a Finsler space of two dimensions the holonomy group is in general an infinite continuous group. This group has one, two or three parameters but this reduces to one parameter for a Landsberg space [67].
5. All non-Riemannian Berwald spaces can be constructed from cartesian products among 54 basic non-Riemannian Berwald spaces devised by Szabo [88].
6. A Finsler space of dimension 2 is a generalized Berwald space if and only if the first derivative of the main scalar by the Landsberg angle gives a differential equation of the form $y = f'(y)$.

7. Lee and Park [54] showed that a Finsler space (M, F) with (α, β) -metric

$$(1.14) \quad L(\alpha, \beta) = \frac{\beta^2}{\beta - \alpha} = \beta \sum_{r=0}^{\infty} \left(\frac{\alpha}{\beta} \right)^r, \quad \alpha = (a_{ij}(x)y^i y^j)^{1/2}, \beta = b_i(x)y^i$$

is projectively flat if and only if it is a Berwald space and the associated Riemannian space is projectively flat.

8. Bácsó and Matsumoto [16] showed that a Finsler n -manifold (M, F) is a projectively flat Berwald space if and only if it belongs to one of the following classes: (1) $n \geq 3$: (a) locally Minkowski spaces, (b) Riemannian spaces of constant curvature; or (2) $n = 2$: (a) locally Minkowski spaces, (b) Riemannian spaces of constant curvature, (c) spaces with $L = \beta^2/\gamma$ and the signature $\epsilon = +1$.

2 Landsberg Geometry

Every Berwald space is a Landsberg space; however, it is not known if the converse is true [83]. Vattamány [95] proved the general case that every Landsberg space with vanishing Douglas tensor is a Berwald space. Hence, a Finsler manifold (M, F) is a Berwald space if there exists a symmetric linear connection ∇ having parallel translation which preserves the metric; if such a connection ∇ is flat, then (M, F) is said to be locally Minkowski [72]. Muzsnay [68] has shed more light on the equivalence problem by establishing conditions under which a second-order homogeneous ordinary differential equation (spray) be Finsler metrizable—ie the geodesic equation of a Finsler space, in terms of the holonomy algebra generated by horizontal vector-fields. Muzsnay obtained similar results for the Landsberg case, in particular, he proved that a quadratic second order differential equation is Landsberg metrizable if and only if it is Finsler metrizable. Most recently, Asanov [10] suggests some new methodology by his introduction of the concept of Finsleroid Finsler spaces.

Mo [66] gave geometric and algebraic characterizations for Finsler spaces with zero Riemann curvature. In particular, such spaces are characterized by the fact that the horizontal distribution of the projective sphere bundle has a flat foliation.

Wang [98] solved the problem of determining all Finsler spaces of dimensions $n > 2$ which admit a $\frac{1}{2}n(n+1)$ parameter group of motions: they are the Riemann spaces of constant curvature. The method of proof uses results on Lie groups and linear groups rather than the classical method of studying integrability conditions. Singh et al. [86] obtained conditions for a vector field to be a Killing field in a Randers space. Killing field properties in Finsler manifolds have been studied by Yawata [101] who gave the form of Killing equations with respect to the Cartan and Berwald connections. Lovas [55] studied Killing fields whose integral curves are geodesics of an associated Finsler manifold, including the case of a Randers metric.

Antonelli and Lackey [5] and Antonelli and Zastawniak [6] provided detailed treatments of the Finsler analogue of Laplace operators and Hodge decomposition, and the classification theorem for 2-dimensional Berwald spaces which are not locally Minkowski. Centore [24] provided a very elegant characterisation of Riemannian spaces as a subset of Finsler spaces and Berwald spaces as a subset of Finsler spaces, solely in terms of the two naturally associated volume forms: Riemannian and Busemann [23]. For further discussion of the role of the Busemann volume form, in the context of developing a Laplacian for Finsler spaces, see Centore [25].

For a positive definite Finsler manifold the associated Levi-Civita connection coincides with the canonical connection if the Finsler space reduces to a Berwald space [88] and if a linear connection on the base manifold is compatible with the horizontal distribution of a Finsler space, then it is compatible with respect to the associated Riemannian metric [96].

A Berwald space (M, F) is locally, respectively globally, symmetric if the Chern connection is locally, respectively globally, symmetric. Deng [28] proved that every locally symmetric Berwald symmetric space is locally isometric to a globally symmetric Berwald space; this extends to locally geodesic Berwald spaces.

Yang [99] reported a necessary and sufficient condition for a Finsler space with (α, β) -metric to be a Berwald space, and studied the conformal changes between two (α, β) -metric Finsler spaces.

Finsler manifolds admit exponential maps and normal neighbourhoods. In particular, Kristaly et al. [50] proved that in a Berwald space (M, F) of non-positive curvature, every point $x \in M$ admits a neighborhood, such that two geodesics $\gamma_1, \gamma_2: [0, 1] \rightarrow M$, emanating from x ($\gamma_1(0) = \gamma_2(0) = x$), satisfy the inequality

$$(2.15) \quad 2d_F(\gamma_1(\tfrac{1}{2}), \gamma_2(\tfrac{1}{2})) \leq d_F(\gamma_1(1), \gamma_2(1)).$$

It follows that the length of a median of a geodesic triangle in M is smaller than or equal to the length of the corresponding side.

Lee and Park [54] studied Finsler spaces (M, F) with (α, β) -metric

$$(2.16) \quad L(\alpha, \beta) = \frac{\beta^2}{\beta - \alpha} = \beta \sum_{r=0}^{\infty} \left(\frac{\alpha}{\beta} \right)^r, \quad \alpha = (a_{ij}(x)y^i y^j)^{1/2}, \beta = b_i(x)y^i$$

They proved that (M, F) is a Berwald space if and only if $b_{j;i} = 0$; then the Berwald connection is Riemannian. (M, F) is projectively flat if and only if it is a Berwald space and the associated Riemannian space is projectively flat. Lee [52] studied a Finsler space with the special (α, β) metric

$$L(\alpha, \beta) = c_1\alpha + c_2\beta + \alpha^2/\beta$$

satisfying some conditions, finding a condition under which this special Finsler space is a Berwald space. If a two-dimensional Finsler space with this metric $L(\alpha, \beta)$ is a Landsberg space, then it is a Berwald space.

Park and Lee [75] considered a Finsler space $F^n = (M, L)$ with a generalized Randers metric

$$L^2(\alpha, \beta) = c_1\alpha^2 + c_2\alpha\beta + c_3\beta^2$$

(where $\alpha^2 = a_{ij}(x)y^i y^j$ may be any quadratic form, $\beta = b_i(x)y^i$ and c_1, c_2, c_3 are nonzero constants). This type of Finsler space is a Landsberg space if and only if b_i is a Killing vector field with constant length, and such a Landsberg space is a Berwald space.

Tamásy [90] studied the class of Finsler spaces that admit metrical linear connections; these are exactly the affine deformations of the associated locally Minkowski spaces. Moreover, a Finsler space admits a metrical linear connection in the tangent bundle TM if and only if it is an affine deformation of a Berwald space with vanishing h -curvature tensor K of its Rund connection.

Pandey and Tiwari [71] studied the Landsberg and semi- C -reducible Landsberg cases and explicitly derived expressions of the h -connection vectors and of the h -covariant derivative of the h - $h\nu$ torsion tensor. They thereby obtained necessary and sufficient conditions for a 4-dimensional semi- C -reducible Landsberg space to be a Berwald space, in terms of relations satisfied by the nontrivial main scalars.

Dragomir [29] reported the following results on harmonic maps: Let $(M^n(c), E)$ be a Finsler space of scalar curvature $c \neq 0$ and vanishing mixed torsion vector $P_j = \dot{\partial}_i N_j^i - F_{ji}^i$. All h -harmonic functions $f(x, y)$ on $T(M^n(c) \setminus \{0\})$ which are positive homogeneous of degree r in the y^i 's and whose h -gradient has compact support are given by $f = aE^{r/2}$, $a \in \mathbb{R}$. The image of a totally geodesic immersion of a Finsler space in a Landsberg space M^{n+p} is not contained in any h -convex supporting set of M^{n+p} .

2.1 Conformal properties

Two Finsler manifolds (M, F) and (M, \tilde{F}) are conformally equivalent if there exists a positive smooth function $\varphi: TM \rightarrow \mathbb{R}$, called the scale function, such that $\tilde{g} = \varphi g$. Aikou [2] used the Weyl structure of a conformal class determined by the associated Riemannian metric to characterise when a Finsler manifold is conformal to a Berwald manifold; Vincze [97] showed that in this case the exterior derivative of the scale function is closed and exact, cf also Tamásy [90]. Aikou [3] treats the complex case.

By Hashiguchi's theorem [31], cf also [32], a Landsberg space remains a Landsberg space under any conformal change of metric if and only if its T -tensor field vanishes identically. Matsumoto [62] gave a necessary and sufficient condition for the Berwald spaces property to persist under conformal change of metrics; in the case of a 2-dimensional Berwald space this condition is if and only if it has constant main scalar. Ikeda [39] provided criteria for conformal flatness of Finsler spaces, developing

further the method and theorem of Kikuchi [45]. Yang [99] studied the conformal changes between two (α, β) -metric Finsler spaces.

Izumi [42] introduced the notion of a curve in a Finsler space F^n as a geodesic circle if its first and second curvatures are constant and zero, respectively. This definition depends on the choice of the connection so various cases were considered. A conformal transformation $(*) F \rightarrow F' = \exp[\sigma(x)]F$ is concircular if it preserves geodesic circles. Necessary and sufficient conditions for concircularity were derived for the various connections in terms of systems of partial differential equations to be satisfied by the conformal function $\sigma(x)$. Subsequently, Izumi [43] called the conformal transformation $(*)$ h -conformal when, for the Finsler objects

$$2g_{hj} = \partial^2 F / \partial y^h \partial y^j, \quad \text{and} \quad 2C_{hjk} = \partial g_{hj} / \partial y^k,$$

his h -condition is satisfied, namely:

$$(n-1)C_{ij}^h \sigma_h = C^h \sigma_h h_{ij}, \quad \text{where} \quad \sigma_h = \partial \sigma / \partial x^h, \quad C_h = C_{hj}^j, \quad h_{ij} = g_{ij} - y_i y_j / F^2.$$

He gave a geometric interpretation of this condition and some h -conformal invariants are displayed, with necessary and sufficient conditions under which F^n is h -conformally flat, that is, h -conformal to a Minkowskian space. Singh and Gupta [87] studied conformal and h -conformal transformation in special Finsler spaces, including the Landsberg case.

Prasad and Dwivedi [77] studied conformal change in three-dimensional Finsler spaces, providing a three-dimensional Finsler space which is conformal to a Berwald space or Landsberg space. Prasad and Dwivedi [76] studied conformal changes: $L(x, y) \rightarrow \bar{L}(x, y) = e^{\alpha(x)}L(x, y)$ and the associated changes of Cartan connection and Berwald connection; they investigated the conditions under which m -th root metric Berwald spaces, \mathbb{S}^3 -like spaces and Landsberg spaces are preserved by a conformal change of the metric.

2.2 Necessary and sufficient conditions for (M, F) to be Landsberg

We collect some characterising conditions from the literature.

1. Along every curve c the parallel translation $P_c: (T_{c(a)}M, g_{c(a)}) \rightarrow (T_{c(b)}M, g_{c(b)})$ is an isometry between the Riemannian spaces [36].
2. The vertical foliation in (TM, G) is totally geodesic, with [1]

$$G = \frac{1}{2} \frac{\partial^2 F^2}{\partial y^i \partial y^j} [dx^i \otimes dx^j + \theta^i \otimes \theta^j]$$

$$\theta^i := dy^i + N_m^i dx^m.$$

3. Each fibre is a totally geodesic submanifold in the total space TM with a Sasaki-type metric [1].
4. The infinitesimal h -mapping

$$(x_0^i, l_0^i) \mapsto (x^i = x_0^i + dx_0^i, l^i = l_0^i - l_0^j \Gamma_{jk}^i(x_0, l_0) dx_0^k)$$

is affine or an isometry [100]. Shen proved that a Finsler space is Berwald if and only if along every curve parallel translation is an isometry between the associated Minkowski spaces.

5. In a Finsler space with a generalized Randers metric

$$L^2(\alpha, \beta) = c_1 \alpha^2 + c_2 \alpha \beta + c_3 \beta^2$$

(where $\alpha^2 = a_{ij}(x)y^i y^j$ may be any quadratic form, $\beta = b_i(x)y^i$ and c_1, c_2, c_3 are nonzero constants) b_i is a Killing vector field with constant length.

6. For dimension 2, every Landsberg space is a Berwald space.

2.3 Landsberg Problems

We mention some areas that seem still to offer challenges in the study of Landsberg spaces.

Local projectivity Characterization of locally projectively Berwald spaces; see Bácsó [12] for a summary.

Berwald equivalence Every Berwald space is a Landsberg space; however, it is not known if the converse is true; see eg Shen [83] and the new work of Muzsnay [68], also Asanov [10] seems to provide a major step forward.

Conformal properties Characterization of conformal classes, eg conformal flatness. Prasad and Dwivedi [76] investigated the conditions under which special Finsler spaces with m -th root metric (i.e., Berwald spaces, S^3 -like spaces and Landsberg spaces) are preserved by a conformal change of the metric.

Jacobi fields Characterization of Jacobi vector fields. Hassan [34, 35] studied sprays and Jacobi fields in Finsler spaces; Tamin [91] discussed some Randers manifolds and Crampin [26] provided a covariant form of the Lagrangian second variation formula and showed that each of the four standard connections encountered in Finsler geometry produces the same result.

Harmonic maps See Dragomir [29] for some work in this area.

Local flow diffeomorphisms isometric, harmonic.

Warped products Udriste [94] and Kozma et al. [48] have begun the extension of the Riemannian methods to Finsler spaces. See also Asanov [8, 9] for study of the Finsler case $M \times \mathbb{R}$.

Acknowledgement

The author is grateful to PROMEP for support to attend this Workshop and to the Universidad Autónoma de San Luis Potosí for hospitality.

References

- [1] Aikou, Tadashi. Some remarks on the geometry of tangent bundle of Finsler manifolds. *Tensor (N.S.)* 52 (1993), no. 3, 234–242.
- [2] Aikou, Tadashi. Locally conformal Berwald spaces and Weyl structures. *Publ. Math. Debrecen* 49, 1-2 (1996) 113-126.
- [3] Aikou, Tadashi. Some remarks on locally conformal complex Berwald spaces. Finsler geometry (Seattle, WA, 1995), 109-120, *Contemp. Math.*, 196, Amer. Math. Soc., Providence, RI, 1996.
- [4] Antonelli, P.L.; Ingarden, R.S. and Matsumoto, M. **The theory of sprays and Finsler spaces with applications in Physics and Biology**, Kluwer Academic Publishers (1993).
- [5] Antonelli, P.L.; Lackey, B.C., Eds, **The theory of Finslerian Laplacians and applications**. Mathematics and its Applications, 459. Kluwer Academic Publishers, Dordrecht, 1998.
- [6] Antonelli, P.L.; Zastawniak, T. J. **Fundamentals of Finslerian diffusion with applications. Handbook of Finsler geometry**. Vol. 1, 2, 177–355, Kluwer Acad. Publ., Dordrecht, 2003.
- [7] Asanov, G.S. **Finsler Geometry, Relativity and Gauge Theories** D. Reidel, Dordrecht 1985.
- [8] Asanov, G. S. Finsler cases of GF-space. *Aequationes Math.* 49, 3 (1995) 234-251.
- [9] Asanov, G. S. Finslerian metric functions over the product $R \times M$ and their potential applications. *Rep. Math. Phys.* 41, 1 (1998) 117-132.

- [10] Asanov, G. S. Finsleroid–Finsler Space with Berwald and Landsberg Conditions. Preprint, 2006 <http://arxiv.org/abs/math.DG/0603472>
- [11] Atkin, C. J. Bounded complete Finsler structures. I. *Studia Math.* 62, 3 (1978) 219-228.
- [12] Bácsó, S. On a problem of M. Matsumoto and Z. Shen. **Finsler and Lagrange geometries** (Iași, 2001), 55-61, Kluwer Acad. Publ., Dordrecht, 2003.
- [13] Bácsó, S.; Ilosvay, F.; Kis, B. Landsberg spaces with common geodesics. *Publ. Math. Debrecen* 42 (1993), no. 1-2, 139-144.
- [14] Bácsó, S.; Matsumoto, M. Reduction theorems of certain Landsberg spaces to Berwald spaces. *Publ. Math. Debrecen* 48 (1996), no. 3-4, 357-366.
- [15] Bácsó, S.; Hashiguchi, M.; Matsumoto, M. Generalized Berwald spaces and Wagner spaces. *An. Științ. Univ. Al. I. Cuza Iași. Mat. (N.S.)* 43,2 (1997) 307-321.
- [16] Bácsó, S.; Matsumoto, M. On Finsler spaces of Douglas type—a generalization of the notion of Berwald space. *Publ. Math. Debrecen* 51,3-4 (1997) 385-406.
- [17] Bácsó, S.; Yoshikawa, R. Weakly-Berwald spaces. *Publ. Math. Debrecen* 61, 1-2 (2002) 219-231.
- [18] Bao, David; Chern, S. S.; Shen, Z. On the Gauss-Bonnet integrand for 4-dimensional Landsberg spaces. Finsler geometry (Seattle, WA, 1995), 15-26, *Contemp. Math.*, 196, Amer. Math. Soc., Providence, RI, 1996.
- [19] Bao, David; Shen, Zhong Min On the volume of unit tangent spheres in a Finsler manifold. *Results Math.* 26,1-2 (1994) 1-17.
- [20] Bao, David; Chern, Shiing-Shen; Shen, Zhongmin. **An Introduction to Riemann-Finsler Geometry**, Graduate Texts in Mathematics 200, Springer-Verlag, New York 2000.
- [21] Chern, Shiing-Shen; Shen, Zhongmin. **Riemann-Finsler geometry**. Nankai Tracts in Mathematics, 6. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2005.
- [22] Bejancu, Aurel; Farran, Hani Reda Generalized Landsberg manifolds of scalar curvature. *Bull. Korean Math. Soc.* 37, 3 (2000) 543-550.
- [23] Busemann; H. Intrinsic area. *Annals of Math.* 48, (1947)234-267.
- [24] Centore, P. Volume forms in Finsler spaces. *Houston J. Math.* 25, 4 (1999) 625-640.
- [25] Centore, P. Finsler Laplacians and minimal-energy maps. *Internat. J. Math.* 11, 1 (2000) 1-13.
- [26] Crampin, M. The second variation formula in Lagrange and Finsler geometry. *Houston J. Math.* 26, 2 (2000) 255-275.
- [27] Dazord, Pierre. Tores finslériens sans points conjugués. *Bull. Soc. Math. France* 99 (1971) 171-192; *erratum, ibid.* 99 (1971) 397.
- [28] Deng, S; Hou, Z. On locally and globally symmetric Berwald spaces. *J. Phys. A* 38, 8 (2005) 1691-1697.
- [29] Dragomir, S. Harmonic functions on Finsler spaces. *Istanbul Ünv. Fen Fak. Mat. Derg.* 48 (1987/89), 67-76.
- [30] Dragomir, S.; Capursi, M. On the topology of Landsberg spaces. *Riv. Mat. Univ. Parma* (4) 13 (1987), 305-313.
- [31] Hashiguchi, Masao. On conformal transformations of Finsler metrics. *J. Math. Kyoto Univ.* 16, 1 (1976) 25-50.
- [32] Hashiguchi, Masao; Ichijyō, Yoshihiro. On conformal transformations of Wagner spaces. *Rep. Fac. Sci. Kagoshima Univ.* 10, (1977), 19-25.

- [33] Hashiguchi, Masao; Hōjō, Shun-ichi; Matsumoto, Makoto, Landsberg spaces of dimension two with (α, β) -metric. *Tensor (N.S.)* 57,2 (1996) 145-153.
- [34] Hassan, B. T. M. **The theory of geodesics in Finsler spaces**, PhD Thesis, Southampton University, 1967.
- [35] Hassan, B. T. M. Sprays and Jacobi fields in Finsler geometry. *An. Univ. Timișoara Ser. Științ. Mat.* 19, 2 (1981) 129-139.
- [36] Ichijyo, Y. On special Finsler connections with the vanishing hv -curvature tensor. *Tensor (N.S.)* 32,2 (1978) 149-155.
- [37] Ikeda, Fumio Some remarks on Landsberg spaces. *TRU Math.* 22,2 (1986) 73-77.
- [38] Ikeda, Fumio. On some properties of three-dimensional Finsler spaces. *Tensor (N.S.)* 55 (1994), no. 1, 66-73.
- [39] Ikeda, Fumio. Criteria for conformal flatness of Finsler spaces. *Balkan J. Geom. Appl.* 2, 2 (1997) 63-68.
- [40] Ikeda, Fumio Landsberg spaces satisfying the T -condition. *Balkan J. Geom. Appl.* 3,1 (1998) 23-28.
- [41] Ikeda, Fumio On S_4 -like Finsler spaces with T -condition. The 5th International Conference of Tensor Society on Differential Geometry and its Applications and Information Sciences their Mathematical Foundations (Calcutta, 1998). *Tensor (N.S.)* 61, 2 (1999) 171-175.
- [42] Izumi, Hideo. Conformal transformations of Finsler spaces. I. Conircular transformations of a curve with Finsler metric. *Tensor (N.S.)* 31, 1 (1977) 33-41.
- [43] Izumi, Hideo. Conformal transformations of Finsler spaces. II. An h -conformally flat Finsler space. *Tensor (N.S.)* 34, 3 (1980) 337-359.
- [44] Jun, Dong-Kum; Lee, Il-Yong On two-dimensional Landsberg spaces of the Finsler space with $L^2(\alpha, \beta) = 2\alpha\beta$ metric. *Far East J. Math. Sci.* 6,1 (1998) 75-88.
- [45] Kikuchi, Shigetaka. On the condition that a Finsler space be conformally flat. *Tensor (N.S.)* 55, 1 (1994) 97-100.
- [46] Kozma, L. On Landsberg spaces and holonomy of Finsler manifolds. Finsler geometry (Seattle, WA, 1995), 177-186, *Contemp. Math.* 196, American Mathematical Society, Providence, RI, 1996.
- [47] Kozma, L. On holonomy groups of Landsberg manifolds. *Tensor (N.S.)* 62, 1 (2000) 87-90.
- [48] Kozma, L.; Peter, R. and Varga, C. Warped product of Finsler manifolds. *Ann. Univ. Sci. Budapest. Eötvös Sect. Math.* 44 (2001) 157-170.
- [49] Kovács, Z. On the Chern-Weil homomorphism in Finsler spaces. *Acta Math. Acad. Paedagog. Nyházi. (N.S.)* 17, 2 (2001) 131-135.
- [50] Kristály, A; Varga, Cs; Kozma, L. The dispersing of geodesics in Berwald spaces of non-positive flag curvature. *Houston J. Math.* 30, 2 (2004) 413-420.
- [51] Lee, Yong-Duk Conformal transformations of difference tensors of Finsler space with an (α, β) -metric. *Commun. Korean Math. Soc.* 12,4 (1997) 975-984.
- [52] Lee, Il-Yong On two-dimensional Landsberg space with a special (α, β) -metric. *J. Korea Soc. Math. Educ. Ser. B Pure Appl. Math.* 10, 4 (2003) 279-288.
- [53] Lee, Il-Yong; Lee, Myung-Han. Landsberg space and differential equation of geodesics of dimension two on Matsumoto metric. *Far East J. Math. Sci.* 12, 2 (2004) 259-275.
- [54] Lee, Il-Yong; Park, Hong-Suh. Finsler spaces with infinite series (α, β) -metric. *J. Korean Math. Soc.* 41, 3 (2004) 567-589.

- [55] Lovas, Rezs L. On the Killing vector fields of generalized metrics. *SUT J. Math.* 40, 2 (2004) 133-156.
- [56] Matsumoto, Makoto Remarks on Berwald and Landsberg spaces. Finsler geometry (Seattle, WA, 1995), 79-82, *Contemp. Math.*, 196, American Mathematical Society, Providence, RI, 1996.
- [57] Matsumoto, Makoto Theory of Finsler spaces with m -th root metric. II. *Publ. Math. Debrecen* 49,1-2 (1996) 135-155.
- [58] Matsumoto, Makoto On the stretch curvature of a Finsler space and certain open problems. Special volume to felicitate Prof. Dr. R. S. Mishra on the occasion of his 80th birthday. *J. Nat. Acad. Math. India* 11 (1997), 22-32.
- [59] Matsumoto, Makoto Reduction theorems of Landsberg spaces with (α, β) -metric. *Tensor (N.S.)* 58,2 (1997) 160-166.
- [60] Matsumoto, Makoto Projective Randers change of P -reducible Finsler spaces. 4th International Conference on Differential Geometry and its Applications (Tsukuba, 1996). *Tensor (N.S.)* 59 (1998), 6-11.
- [61] Matsumoto, Makoto A theory of three-dimensional Finsler spaces in terms of scalars and its applications. *An. Ştiinţ. Univ. Al. I. Cuza Iaşi. Mat. (N.S.)* 45, 1 (1999) 115-140.
- [62] Matsumoto, Makoto Conformally closed Finsler spaces. *Balkan J. Geom. Appl.* 4,1 (1999) 117-128.
- [63] Matsumoto, M. Finsler geometry in the 20th-century. **Handbook of Finsler geometry**. Vol. 1, 2, 557-966, Kluwer Acad. Publ., Dordrecht, 2003.
- [64] Mishra, C. K.; Yadav, D. D. S. On special Finsler spaces with Landsberg space. *Acta Cienc. Indica Math.* 30, 3 (2004) 605-608.
- [65] Mo, X. The flag curvature tensor on a closed Finsler space. *Results Math.* 36, 1-2 (1999) 149-159.
- [66] Mo, X. Structure of Finsler space for which the R part of the curvature is zero. *J. Fudan Univ. Nat. Sci.* 39, 5,5 (2000) 518-524.
- [67] Mor, Arthur. Über die Torsions- und Krümmungsinvarianten der dreidimensionalen Finslerschen Räume. *Math. Nachr.* 16 (1957), 85-99.
- [68] Muzsnay, Z. The Euler-Lagrange PDE and Finsler metrizable. *Houston J. Math.* 32, 1 (2006) 79-98.
- [69] Pande, H. D.; Tripathi, P. N.; Prasad, B. N. On a special form of the hv -curvature tensor of Berwald's connection $B\Gamma$ of Finsler space. *Indian J. Pure Appl. Math.* 25,12 (1994) 1275-1280.
- [70] Pandey, T. N.; Diwedi, D. K. Normalized semi-parallel Ch -vector field in special Finsler spaces. *Indian J. Pure Appl. Math.* 30,3 (1999) 307-315.
- [71] Pandey, T. N.; Tiwari, Banktेशwar, On four dimensional semi- C -reducible Landsberg space. *Bull. Calcutta Math. Soc.* 94, 6 (2002) 421-430.
- [72] Park, Hong-Suh(; Lee, Il-Young. A Finsler space with a special metric function. *Commun. Korean Math. Soc.* 11, 2 (1996) 415-421.
- [73] Park, Hong-Suh; Lee, Il-Yong On Landsberg spaces of dimension two with a special (α, β) -metric. *Mem. Sect. Ştiinţ. Acad. Romnă Ser. IV* 21 (1998), 27-36.
- [74] Park, Hong S.; Lee, Il-Yong Landsberg spaces of dimension two with some (α, β) -metrics. *Panamer. Math. J.* 9,3 (1999) 41-56.
- [75] Park, Hong-Suh; Lee, Il-Yong On the Landsberg spaces of dimension two with a special (α, β) -metric. *J. Korean Math. Soc.* 37, 1 (2000) 73-84.

- [76] Prasad, B. N.; Dwivedi, Ashwini Kumar, On conformal transformation of Finsler spaces with m th root metric. *Indian J. Pure Appl. Math.* 33, 6 (2002) 789-796.
- [77] Prasad, B. N.; Diwedi, D. K. Conformal change of three-dimensional Finsler space. The 5th International Conference of Tensor Society on Differential Geometry and its Applications and Information Sciences and their Mathematical Foundations (Calcutta, 1998). *Tensor (N.S.)* 61, 2 (1999) 147-157.
- [78] Rademacher, Hans-Bert. A sphere theorem for non-reversible Finsler metrics. *Math. Ann.* 328, 3 (2004) 373-387.
- [79] Rademacher, Hans-Bert. Nonreversible Finsler metrics of positive flag curvature. **A sampler of Riemann-Finsler geometry**, 261–302, *Math. Sci. Res. Inst. Publ.*, 50, Cambridge Univ. Press, Cambridge, 2004.
- [80] Sakaguchi, Toshio On generalized metric spaces with $C_{ijk} = F^{-1}C_{ik}l_j + C^{-2}C_iC_jC_k$. *Tensor (N.S.)* 58,2 (1997) 167-170.
- [81] Shen, Zhongmin Some formulas of Gauss-Bonnet-Chern type in Riemann-Finsler geometry. *J. Reine Angew. Math.* 475 (1996), 149-165.
- [82] Shen, Zhongmin. Geometric meanings of curvatures in Finsler geometry. In Proceedings of the 20th Winter School "Geometry and Physics" (Srn, 2000). *Rend. Circ. Mat. Palermo (2) Suppl.* 66 (2001), 165-178.
- [83] Shen, Zhongmin. On R -quadratic Finsler spaces. *Publ. Math. Debrecen* 58, 1-2 (2001) 263-274.
- [84] Shen, Zhongmin. **Lectures on Finsler geometry**. World Scientific Publishing Co., Singapore, 2001.
- [85] Shimada, Hideo; Sabău, Vasile Sorin. Finsler geometry. Finslerian geometries (Edmonton, AB, 1998), 15-24, *Fund. Theories Phys.* 109, Kluwer Acad. Publ., Dordrecht, 2000.
- [86] Singh, U. P.; John, V. N.; Prasad, B. N. Finsler spaces preserving Killing vector fields. *J. Math. Phys. Sci.* 13, 3 (1979) 265-271.
- [87] Singh, U. P.; Gupta, B. N. Conformal and h -conformal transformation in special Finsler spaces. *Nepali Math. Sci. Rep.* 11, 1 (1986) 25-35.
- [88] Szabó, Z. I. Positive definite Berwald spaces. Structure theorems on Berwald spaces. *Tensor (N.S.)* 35, 1 (1981) 25-39.
- [89] Szabó, Z. I. Berwald metrics constructed by Chevalley's polynomials. Preprint: arXiv.org math.DG/0601522 (2006).
- [90] Tamássy, L. Finsler spaces with metrical linear connections in TM and conformal mappings. *Tensor (N.S.)* 64, 1 (2003) 34-40.
- [91] Tamim, Aly A. On Jacobi fields in Finsler geometry. *Proc. Math. Phys. Soc. Egypt* 76, (2001) 71-88.
- [92] Tamim, Aly A. Hereditary properties of special Finsler manifolds. *Publ. Math. Debrecen* 62, 1-2 (2003) 141-163.
- [93] Tamin, Aly A.; Youssef, Nabil L. Two nonrelated Finsler structures on a manifold. *Rev. Roumaine Math. Pures Appl.* 45, 4 (2000) 713-722.
- [94] Udriste, Constantin. Completeness of Finsler manifolds. *Publ. Math. Debrecen* 42, 1-2 (1993) 45-50.
- [95] Vattamány, Sz. Projection onto the indicatrix bundle of a Finsler manifold. *Publ. Math. Debrecen* 58, 1-2 (2001) 193-221.

- [96] Vincze, Cs. A new proof of Szabó's theorem on the Riemann-metrizability of Berwald manifolds. *Acta Math. Acad. Paedagog. Nyhzi. (N.S.)* 21, 2 (2005) 199-204.
- [97] Vincze, Cs. On a scale function for testing the conformality of a Finsler manifold to a Berwald manifold. *J. Geom. Phys.* 54, 4 (2005) 454-475.
- [98] Wang, Hsien-Chung. On Finsler spaces with completely integrable equations of Killing. *J. London Math. Soc.* 22, (1947). 5-9.
- [99] Yang, Guo Jun. Applications of a connection in Finsler spaces with (α, β) -metrics. *Xinan Shifan Daxue Xuebao Ziran Kexue Ban* 25, 3 (2000) 216-220.
- [100] Yasuda, Hiroshi. On Landsberg spaces. *Tensor (N.S.)* 34, 1 (1980) 77-84.
- [101] Yawata, Makoto. Killing equations in tangent bundle. Lagrange and Finsler geometry, 189-194, *Fund. Theories Phys.*, 76, Kluwer Acad. Publ., Dordrecht, 1996.