

*The Power of Bidiagonal Matrices*

Higham, Nicholas J.

2023

MIMS EPrint: **2023.3**

Manchester Institute for Mathematical Sciences  
School of Mathematics

The University of Manchester

Reports available from: <http://eprints.maths.manchester.ac.uk/>

And by contacting: The MIMS Secretary  
School of Mathematics  
The University of Manchester  
Manchester, M13 9PL, UK

ISSN 1749-9097

# THE POWER OF BIDIAGONAL MATRICES\*

NICHOLAS J. HIGHAM<sup>†</sup>

**Abstract.** Bidiagonal matrices are widespread in numerical linear algebra, not least because of their use in the standard algorithm for computing the singular value decomposition and their appearance as LU factors of tridiagonal matrices. We show that bidiagonal matrices have a number of interesting properties that make them powerful tools in a variety of problems, especially when they are multiplied together. We show that the inverse of a product of bidiagonal matrices is insensitive to small componentwise relative perturbations in the factors if the factors or their inverses are nonnegative. We derive componentwise rounding error bounds for the solution of a linear system  $Ax = b$ , where  $A$  or  $A^{-1}$  is a product  $B_1 B_2 \dots B_k$  of bidiagonal matrices, showing that strong results are obtained when the  $B_i$  are nonnegative or have a checkerboard sign pattern. We show that given the factorization of an  $n \times n$  totally nonnegative matrix  $A$  into the product of bidiagonal matrices,  $\|A^{-1}\|_\infty$  can be computed in  $O(n^2)$  flops and that in floating-point arithmetic the computed result has small relative error, no matter how large  $\|A^{-1}\|_\infty$  is. We also show how factorizations involving bidiagonal matrices of some special matrices, such as the Frank matrix and the Kac–Murdock–Szegő matrix, yield simple proofs of the total nonnegativity and other properties of these matrices.

**Key words.** Bidiagonal matrix, totally nonnegative matrix, condition number, matrix function, Vandermonde system, Toeplitz matrix, the Frank matrix, the Pascal matrix, the Kac–Murdock–Szegő Matrix.

**AMS subject classifications.** 15A06, 15A12, 15A23, 65F35

## 1. Introduction. Bidiagonal matrices

$$B = \begin{bmatrix} b_{11} & b_{12} & & & \\ & b_{22} & \ddots & & \\ & & \ddots & b_{n-1,n} & \\ & & & b_{nn} & \\ & & & & \end{bmatrix} \in \mathbb{C}^{n \times n}$$

have  $2n - 1$  parameters, appearing on two diagonals. Despite their simplicity, bidiagonal matrices are powerful tools in a variety of problems, especially when they are multiplied together. Their properties and uses have been explained by various authors, but the full range of them may be underappreciated. Indeed, in the 1139-page book *Matrix Mathematics* [4] the word “bidiagonal” appears on only one page and bidiagonal matrices appear little in the *Handbook of Linear Algebra* [31] apart from in the chapter by Fallat [17].

The purpose of this work is to show the utility of bidiagonal matrices, and in particular to show how factorizations of matrices into bidiagonal factors can be exploited. Our main contributions are as follows, where  $A = B_1 B_2 \dots B_k$  with each  $B_i$  either upper bidiagonal or lower bidiagonal.

- We show that small componentwise perturbations in the  $B_i$  produce small componentwise perturbations in  $A^{-1}$  if the  $B_i$  or the  $B_i^{-1}$  are nonnegative (Theorem 2.3).
- We show that the condition number  $\kappa_\infty(A) = \|A\|_\infty \|A^{-1}\|_\infty$  can be computed in  $O(kn)$  flops when the  $B_i$  are nonnegative or have a checkerboard sign pattern, without explicitly forming  $A$  (section 3).
- We give a unified derivation of backward error bounds and forward error bounds for the computed solution of  $Ax = b$  when  $A$  or  $A^{-1}$  is a product of bidiagonal matrices and the system is solved using the factors (section 4).

\*Draft version of November 6, 2023. This paper is based on the Hans Schneider Prize talk given at the 25th International Linear Algebra Society (ILAS), Conference, Madrid, June 12–16, 2023.

<sup>†</sup>Department of Mathematics, University of Manchester, Manchester, M13 9PL, UK (nick.higham@manchester.ac.uk).

- We show that for a totally nonnegative  $n \times n$  matrix  $A$ ,  $\kappa_\infty(A)$  can be computed in  $O(n^2)$  flops, given a factorization of  $A$  into a product of bidiagonal matrices and that the computed solution is highly accurate (Algorithm 5.5).
- We explore functions of bidiagonal matrices and show that the exponential of a totally nonnegative bidiagonal matrix is totally nonnegative.
- We give new observations on how factorizations involving bidiagonal matrices can help us to understand properties of some well-known matrices (section 8).

Bidiagonal matrices arise in some classical contexts in numerical linear algebra, which we briefly summarize as they will not be the focus of our attention.

*Computing the singular value decomposition (SVD).* The first step of the Golub–Reinsch algorithm for computing the SVD is a two-sided reduction by Householder transformations to upper bidiagonal form  $B$ , as proposed by Golub and Kahan [22]. The SVD of  $B$  is then computed by the QR algorithm implicitly applied to  $B^*B$ , and this can be done in a way that guarantees high relative accuracy in all the computed singular values of  $B$  [10].

*LU factorization of tridiagonal matrices.* If  $A \in \mathbb{C}^{n \times n}$  is tridiagonal and has an LU factorization  $A = LU$  then  $L$  is unit lower bidiagonal and  $U$  is upper bidiagonal.

*Lanczos bidiagonalization.* For large, sparse matrices the solution to a linear system or the least squares solution to an overdetermined system can be computed using a method based on unitary reduction to bidiagonal form by the Lanczos process [5, sec. 7.6], [22], [41].

In perturbation and rounding error analyses products of terms of the form  $1 + \delta_i$  arise. Their distance from 1 will be bounded using the following result [28, Lem. 3.1].

LEMMA 1.1. *If  $|\delta_i| \leq \delta$  and  $\rho_i = \pm 1$  for  $i = 1 : n$ , and  $n\delta < 1$ , then*

$$(1.1) \quad \prod_{i=1}^n (1 + \delta_i)^{\rho_i} = 1 + \theta_n, \quad |\theta_n| \leq \frac{n\delta}{1 - n\delta}.$$

We also need a componentwise bound for perturbations in a matrix product [28, Lem. 3.8]. Here and throughout,  $|A| = (|a_{ij}|)$  and inequalities between matrices hold componentwise.

LEMMA 1.2. *If  $X_j + \Delta X_j \in \mathbb{C}^{n \times n}$  satisfies  $|\Delta X_j| \leq \delta_j |X_j|$  for  $j = 1 : m$  then*

$$\left| \prod_{j=1}^m (X_j + \Delta X_j) - \prod_{j=1}^m X_j \right| \leq \left( \prod_{j=1}^m (1 + \delta_j) - 1 \right) \prod_{j=1}^m |X_j|.$$

We use the standard model of floating-point arithmetic [28, sec. 2.2] and denote by  $u$  the unit roundoff. We need the constant, for  $nu < 1$ ,

$$\gamma_n = \frac{nu}{1 - nu}.$$

We will make use of the one-parameter bidiagonal matrix

$$(1.2) \quad T_n(\theta) = \begin{bmatrix} 1 & \theta & & & \\ & 1 & \theta & & \\ & & 1 & \ddots & \\ & & & \ddots & \theta \\ & & & & 1 \end{bmatrix} \in \mathbb{C}^{n \times n}.$$

**2. Basic Properties of Bidiagonal Matrices.** First we consider the inverse of a nonsingular bidiagonal matrix. It is instructive to look at the  $4 \times 4$  case:

$$\begin{bmatrix} a & x & 0 & 0 \\ & b & y & 0 \\ & & c & z \\ & & & d \end{bmatrix}^{-1} = \begin{bmatrix} \frac{1}{a} & -\frac{x}{ab} & \frac{xy}{abc} & -\frac{xyz}{abcd} \\ & \frac{1}{b} & -\frac{y}{bc} & \frac{yz}{bcd} \\ & & \frac{1}{c} & -\frac{z}{cd} \\ & & & \frac{1}{d} \end{bmatrix}.$$

Notice that every element in the upper triangle is a product of off-diagonal elements of  $B$  and inverses of diagonal elements, that the superdiagonals have alternating signs attached, and that there are no additions. These properties hold for general  $n$ , as the explicit form of the inverse in the following result shows.

LEMMA 2.1. *If  $B \in \mathbb{C}^{n \times n}$  is nonsingular and upper bidiagonal then*

$$(2.1) \quad (B^{-1})_{ij} = \frac{1}{b_{jj}} \prod_{k=i}^{j-1} \left( \frac{-b_{k,k+1}}{b_{kk}} \right), \quad j \geq i.$$

We will make use of the fact that when  $B$  has nonnegative elements,  $B^{-1}$  has a checkerboard (alternating) sign pattern.

We introduce the comparison matrix  $M(A)$  of  $A \in \mathbb{C}^{n \times n}$ :

$$(M(A))_{ij} = \begin{cases} |a_{ii}|, & i = j, \\ -|a_{ij}|, & i \neq j. \end{cases}$$

It is easy to see that

$$(2.2) \quad |B^{-1}| = M(B)^{-1},$$

an observation that we will need later.

Using the representation (2.1) of the inverse we can bound the effect of a componentwise perturbation of  $B$ . Let

$$(2.3) \quad \tau = \frac{(2n-1)\delta}{1-(2n-1)\delta}.$$

THEOREM 2.2. *If  $B \in \mathbb{C}^{n \times n}$  is a nonsingular bidiagonal matrix and  $\Delta B$  is a perturbation satisfying  $|\Delta B| \leq \delta |B|$  then*

$$|(B + \Delta B)^{-1} - B^{-1}| \leq \tau |B^{-1}|,$$

where  $\tau$  is defined in (2.3).

*Proof.* Assume, without loss of generality, that  $B$  is upper bidiagonal. Write  $\Delta b_{ij} = \delta_{ij} b_{ij}$ , where  $|\delta_{ij}| \leq \delta$ . From (2.1) we obtain

$$\begin{aligned} (B + \Delta B)^{-1}_{ij} - (B^{-1})_{ij} &= \frac{1}{b_{jj}(1 + \delta_{jj})} \prod_{k=i}^{j-1} \left( \frac{-b_{k,k+1}(1 + \delta_{k,k+1})}{b_{kk}(1 + \delta_{kk})} \right) - \frac{1}{b_{jj}} \prod_{k=i}^{j-1} \left( \frac{-b_{k,k+1}}{b_{kk}} \right) \\ &= (B^{-1})_{ij} \left( \frac{1}{1 + \delta_{jj}} \prod_{k=i}^{j-1} \left( \frac{1 + \delta_{k,k+1}}{1 + \delta_{kk}} \right) - 1 \right) \\ &= (B^{-1})_{ij} \theta_{2(j-i)+1}, \end{aligned}$$

where  $|\theta_k| \leq \gamma_k = k\delta/(1 - k\delta)$  by Lemma 1.1. □

This result, which is essentially the same as [28, Prob. 22.8], shows that a componentwise relative perturbation in  $B$  produces a componentwise relative perturbation in  $B^{-1}$  at most about  $2n$  times larger: a strong result that does not hold for triangular matrices in general.

We now extend this result to a product of bidiagonal matrices. In all the products of bidiagonal matrices in this paper each matrix can be upper bidiagonal or lower bidiagonal.

**THEOREM 2.3.** *Let  $B = B_1 B_2 \dots B_k \in \mathbb{C}^{n \times n}$ , where the  $B_i$  are nonsingular bidiagonal matrices, and let  $B + \Delta B = (B_1 + \Delta B_1)(B_2 + \Delta B_2) \dots (B_k + \Delta B_k)$ , where  $|\Delta B_i| \leq \delta |B_i|$  for all  $i$ . Then*

$$(2.4) \quad |(B + \Delta B)^{-1} - B^{-1}| \leq ((1 + \tau)^k - 1) |B_k^{-1}| |B_{k-1}^{-1}| \dots |B_1^{-1}|,$$

where  $\tau$  is defined in (2.3), and if the  $B_i$  or the  $B_i^{-1}$  are all nonnegative then

$$(2.5) \quad |(B + \Delta B)^{-1} - B^{-1}| \leq ((1 + \tau)^k - 1) |B^{-1}|.$$

*Proof.* We have

$$\begin{aligned} (B + \Delta B)^{-1} &= (B_k + \Delta B_k)^{-1} (B_{k-1} + \Delta B_{k-1})^{-1} \dots (B_1 + \Delta B_1)^{-1} \\ &= (B_k^{-1} + E_k)(B_{k-1}^{-1} + E_{k-1}) \dots (B_1^{-1} + E_1), \end{aligned}$$

where by Theorem 2.2,  $|E_i| \leq \tau |B_i^{-1}|$ ,  $i = 1 : k$ . Hence by Lemma 1.2,

$$|(B + \Delta B)^{-1} - B^{-1}| \leq ((1 + \tau)^k - 1) |B_k^{-1}| |B_{k-1}^{-1}| \dots |B_1^{-1}|.$$

The bound (2.5) is immediate if the  $B_i^{-1}$  are all nonnegative. If the  $B_i$  are all nonnegative, then (2.5) follows from considering the checkerboard sign pattern of the inverses; see Theorem 3.2 below.  $\square$

The bound (2.5) shows that if the  $B_i$  or the  $B_i^{-1}$  are all nonnegative then componentwise relative perturbations in the  $B_i$  produce componentwise relative perturbation in the inverse of the product at most about a factor  $2nk$  times larger.

Like the inverse, the singular values of a bidiagonal matrix are very well behaved under componentwise perturbations. Let  $\sigma_i(B)$  denote the  $i$ th largest singular value of  $B$ .

**THEOREM 2.4.** *Let  $B \in \mathbb{C}^{n \times n}$  and  $B + \Delta B$  be upper bidiagonal and suppose that  $(B + \Delta B)_{ii} = \alpha_{2i-1} b_{ii}$  and  $(B + \Delta B)_{i,i+1} = \alpha_{2i} b_{i,i+1}$ , where the  $\alpha_i$  are nonzero. Then*

$$\frac{\sigma_i(B)}{\mu} \leq \sigma_i(B + \Delta B) \leq \mu \sigma_i(B), \quad i = 1 : n,$$

where

$$\mu = \prod_{i=1}^{2n-1} \max(|\alpha_i|, |\alpha_i^{-1}|).$$

*Proof.* We can write  $B + \Delta B = D_1 B D_2$ , where

$$D_1 = \text{diag}\left(\alpha_1, \frac{\alpha_1 \alpha_3}{\alpha_2}, \frac{\alpha_1 \alpha_3 \alpha_5}{\alpha_2 \alpha_4}, \dots\right), \quad D_2 = \text{diag}\left(1, \frac{\alpha_2}{\alpha_1}, \frac{\alpha_2 \alpha_4}{\alpha_1 \alpha_3}, \frac{\alpha_2 \alpha_4 \alpha_6}{\alpha_1 \alpha_3 \alpha_5}, \dots\right).$$

An extension for singular values of a result of Ostrowski for eigenvalues [15, Thm. 3.1] gives

$$\frac{\sigma_i(B)}{\|D_1^{-1}\|_2 \|D_2^{-1}\|_2} \leq \sigma_i(B + \Delta B) \leq \sigma_i(B) \|D_1\|_2 \|D_2\|_2.$$

Using  $\|D_1\|_2 \|D_2\|_2 = \max_i |(D_1)_{ii}| \max_i |(D_2)_{ii}| \leq \mu$  (taking account of cancellation in the product) and  $\|D_1^{-1}\|_2 \|D_2^{-1}\|_2 \leq \mu$  gives the result.  $\square$

Theorem 2.4 is from Demmel and Kahan [10, Cor. 2] and the proof is from Eisenstat and Ipsen [15, Cor. 4.2]. The theorem shows that relative perturbations of magnitude at most  $\tau = \max_i |1 - \alpha_i| \ll 1$  to the elements on the diagonal and superdiagonal of an upper bidiagonal matrix produce relative changes of at most  $(1 - \tau)^{2n-1} - 1 \approx (2n - 1)\tau$  in each singular value. This is a much stronger result than for general perturbations of a general  $n \times n$  matrix, where it is only the absolute changes in the singular values that are bounded:  $|\sigma_k(A + \Delta A) - \sigma_k(A)| \leq \sigma_1(\Delta A) = \|\Delta A\|_2$ ,  $k = 1 : n$  [32, Cor. 7.3.5].

Theorem 2.4 does not extend to a product of bidiagonal matrices, as the following example shows. Let

$$A = I = \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -x \\ 0 & 1 \end{bmatrix} =: B_1 B_2,$$

$$A + \Delta A = \begin{bmatrix} 1 & 2x\delta \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & x(1 + \delta) \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -x(1 - \delta) \\ 0 & 1 \end{bmatrix} =: (B_1 + \Delta B_1)(B_2 + \Delta B_2),$$

where  $\delta > 0$ ,  $x > 0$ , and  $x\delta \gg 1$ . Here,  $B_1$  and  $B_2$  have undergone a componentwise relative change  $\delta$ . The singular values of  $A$  are  $\sigma_1 = 1$  and  $\sigma_2 = 1$ , and those of  $A + \Delta A$  are approximately  $\hat{\sigma}_1 = 2x\delta$  and  $\hat{\sigma}_2 = (2x\delta)^{-1}$  (since  $x\delta \gg 1$ ). Hence the relative change in  $\sigma_1$  is  $|\sigma_1 - \hat{\sigma}_1|/\sigma_1 \approx 2x\delta \gg 1$  and that in  $\sigma_2$  is  $|\sigma_2 - \hat{\sigma}_2|/\sigma_2 \approx 1 - 1/(2x\delta) \approx 1$ . We conclude that relative changes in bidiagonal matrices  $B_1, B_2, \dots, B_k$  can induce a much larger relative change in the singular values of their product. The situation is different for a product of *nonnegative* bidiagonal matrices  $B_1, B_2, \dots, B_k$ : small componentwise relative changes in the  $B_i$  produce only small relative changes in the singular values of the product  $B_1, B_2, \dots, B_k$ , as shown by Koev [34, Cor. 7.3].

The next result reveals some further interesting properties of the singular values of a bidiagonal matrix.

**THEOREM 2.5.** *Let  $B \in \mathbb{C}^{n \times n}$  be bidiagonal.*

- (a)  *$|B| = DBF$ , where  $D$  and  $F$  are unitary diagonal matrices. Hence  $B$  and  $|B|$  have the same singular values.*
- (b) *If  $b_{ii}$  and  $b_{i,i+1}$  are nonzero for all  $i$  then the singular values of  $B$  are distinct.*

*Proof.* (a): Let  $D = \text{diag}(d_i)$  and  $F = \text{diag}(f_i)$  with  $f_1 = 1$ . We take  $d_1 = \text{sign}(b_{11})^*$ ,  $f_2 = \text{sign}(d_1 b_{12})^*$ ,  $d_2 = \text{sign}(b_{22} f_2)^*$ ,  $f_3 = \text{sign}(d_2 b_{23})^*$ , and so on, where  $\text{sign}(z) = z/|z|$  if  $z \neq 0$  or 1 otherwise. Then  $|B| = DBF$ , where  $D$  and  $F$  have diagonal elements of modulus 1 and so are unitary. Therefore if  $B = U\Sigma V^*$  is an SVD of  $B$  then  $|B| = (DU)\Sigma(V^*F)$  is an SVD of  $|B|$ .

(b): The singular values of  $B$  are the square roots of the eigenvalues of  $T = |B|^*|B|$ , by (a). The matrix  $T$  is symmetric tridiagonal with positive superdiagonal and subdiagonal elements, so the eigenvalues of  $T$  are distinct [42, Lem. 7.7.1], and hence so are the singular values of  $B$ .  $\square$

It is interesting to note that the SVD codes in both LINPACK [12] and LAPACK [3] reduce  $A \in \mathbb{C}^{m \times n}$  to a real bidiagonal matrix, so that the QR iteration can be carried out in real arithmetic, but they do so in different ways. LINPACK reduces  $A$  to bidiagonal form by Householder transformations and then explicitly carries out the diagonal scaling given in part (a) of Theorem 2.5. LAPACK reduces  $A$  to bidiagonal form using elementary unitary matrices of the form  $P = I - \rho v v^*$  with generally nonreal  $\rho$  that are chosen so that the reduced bidiagonal matrix is real [36].

**3. The Condition Number of a Matrix Product.** Suppose a matrix  $X \in \mathbb{C}^{n \times n}$  is given in factored form  $X = A_1 A_2 \dots A_k$ , where  $A_i \in \mathbb{C}^{n \times n}$  for all  $i$ , and that we wish to compute or estimate the condition number  $\kappa_\infty(X) = \|X\|_\infty \|X^{-1}\|_\infty$  without explicitly forming  $X$ . Initially

we will make no assumptions about the  $A_i$ , but later we will specialize to bidiagonal  $A_i$ . For dense matrices the cost of forming  $X$  is  $2(k-1)n^3$  flops, whereas we would like to compute or estimate  $\kappa_\infty(X)$  at the cost of a few matrix–vector products with  $X$ , that is, in a small multiple of  $2(k-1)n^2$  flops.

The condition number estimation problem is well studied [28, Chap. 15]. Here we focus on the problem of *exactly* computing the condition number. Recall that the  $\infty$ -norm satisfies

$$\|X\|_\infty = \| |X| \|_\infty = \| |X|e \|_\infty,$$

where  $e = [1, 1, \dots, 1]^T$ .

In general we cannot compute  $\|A_1 A_2 \dots A_k\|_\infty$  without forming the matrix product. However, if the equality

$$(3.1) \quad |A_1 A_2 \dots A_k| = |A_1| |A_2| \dots |A_k|$$

holds then

$$(3.2) \quad \|A_1 A_2 \dots A_k\|_\infty = \| |A_1| |A_2| \dots |A_k| \|_\infty = \| |A_1| |A_2| \dots |A_k| e \|_\infty$$

and we can evaluate the right-hand side in  $O(kn^2)$  flops as opposed to the  $O(kn^3)$  flops that are required if we explicitly form the product. If the  $A_i$  are bidiagonal then the costs are  $3kn$  flops compared with up to  $O(kn^2)$  flops if the product is explicitly formed, since in general the product fills in.

The equality (3.1) obviously holds when the  $B_i$  are all nonnegative. It can also hold because all additions in the product  $A_1 A_2 \dots A_k$  are of like-signed numbers, so that there is no cancellation. Important such cases are when the  $A_i$  are nonnegative and when each  $A_i$  has a checkerboard (alternating) sign pattern, which can be expressed as

$$(3.3) \quad A_i = \pm \Sigma |A_i| \Sigma, \quad i = 1 : k,$$

where

$$(3.4) \quad \Sigma = \text{diag}(1, -1, 1, \dots, (-1)^{n-1}).$$

**THEOREM 3.1.** *If the matrices  $A_i$ ,  $i = 1 : k$ , satisfy (3.3) then*

$$(3.5) \quad A_1 A_2 \dots A_k = \pm \Sigma |A_1| |A_2| \dots |A_k| \Sigma$$

and hence

$$(3.6) \quad |A_1 A_2 \dots A_k| = |A_1| |A_2| \dots |A_k|.$$

*Proof.* If the  $A_i$  satisfy (3.3) then

$$A_1 A_2 \dots A_k = \pm \Sigma |A_1| \Sigma \cdot \Sigma |A_2| \Sigma \dots \Sigma |A_k| \Sigma = \pm \Sigma |A_1| |A_2| \dots |A_k| \Sigma,$$

which is (3.5), and (3.6) follows immediately.  $\square$

We conclude that if the  $A_i$  are nonnegative or have a checkerboard sign pattern then we can compute  $\|A_1 A_2 \dots A_k\|_\infty$  in  $O(kn^2)$  flops.

If  $B_1, B_2, \dots, B_k$  are bidiagonal and nonnegative then from Lemma 2.1 it is clear that  $B_i^{-1}$  has a checkerboard sign pattern, that is, it satisfies (3.3). Therefore by (3.6),

$$(3.7) \quad |B_k^{-1} B_{k-1}^{-1} \dots B_1^{-1}| = |B_k^{-1}| |B_{k-1}^{-1}| \dots |B_1^{-1}|.$$

The same is true if the  $B_i$  have a checkerboard sign pattern.

**THEOREM 3.2.** *Let  $B_1, B_2, \dots, B_k \in \mathbb{R}^{n \times n}$  be nonsingular bidiagonal matrices. If  $B_i$  is nonnegative for all  $i$  or has a checkerboard sign pattern for all  $i$  then*

$$(3.8) \quad |B_k^{-1} B_{k-1}^{-1} \dots B_1^{-1}| = |B_k^{-1}| |B_{k-1}^{-1}| \dots |B_1^{-1}| = M(B_k)^{-1} M(B_{k-1})^{-1} \dots M(B_1)^{-1}.$$

*Proof.* For nonnegative  $B_i$  the result follows from (3.7) on recalling (2.2). From (2.1) it is clear that  $B_i$  having a checkerboard sign pattern is equivalent to either  $B_i^{-1}$  or  $-B_i^{-1}$  being nonnegative and equal to  $M(B_i)^{-1}$ , which gives the second part of the result.  $\square$

From (3.8) we have

$$(3.9) \quad \|B_k^{-1} B_{k-1}^{-1} \dots B_1^{-1}\|_\infty = \|M(B_k)^{-1} M(B_{k-1})^{-1} \dots M(B_1)^{-1} e\|_\infty,$$

and the right-hand side can be computed in  $3kn$  flops, whereas explicitly forming the product on the left (using substitutions) costs  $3kn^2/2$  flops. We conclude that when the  $B_i$  are nonnegative for all  $i$  or all have a checkerboard sign pattern,  $\kappa_\infty(B_1 B_2 \dots B_k)$  can be computed exactly in  $6kn$  flops. Since  $\|A\|_1 = \|A^T\|_\infty$ , the 1-norm condition number can be computed at the same cost by working with the transpose of the product.

In the case  $k = 1$ , (3.9) reduces to the result that  $\|B^{-1}\|_\infty = \|M(B)^{-1}\|_\infty = \|M(B)^{-1} e\|_\infty$  [25, sec. 2].

We can also compute the condition number of Skeel [44],

$$\text{cond}(A, x) = \frac{\| |A^{-1}| |A| |x| \|_\infty}{\|x\|_\infty},$$

exactly in  $6kn$  flops for  $A = B_1 B_2 \dots B_k$  with nonnegative  $B_i$ :

$$\text{cond}(B_1 B_2 \dots B_k, x) = \frac{\| M(B_k)^{-1} \dots M(B_1)^{-1} B_1 \dots B_k |x| \|_\infty}{\|x\|_\infty}.$$

If the  $B_i$  have checkerboard sign patterns then the same formula holds with  $B_1 B_2 \dots B_k$  replaced by  $|B_1| |B_2| \dots |B_k|$ .

We will make use of (3.9) for totally nonnegative matrices in Section 5.

**4. Linear Systems.** We consider a linear system  $Ax = b$  in which  $A$  is either a product of bidiagonal matrices or a product of inverses of bidiagonal matrices. Our interest is in what can be said about the backward error and forward error when such a system is solved in floating-point arithmetic.

**4.1. Product of Bidiagonal Matrices.** Suppose  $A = B_1 B_2 \dots B_k$  is a product of  $k$  bidiagonal matrices. We can solve the system by solving  $k$  bidiagonal systems by substitution. Standard rounding error analysis [28, Lem. 8.2] shows that the computed  $\widehat{x}$  satisfies

$$(4.1) \quad (B_1 + \Delta B_1)(B_2 + \Delta B_2) \dots (B_k + \Delta B_k) \widehat{x} = b, \quad |\Delta B_i| \leq \gamma_2 |B_i|, \quad i = 1 : k.$$

Hence the residual is

$$\begin{aligned} |b - B_1 B_2 \dots B_k \widehat{x}| &= |((B_1 + \Delta B_1)(B_2 + \Delta B_2) \dots (B_k + \Delta B_k) - B_1 B_2 \dots B_k) \widehat{x}| \\ &\leq ((1 + \gamma_2)^k - 1) |B_1| |B_2| \dots |B_k| |\widehat{x}|, \end{aligned}$$

by Lemma 1.2. If the  $B_i$  are all nonnegative or, by Theorem 3.1, if they have a checkerboard sign pattern, then the bound becomes

$$(4.2) \quad |b - A \widehat{x}| \leq ((1 + \gamma_2)^k - 1) |A| |\widehat{x}| = (2ku + O(u^2)) |A| |\widehat{x}|,$$

which shows that the componentwise relative backward error is small—an ideal backward error result. We note that this result has used the triangularity of the  $B_i$  but not their bidiagonal structure (except through the constant in (4.1)).

To obtain a forward error bound we rewrite (4.1) as

$$\widehat{x} = (B_k + \Delta B_k)^{-1}(B_{k-1} + \Delta B_{k-1})^{-1} \dots (B_1 + \Delta B_1)^{-1}b.$$

Then

$$\begin{aligned} |\widehat{x} - x| &\leq |(B_k + \Delta B_k)^{-1}(B_{k-1} + \Delta B_{k-1})^{-1} \dots (B_1 + \Delta B_1)^{-1} - B_k^{-1}B_{k-1}^{-1} \dots B_1^{-1}| |b| \\ (4.3) \quad &\leq ((1 + \tau)^k - 1) |B_k^{-1}| |B_{k-1}^{-1}| \dots |B_1^{-1}| |b| \end{aligned}$$

by Theorem 2.3, where

$$(4.4) \quad \tau = \frac{(2n-1)\gamma_2}{1 - (2n-1)\gamma_2}.$$

If the  $B_i$  are all nonnegative or have a checkerboard sign pattern then by Theorem 3.2 this inequality becomes

$$(4.5) \quad |\widehat{x} - x| \leq (2k(2n-1)u + O(u^2)) |A^{-1}| |b|.$$

The bound (4.5) is a strong forward error bound because it is the same as a bound for the change in  $x$  induced by a small componentwise relative perturbation of  $b$ :  $b \rightarrow b + \Delta b$  with  $|\Delta b| \leq 4knu|b|$  [28, Thm. 7.4].

**4.2. Product of Inverses of Bidiagonal Matrices.** Now suppose that it is  $A^{-1}$  rather than  $A$  that is a product of bidiagonal matrices:  $A^{-1} = B_1 B_2 \dots B_k$ . Now we solve  $Ax = b$  by forming  $x = A^{-1}b = B_1 B_2 \dots B_k b$  and the computed  $\widehat{x}$  satisfies

$$(4.6) \quad \widehat{x} = (B_1 + \Delta B_1)(B_2 + \Delta B_2) \dots (B_k + \Delta B_k)b, \quad |\Delta B_i| \leq \gamma_2 |B_i|, \quad i = 1 : k.$$

Then the forward error is

$$\begin{aligned} |\widehat{x} - x| &= |((B_1 + \Delta B_1)(B_2 + \Delta B_2) \dots (B_k + \Delta B_k) - B_1 B_2 \dots B_k)b|, \\ (4.7) \quad &\leq ((1 + \gamma_2)^k - 1) |B_1| |B_2| \dots |B_k| |b|, \end{aligned}$$

by Lemma 1.2. If the  $B_i$  are all nonnegative or have a checkerboard sign pattern then by Theorem 3.1,  $|B_1| |B_2| \dots |B_k| = |B_1 B_2 \dots B_k|$ , so

$$(4.8) \quad |\widehat{x} - x| \leq ((1 + \gamma_2)^k - 1) |A^{-1}| |b|.$$

Now we turn to the residual. Note first that by (4.6),

$$b = (B_k + \Delta B_k)^{-1}(B_{k-1} + \Delta B_{k-1})^{-1} \dots (B_1 + \Delta B_1)^{-1} \widehat{x}.$$

Hence

$$|b - A\widehat{x}| = |[(B_k + \Delta B_k)^{-1}(B_{k-1} + \Delta B_{k-1})^{-1} \dots (B_1 + \Delta B_1)^{-1} - B_k^{-1}B_{k-1}^{-1} \dots B_1^{-1}] \widehat{x}|$$

and by Lemma 1.2 and Theorem 2.3 we obtain, with  $\tau$  given by (4.4),

$$\begin{aligned} |b - A\widehat{x}| &\leq ((1 + \tau)^k - 1) |B_k^{-1}| |B_{k-1}^{-1}| \dots |B_1^{-1}| |\widehat{x}| \\ &= (2k(2n-1)u + O(u^2)) |B_k^{-1}| |B_{k-1}^{-1}| \dots |B_1^{-1}| |\widehat{x}|. \end{aligned}$$

If the  $B_i$  are all nonnegative or have a checkerboard sign pattern then by Theorem 3.2 this bound can be written

$$(4.9) \quad |b - A\hat{x}| \leq (2k(2n-1)u + O(u^2))|A||\hat{x}|,$$

which again shows a small componentwise relative backward error.

Our conclusion is that whether it is  $A$  or  $A^{-1}$  that is a product of bidiagonal matrices we have the same satisfactory form of forward error bounds (4.5) and (4.8) and residual bounds (4.2) and (4.9) when the  $B_i$  are all nonnegative or have a checkerboard sign pattern.

**4.3. Application to Vandermonde Systems.** An application of these results is to the Björck–Pereyra algorithm for solving a Vandermonde system  $Vy = b$  in  $O(n^2)$  flops [6], where  $V = (x_j^{i-1}) \in \mathbb{C}^{n \times n}$  for given points  $x_i \in \mathbb{C}$ . This algorithm uses a factorization of  $V^{-1}$  into a product of  $2n-2$  bidiagonal matrices  $B_{2n-2}, \dots, B_1$  given in terms of the points  $x_i$ . When  $0 \leq x_1 < x_2 < \dots < x_n$  the bidiagonal factors have positive diagonal and nonpositive off-diagonal elements. Therefore the  $B_i$  have a checkerboard sign pattern and so  $|B_{2n-2}| \dots |B_1| = |B_{2n-2} \dots B_1| = |A^{-1}|$  by (3.7). From (4.8) and (4.9) we have

$$\begin{aligned} |\hat{y} - y| &\leq (2(2n-2)u + O(u^2))|V^{-1}||b|, \\ |b - V\hat{y}| &\leq (2(2n-2)(2n-1)u + O(u^2))|V||\hat{y}|, \end{aligned}$$

which reproduce [26, Thm. 2.3] and the monomial case of [27, Cor. 4.1], respectively. Since  $V^{-1}$  has a checkerboard sign pattern, if  $(-1)^i b_i \geq 0$  then  $|V^{-1}||b| = |V^{-1}b| = |y|$ , and  $\hat{y}$  therefore has a small componentwise relative error. The analysis in [27] makes use of the bidiagonal factorization, but that in [26] does not.

**4.4. Application to Pascal Systems.** We give a numerical illustration of the use of the bidiagonal factorization for solving the linear system  $P_n x = b$ , where  $P_n$  is the symmetric positive definite  $n \times n$  Pascal matrix with

$$(4.10) \quad p_{ij} = \binom{i+j-2}{j-1} = \frac{(i+j-2)!}{(i-1)!(j-1)!}$$

and  $b = e_n/n$ , where  $e_n$  is the  $n$ th unit vector. The Pascal matrix has a known factorization as a product of  $2n-1$  bidiagonal matrices, as we explain in section 8.3. We solve the system using the bidiagonal factorization, solving the bidiagonal systems by substitution. We also solve the system for the explicitly formed  $P$  using the MATLAB backslash operator (which exploits the symmetric positive definiteness of  $P_n$  but not its bidiagonal factorization). The working precision is double precision, with  $u \approx 1.1 \times 10^{-16}$ . Table 4.1 shows the relative errors  $\|x - \hat{x}\|_\infty / \|x\|_\infty$ , for which we take as the exact solution  $x$  the solution computed at a precision of 500 decimal digits using the Multiprecision Computing Toolbox [39] and then rounded to double precision. We restrict to  $n \leq 25$  to ensure that  $P$  is exactly representable at the working precision. We see that substitution with the bidiagonal factorization yields errors of  $O(u)$  that satisfy the bound (4.5), whereas the MATLAB backslash function produces much larger errors, which usually exceed (4.5).

**5. Totally Nonnegative Matrices.** A matrix  $A \in \mathbb{R}^{n \times n}$  is totally nonnegative if every minor (determinant of a square submatrix) is nonnegative and totally positive if every minor is positive. We will need the following key result, which is a direct consequence of the Binet–Cauchy theorem on determinants [32, sec. 0.8.7].

**THEOREM 5.1.** *If  $A, B \in \mathbb{R}^{n \times n}$  are totally nonnegative then so is  $AB$ .*

Table 4.1: Relative errors for the computed solution to a linear system  $P_n x = b$  with  $P_n$  the  $n \times n$  Pascal matrix.

$n$	Relative errors		
	Bidiagonal factorization	$P \backslash b$	Error bound (4.5)
5	9.25e-17	9.25e-16	7.99e-15
10	1.50e-16	4.94e-9	3.80e-14
15	6.36e-17	1.05e-3	9.02e-14
20	1.34e-16	3.12e-12	1.65e-13
25	1.68e-16	2.76e-11	2.61e-13

Bidiagonal matrices play a key role in the theory of totally nonnegative matrices. Indeed a nonnegative bidiagonal matrix is totally nonnegative. In the proof of this result we will need the elementary lower bidiagonal matrix

$$(5.1) \quad L_k(\ell_{k+1,k}) = I + \ell_{k+1,k} e_{k+1} e_k^T,$$

which differs from the identity matrix only in the  $(k+1, k)$  position, which contains  $\ell_{k+1,k}$ .

**THEOREM 5.2.** *A bidiagonal matrix  $B \in \mathbb{R}^{n \times n}$  with nonnegative elements is totally nonnegative.*

*Proof.* Without loss of generality we take  $B$  to be lower bidiagonal. We first assume that  $B$  is nonsingular. Since  $0 \neq \det(B) = b_{11} b_{22} \dots b_{nn}$ , the  $b_{ii}$  are all positive, so with  $D = \text{diag}(b_{ii})$  and  $\ell_{i+1,i} = b_{i+1,i}/b_{i+1,i+1} \geq 0$ ,  $i = 1 : n-1$ , we can write

$$(5.2) \quad B = D \begin{bmatrix} 1 & & & \\ \ell_{21} & 1 & & \\ & \ell_{32} & \ddots & \\ & & \ddots & \ddots \\ & & & \ell_{n,n-1} & 1 \end{bmatrix} \equiv DL.$$

Since  $D$  is clearly totally nonnegative, by Theorem 5.1 it suffices to show that  $L$  is totally nonnegative.

For  $n = 4$  we have

$$L = \begin{bmatrix} 1 & & & \\ \ell_{21} & 1 & & \\ & \ell_{32} & 1 & \\ & & \ell_{43} & 1 \end{bmatrix} = \begin{bmatrix} 1 & & & \\ \ell_{21} & 1 & & \\ & & 1 & \\ & & & 1 \end{bmatrix} \begin{bmatrix} 1 & & & \\ & 1 & & \\ & \ell_{32} & 1 & \\ & & & 1 \end{bmatrix} \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & \ell_{43} & 1 \end{bmatrix},$$

and this factorization clearly generalizes to

$$(5.3) \quad L = L_1(\ell_{21}) L_2(\ell_{32}) \dots L_{n-1}(\ell_{n,n-1}),$$

where  $L_k(\ell_{k+1,k})$  is the elementary lower bidiagonal matrix (5.1). It is easy to see that  $L_k(\ell_{k+1,k})$  is totally nonnegative for all  $k$ , so  $L$  is totally nonnegative by Theorem 5.1.

If  $B$  is singular then consider the bidiagonal matrix  $B(\epsilon) = B + \epsilon I$ , which is nonsingular for  $\epsilon > 0$ . By the argument above,  $B(\epsilon)$  is totally nonnegative for  $\epsilon > 0$ . Any minor of  $B(\epsilon)$

is the determinant of a submatrix of  $B(\epsilon)$ , which is a polynomial in  $\epsilon$ , so it is continuous in  $\epsilon$ . This minor is nonnegative for all  $\epsilon > 0$  and so must remain nonnegative in the limit as  $\epsilon \rightarrow 0$ . Therefore  $B = B(0)$  is totally nonnegative.  $\square$

Even if  $B$  is not totally nonnegative, there is an associated totally nonnegative matrix.

**THEOREM 5.3.** *If  $B \in \mathbb{R}^{n \times n}$  is nonsingular and bidiagonal then  $M(B)^{-1}$  is totally nonnegative.*

*Proof.* Assuming that  $B = L$  is lower bidiagonal, by (5.2) and (5.3),

$$M(B) = M(DL) = |D|M(L) = |D|L_1(-|\ell_{21}|)L_2(-|\ell_{32}|) \dots L_{n-1}(-|\ell_{n,n-1}|)$$

and  $L_k(-|\ell_{k+1,k}|)^{-1} = L_k(|\ell_{k+1,k}|)$ , so  $M(B)^{-1} = L_{n-1}(|\ell_{n,n-1}|)L_{n-2}(|\ell_{n-1,n-2}|) \dots L_1(|\ell_{21}|) \times |D|^{-1}$ , which is a product of totally nonnegative matrices and hence is totally nonnegative.  $\square$

The next result shows that any nonsingular totally nonnegative matrix can be written as a product of nonnegative bidiagonal matrices.

**THEOREM 5.4.** *A nonsingular matrix  $A \in \mathbb{R}^{n \times n}$  is totally nonnegative if and only if it can be factorized as*

$$(5.4) \quad A = L_{n-1}L_{n-2} \dots L_1 D U_1 U_2 \dots U_{n-1},$$

where  $D$  is a diagonal matrix with positive diagonal entries and  $L_i$  and  $U_i$  are unit lower and unit upper bidiagonal matrices, respectively, with the first  $i-1$  entries along the subdiagonal of  $L_i$  and  $U_i^T$  zero and the rest nonnegative.

The factorization (5.4) is essentially an LU factorization in which  $L$  and  $U$  have been factorized into a product of specially structured nonnegative bidiagonal matrices.

Theorem 5.4 is from Gasca and Peña [21, Thm. 4.2]. Fallat and Johnson [19, sec. 2.0] summarize the history of different forms of this factorization.

Since the bidiagonal matrices in the factorization (5.4) are all nonnegative, by (3.9) we have

$$(5.5) \quad \|A^{-1}\|_\infty = \|M(U_{n-1})^{-1} \dots M(U_1)^{-1} D^{-1} M(L_1)^{-1} \dots M(L_{n-1})^{-1} e\|_\infty,$$

and so we can compute  $\|A^{-1}\|_\infty$  by  $2(n-1)$  substitutions in  $O(n^2)$  flops for any nonsingular totally nonnegative matrix given the factorization (5.4).

Let  $\widehat{c} = \text{fl}(\|A^{-1}\|_\infty)$ . Taking  $\infty$ -norms in (4.5) with  $b = e$  gives, using the triangle inequality,

$$(5.6) \quad \frac{|\widehat{c} - \|A^{-1}\|_\infty|}{\|A^{-1}\|_\infty} \leq dn^2 u$$

for a modest constant  $d$ . Therefore  $\widehat{c}$  is highly accurate, essentially because there is no cancellation in evaluating (5.5): all additions are of nonnegative quantities. Standard methods for evaluating  $\|A^{-1}\|_\infty$  for general  $A$  only satisfy  $|\widehat{c} - \|A^{-1}\|_\infty|/\|A^{-1}\|_\infty \leq cn^3 \kappa_\infty(A)u$ , which is the best that can be expected in general because the condition number of  $\kappa_\infty(A)$  is  $\kappa_\infty(A)$  [24].

To obtain  $\kappa_\infty(A)$  we need  $\|A\|_\infty$ , which can either be computed from  $A$  if it is explicitly known, or from  $\|A\|_\infty = \|L_{n-1}L_{n-2} \dots L_1 D U_1 U_2 \dots U_{n-1} e\|_\infty$  otherwise. We summarize the computations in an algorithm.

**ALGORITHM 5.5.** *This algorithm computes  $c = \kappa_\infty(A)$  for a totally nonnegative matrix  $A$  given the factorization (5.4).*

Table 5.1: Condition numbers and relative errors for the Hilbert matrix.

$n$	$\kappa_\infty(H_n)$	Relative error for Algorithm 5.5
4	2.84e4	1.28e-16
8	3.39e10	2.25e-16
16	5.06e22	3.67e-17
32	1.36e47	1.75e-15
64	1.10e96	1.77e-15

```

1 If A is explicitly known
2    $\alpha = \|A\|_\infty$ 
3 else
4    $\alpha = \|L_{n-1}L_{n-2} \dots L_1 D U_1 U_2 \dots U_{n-1} e\|_\infty$ 
5 end
6 Compute  $\beta = \|M(U_{n-1})^{-1} \dots M(U_1)^{-1} D^{-1} M(L_1)^{-1} \dots M(L_{n-1})^{-1} e\|_\infty$ 
   by substitutions.
7  $c = \alpha\beta$ 

```

How do we obtain the parameters in the factorization (5.4)? In some cases they are known from the construction of the matrix. Formulas are known for totally positive Vandermonde matrices and Cauchy matrices [34, eqs. (3.5), (3.6)] and a variety of Vandermonde-type matrices [9]. For totally positive matrices determinantal formulas for the parameters are available [34, Prop. 3.1]. Assuming the determinants can be computed accurately, in all these cases the parameters can be evaluated to high relative accuracy, and so in view of Theorem 2.3 the errors in the evaluation of the parameters do not affect the form of the bound (5.6).

We give two numerical experiments in MATLAB to illustrate the accuracy of the condition number evaluation. We take as the exact condition number the one computed at a precision of 500 decimal digits using the Multiprecision Computing Toolbox [39] and then rounded to double precision.

First, in Table 5.1 we show the relative errors in computing the  $\infty$ -norm condition number of the Hilbert matrix  $H_n$ , which has  $(i, j)$  element  $1/(i + j - 1)$  and is totally positive. The parameters in the bidiagonal factorization (5.4) are computed using the function `TNcauchyBD` from the `TNTool` toolbox.<sup>1</sup> We see that even extremely large condition numbers are obtained to high accuracy.

Next we consider the Pascal matrix (4.10), which is totally positive [19, Ex. 0.1.6]. Since this matrix is exactly representable at the working precision for  $n$  up to around 25, we can compare Algorithm 5.5 with the MATLAB `cond` function. We see from the results in Table 5.2 that the MATLAB function loses accuracy as  $n$  increases while Algorithm 5.5 returns a result correct to the working precision.

Another use of the factorization of Theorem 5.4 is to construct totally nonnegative matrices by choosing the  $n^2$  parameters that make up the  $L_i$ ,  $D$ , and the  $U_i$ . The function call

```
A = anymatrix('core/totally_nonneg', X)
```

in the `Anymatrix` toolbox [30] constructs an  $n \times n$  totally nonnegative matrix  $A$  from parameters given in the  $n \times n$  matrix  $X$ , whose format is as suggested in [34, sec. 4]. The Pascal matrix is generated when  $X = \text{ones}(n)$ . In a call

<sup>1</sup><http://www.math.sjsu.edu/~koev/software/TNTool.html>

Table 5.2: Condition numbers and relative errors for the Pascal matrix.

$n$	$\kappa_\infty(P_n)$	Relative errors	
		Algorithm 5.5	cond(P_n, inf)
5	1.56e4	0.00	0.00
10	8.13e9	0.00	1.49e-11
15	5.77e15	0.00	2.19e-8
20	4.50e21	4.66e-17	3.41e-4
25	3.81e27	1.70e-17	3.17e-2

$A = \text{anymatrix}(\text{'core/totally\_nonneg'}, n)$

the parameters are chosen randomly, and this is a convenient way to generate random totally nonnegative matrices.

Koev [34, sec. 7], [35] shows that small relative changes in the parameters in the factorization (5.4) produce small relative changes in the determinant, the eigenvalues, and the singular values. In [34] he develops algorithms for accurate computation of eigenvalues and the SVD of nonsingular totally nonnegative matrices, given an accurate bidiagonal factorization, by carrying out transformations on the bidiagonal factorization in such a way that no subtractions occur.

For later use, we note a useful theorem about the eigenvalues of a totally nonnegative matrix [18, Thm. 3.3].

**THEOREM 5.6.** *If  $A \in \mathbb{R}^{n \times n}$  is totally nonnegative and irreducible then its eigenvalues are real and nonnegative and the positive eigenvalues are distinct.*

Note that the irreducibility requirement in the theorem means that it cannot be applied to triangular matrices, so there is no contradiction to the fact that the totally nonnegative matrix  $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$  (for example) has repeated nonzero eigenvalues.

**6. Matrix Functions and Polynomial Evaluation and Interpolation.** Bidiagonal matrices are intimately connected with polynomial evaluation and interpolation. Horner's method for evaluating a polynomial at a point  $\alpha$  can be expressed as the solution of a linear system with coefficient matrix  $T_n(-\alpha)$  [28, sec. 5.2], where  $T_n$  is defined in (1.2). Premultiplying a vector by  $T_n(-1)^T$  corresponds to forming a backward difference, and a subsequent multiplication by a diagonal matrix yields divided differences [28, sec. 5.3]. In fact, an explicit formula for a function of a bidiagonal matrix is available in terms of divided differences. Recall that divided differences of a function  $f$  at points  $x_k$  are defined recursively by (see, e.g. [7, Chap. 2] or [29, sec. B.16])

$$\begin{aligned}
 f[x_k] &= f(x_k), \\
 f[x_0, x_1, \dots, x_{k+1}] &= \begin{cases} \frac{f[x_1, x_2, \dots, x_{k+1}] - f[x_0, x_1, \dots, x_k]}{x_{k+1} - x_0}, & x_0 \neq x_{k+1}, \\ \frac{f^{(k+1)}(x_{k+1})}{(k+1)!}, & x_0 = x_{k+1}, \end{cases}
 \end{aligned}
 \tag{6.1}$$

where, since  $f[x_1, x_2, \dots, x_{k+1}]$  does not depend on the order of its arguments, we assume without loss of generality that equal points are contiguous.

THEOREM 6.1. If  $B \in \mathbb{C}^{n \times n}$  is upper bidiagonal then  $F = f(B)$  is upper triangular with  $f_{ii} = f(t_{ii})$  and

$$(6.2) \quad f_{ij} = b_{i,i+1} b_{i+1,i+2} \cdots b_{j-1,j} f[b_{ii}, b_{i+1,i+1}, \dots, b_{jj}], \quad j > i.$$

*Proof.* The formula (6.2) is a special case of the formula for  $f(T)$ , where  $T$  is upper triangular, given in Davis [8], Descoux [11], and Van Loan [45].  $\square$

Lemma 2.1 is the special case of Theorem 6.1 with  $f(x) = 1/x$ . Since  $f[\lambda, \lambda, \dots, \lambda] = f^{(n-1)}(\lambda)/(n-1)!$ , another special case is the formula for a function of an  $m \times m$  Jordan block [29, sec. 1.2]

$$(6.3) \quad f\left(\begin{bmatrix} \lambda & 1 & & \\ & \lambda & \ddots & \\ & & \ddots & 1 \\ & & & \lambda \end{bmatrix}\right) = \begin{bmatrix} f(\lambda) & f'(\lambda) & \cdots & \frac{f^{(m-1)}(\lambda)}{(m-1)!} \\ & f(\lambda) & \ddots & \vdots \\ & & \ddots & f'(\lambda) \\ & & & f(\lambda) \end{bmatrix}.$$

Yet another special case is

$$f\left(\begin{bmatrix} \lambda_1 & 1 & & \\ & \lambda_2 & \ddots & \\ & & \ddots & 1 \\ & & & \lambda_n \end{bmatrix}\right)_{1n} = f[\lambda_1, \lambda_2, \dots, \lambda_n],$$

which is a result of Opitz [40] and is used in computing divided differences of the exponential by McCurdy, Ng, and Parlett [37].

A natural question is whether a function of a nonnegative bidiagonal matrix is totally nonnegative. For the exponential, the answer is yes.

THEOREM 6.2. If  $B \in \mathbb{R}^{n \times n}$  is a nonnegative bidiagonal matrix then  $e^B$  is totally nonnegative.

*Proof.* Consider the formula [29, sec. 10.1]  $e^A = \lim_{m \rightarrow \infty} (I + A/m)^m$ , valid for any  $A$ , where  $m \in \mathbb{Z}$ . For nonnegative bidiagonal  $B$ ,  $I + B/m \geq 0$  for all  $m > 0$ , so by Theorem 5.2  $I + B/m$  is totally nonnegative and therefore  $X_m = (I + B/m)^m$  is totally nonnegative for all  $m > 0$  by Theorem 5.1. Suppose that  $\lim_{m \rightarrow \infty} X_m$  is not totally nonnegative, so that some submatrix with indices  $(\alpha, \beta)$  has negative determinant. Let  $x_m = \det(X_m(\alpha, \beta))$ . Then  $\lim_{m \rightarrow \infty} x_m < 0$  but  $x_m > 0$  for all  $m$ , which is a contradiction, so  $e^B$  is totally nonnegative.  $\square$

Note that Theorem 6.2 does not generalize to wider bandwidths, as the example

$$\exp\left(\begin{bmatrix} 1 & 1 & 1 \\ & 1 & 1 \\ & & 1 \end{bmatrix}\right) = \begin{bmatrix} e & e & 3e/2 \\ & e & e \\ & & e \end{bmatrix}$$

shows, since the  $(1:2, 3:4)$  submatrix has negative determinant.

**7. Upper Triangular Toeplitz matrices.** Upper triangular Toeplitz matrices  $T \in \mathbb{C}^{n \times n}$  can be written in the form

$$T = \begin{bmatrix} t_0 & t_1 & \cdots & t_{n-1} \\ & t_0 & \ddots & \vdots \\ & & \ddots & t_1 \\ & & & t_0 \end{bmatrix} = \sum_{j=1}^n t_{j-1} N^{j-1},$$

where  $N$  is upper bidiagonal with a superdiagonal of ones:

$$N = \begin{bmatrix} 0 & 1 & & \\ & 0 & \ddots & \\ & & \ddots & 1 \\ & & & 0 \end{bmatrix},$$

Note that  $N^n = 0$ . It follows that the product of two upper triangular Toeplitz matrices is again upper triangular Toeplitz and that upper triangular Toeplitz matrices commute. Furthermore, since  $f(T)$  is a polynomial in  $T$ , it follows that  $f(T)$  is also upper triangular and Toeplitz. Note that as a special case, if  $B$  is a Toeplitz bidiagonal matrix with  $b_{ii} = b$  and  $b_{i,i+1} = c$  then Theorem 6.1 gives  $f(B)_{ij} = c^{j-i} f[b, b, \dots, b] = c^{j-i} f^{(j-i)}(b)/(j-i)!$ , of which (6.3) is a special case.

**8. Exploiting Factorizations Into Products of Bidiagonal Matrices.** In this section we show how factorizations involving bidiagonal matrices or their inverses can provide valuable information about particular matrices.

**8.1. The Frank Matrix.** In 1958 Frank [20] reported that his algorithms had difficulty computing accurately the smaller eigenvalues of the  $n \times n$  upper Hessenberg matrix

$$F_n = \begin{bmatrix} n & n-1 & n-2 & \dots & 2 & 1 \\ n-1 & n-1 & n-2 & \dots & 2 & 1 \\ 0 & n-2 & n-2 & \dots & 2 & 1 \\ \vdots & 0 & \ddots & \ddots & \vdots & 1 \\ \vdots & \vdots & \dots & 2 & 2 & 1 \\ 0 & 0 & \dots & 0 & 1 & 1 \end{bmatrix}.$$

Wilkinson [47, sec. 8] [48, pp. 92–93] showed that the difficulties are caused by the sensitivity of the eigenvalues to perturbations in the matrix, which can be measured by the condition number of a simple eigenvalue  $\lambda$ :  $\kappa_2(\lambda) = \|y\|_2 \|x\|_2 / |y^* x|$ , where  $x$  and  $y$  are right and left eigenvectors, respectively, corresponding to  $\lambda$ . The eigenvalues are known to be real and positive, and they can be expressed in terms of the zeros of Hermite polynomials [13], [46]. However, in none of these references is it shown that the eigenvalues are distinct, which is necessary for the eigenvalue condition numbers to be defined.

If we subtract row  $k+1$  from row  $k$  for  $k = 1 : n-1$ , we obtain a lower bidiagonal matrix. For  $n = 4$  this transformation can be written

$$\begin{bmatrix} 1 & -1 & & \\ & 1 & -1 & \\ & & 1 & -1 \\ & & & 1 \end{bmatrix} \begin{bmatrix} 4 & 3 & 2 & 1 \\ 3 & 3 & 2 & 1 \\ 2 & 2 & 1 & 1 \\ 1 & 1 & & \end{bmatrix} = \begin{bmatrix} 1 & & & \\ 3 & 1 & & \\ & 2 & 1 & \\ & & 1 & 1 \end{bmatrix},$$

and in general we have

$$(8.1) \quad F_n = T_n(-1)^{-1} \begin{bmatrix} 1 & & & \\ n-1 & 1 & & \\ & n-2 & 1 & \\ & & \ddots & \ddots \\ & & & 1 & 1 \end{bmatrix} \equiv T_n(-1)^{-1} L,$$

where  $T_n$  is defined in (1.2). This is equivalent to a factorization noted by Rutishauser [43, sec. 9]. Note that this is a  $UL$  factorization, not an  $LU$  factorization, and it takes advantage

of the rank-1 nature of the upper triangle of  $F_n$ . This factorization shows that the inverse  $F_n^{-1} = L^{-1}T_n(-1)$  is lower Hessenberg with integer entries and that  $\det(F_n) = 1$ . Furthermore,  $L$  is totally nonnegative by Theorem 5.2 and  $T_n(-1)^{-1} = M(T_n(-1))^{-1}$  is totally nonnegative by Theorem 5.3, so  $F_n$  is the product of two totally nonnegative matrices and so is totally nonnegative by Theorem 5.1—a property that to our knowledge has not previously been noted. Since  $F_n$  is nonsingular, irreducible (being upper Hessenberg with nonzero subdiagonal), and totally nonnegative it follows from by Theorem 5.6 that  $F_n$  has distinct eigenvalues. The distinctness of the eigenvalues also follows from some rather lengthy analysis of the characteristic polynomial in [38, Thm. 2.5].

Frank discussed two matrices in his paper. The other matrix is obtained from  $A_n = (\min(i, j)) \in \mathbb{R}^{n \times n}$  by taking the rows and columns in reverse order. We will focus on  $A_n$ . For example,

$$A_4 = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 2 & 2 \\ 1 & 2 & 3 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix}.$$

The determinant, the inverse, and the eigenvalues of  $A_n$  can all be easily found by constructing a factorization involving a bidiagonal matrix. Consider subtracting row  $k - 1$  from row  $k$  for  $k = n : -1 : 2$ . For  $A_4$  this yields

$$\begin{bmatrix} 1 & & & \\ -1 & 1 & & \\ & -1 & 1 & \\ & & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 2 & 2 \\ 1 & 2 & 3 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ & 1 & 1 & 1 \\ & & 1 & 1 \\ & & & 1 \end{bmatrix}.$$

In general,  $T_n(-1)^T A_n = U$ , where  $U$  is the upper triangular matrix of 1s. Hence  $A_n = T_n(-1)^{-T} U$ , which is a Cholesky factorization  $A_n = U^T U$  since  $T_n(-1)^{-1} = U$ , which shows that  $A_n$  is symmetric positive definite. Furthermore,  $\det(A) = \det(U)^2 = 1$  and  $A_n^{-1} = U^{-1} U^{-T} = T_n(-1) T_n(-1)^T$ , which is tridiagonal since  $T_n$  is upper bidiagonal. Now  $T_n(-1)^{-1}$  is totally nonnegative, as noted above; hence  $A_n$  is the product of two totally nonnegative matrices and therefore is totally nonnegative. By Theorem 5.6, the eigenvalues of  $A_n$  are distinct. In fact,  $A_n^{-1}$  is the almost-Toeplitz tridiagonal matrix

$$A_n^{-1} = \begin{bmatrix} 2 & -1 & & & \\ -1 & 2 & -1 & & \\ & -1 & \ddots & \ddots & \\ & & \ddots & 2 & -1 \\ & & & -1 & 1 \end{bmatrix},$$

and its eigenvalues are [16], [23, Chap. 7] (and as given by Frank)

$$\mu_k = 2 \left( 1 + \cos \left( \frac{2k\pi}{2n+1} \right) \right), \quad k = 1 : n.$$

The eigenvalues of  $A_n$  are the reciprocals of the  $\mu_k$ .

**8.2. The Kac–Murdock–Szegő Matrix.** The Kac–Murdock–Szegő matrix is the symmetric Toeplitz matrix, depending on a single parameter  $\rho \in \mathbb{R}$ ,

$$(8.2) \quad A_n(\rho) = \begin{bmatrix} 1 & \rho & \rho^2 & \dots & \rho^{n-1} \\ \rho & 1 & \rho & \dots & \rho^{n-2} \\ \rho^2 & \rho & 1 & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & \rho \\ \rho^{n-1} & \rho^{n-2} & \dots & \rho & 1 \end{bmatrix} \in \mathbb{R}^{n \times n}.$$

It was considered by Kac, Murdock, and Szegő [33, p. 784 ff.], who investigated its spectral properties. It arises in the autoregressive AR(1) model in statistics and signal processing.

It is straightforward to verify that  $A_n$  has a factorization  $A_n = LDL^T$  with

$$(8.3) \quad L = T_n(-\rho)^{-T}, \quad D = \text{diag}(1, 1 - \rho^2, 1 - \rho^2, \dots, 1 - \rho^2).$$

This factorization reveals several properties.

(1)  $\det(A_n(\rho)) = (1 - \rho^2)^{n-1}$ .

(2) For  $\rho \neq \pm 1$ ,  $A_n$  is nonsingular and  $A_n(\rho)^{-1} = T_n(-\rho)D^{-1}T_n(-\rho)^T$  is the tridiagonal (but not Toeplitz) matrix

$$(8.4) \quad A_n(\rho)^{-1} = \frac{1}{1 - \rho^2} \begin{bmatrix} 1 & -\rho & & & \\ -\rho & 1 + \rho^2 & -\rho & & \\ & -\rho & 1 + \rho^2 & \ddots & \\ & & \ddots & \ddots & \ddots \\ & & & -\rho & 1 + \rho^2 & -\rho \\ & & & & -\rho & 1 \end{bmatrix}.$$

(3) For  $0 \leq \rho \leq 1$ ,  $T_n(-\rho) = M(T_n(-\rho))$  and so by Theorem 5.3  $M(T_n(-\rho))^{-1} = T_n(-\rho)^{-1} = L^T$  is totally nonnegative, so  $A_n(\rho)$  is the product of three totally nonnegative matrices and is therefore totally nonnegative. For  $0 < \rho < 1$ ,  $A_n(\rho)$  is also nonsingular and irreducible, so the eigenvalues are distinct by Theorem 5.6. Since  $A_n(\rho) = \Sigma A_n(-\rho)\Sigma$  for  $\Sigma$  in (3.4),  $A_n(\rho)$  is similar to  $A_n(-\rho)$  and therefore  $A_n(\rho)$  has distinct eigenvalues for  $0 \neq \rho \in (-1, 1)$ .

**8.3. The Pascal Matrix.** The Pascal matrix  $P_n \in \mathbb{R}^{n \times n}$ , defined in (4.10), contains the rows of Pascal's triangle along the antidiagonals. For example:

$$P_5 = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 & 5 \\ 1 & 3 & 6 & 10 & 15 \\ 1 & 4 & 10 & 20 & 35 \\ 1 & 5 & 15 & 35 & 70 \end{bmatrix}.$$

This matrix is much-studied and most analyses involve the use of combinatorial identities. A number of key properties can be obtained from a factorization of  $P_n$  into a product of bidiagonal matrices.

The key observation is that  $P_n$  can be reduced to upper triangular form by repeatedly subtracting a row from the row below. For  $n = 5$ , with  $L_k(-1)$  denoting the unit lower

bidiagonal matrix with  $-1$ s in subdiagonal elements  $(k+1, k), \dots, (n-1, n)$ ,

$$\begin{aligned}
L_4(-1)L_3(-1)L_2(-1)L_1(-1)P_5 &= \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & -1 & 1 \end{bmatrix} \\
&\times \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & -1 & 1 \end{bmatrix} P_5 \\
&= \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 3 & 4 \\ 0 & 0 & 1 & 3 & 6 \\ 0 & 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} = R.
\end{aligned}$$

In general, we have

$$P_n = L_1(-1)^{-1}L_2(-1)^{-1} \dots L_{n-1}(-1)^{-1}R_n = L_nR_n,$$

where  $L_n$  is unit lower triangular and  $R_n$  is unit upper triangular. By the uniqueness of the LU and Cholesky factorizations of a positive definite matrix we must have  $L_n = R_n^T$ , so  $P_n = R_n^T R_n$ , and it can be shown that  $R_n = L_{n-1}(1)^T L_{n-2}(1)^T \dots L_1(1)^T$ , which contains the binomial coefficients down its columns.

This is the factorization (5.4) in Theorem 5.4: all the parameters are equal to 1.

We can make several deductions.

(1)  $P_n$  is symmetric positive definite.

(2)  $\det(P_n) = 1$ .

(3)  $P_n$  and  $R_n$  are both totally nonnegative, since they are products of bidiagonal matrices  $L_i(1)$ , each of which is totally nonnegative by Theorem 5.2. Hence the eigenvalues of  $P_n$  are distinct by Theorem 5.6.

(4) The matrix  $S_n = \Sigma R_n$  (where  $\Sigma$  is defined in (3.4)) is involutory, that is,  $S_n^2 = I$ . This can be proved with the aid of the bidiagonal factorization but we omit the rather tedious details. Since  $P_n = S_n^T S_n$ , we have  $P_n^{-1} = S_n^{-1} S_n^{-T} = S_n S_n^T = S_n^T P_n S_n^T$ , so  $P_n^{-1}$  is similar to  $P_n$ , which means that the eigenvalues of  $P_n$  occur in reciprocal pairs. It follows, in particular, that  $\|P_n\|_2 = \|P_n^{-1}\|_2$  and so  $\kappa_2(P_n) = \|P_n\|_2^2$ .

It is also interesting to note that, as an instance of Theorem 6.2, the Cholesky factor  $R_n$  is the exponential of a bidiagonal matrix:  $R_n = e^{C_n}$ , where [2], [14]

$$C_n = \begin{bmatrix} 0 & 1 & & & \\ & 0 & 2 & & \\ & & \ddots & \ddots & \\ & & & 0 & n-1 \\ & & & & 0 \end{bmatrix} \in \mathbb{R}^{n \times n}.$$

The matrix  $C_n$  is called the creation matrix in [1], [2] because of its role in generating matrix representations of polynomials and providing simple proofs of identities.

**8.4. Tridiagonal Matrices from Partial Differential Equations.** Consider a linear system  $Ax = b$ , where  $A = D + L + U$  with  $D = \text{diag}(A)$  and  $L$  and  $U$  the strictly lower triangular and strictly upper triangular parts of  $A$ , respectively. The powers of the matrix  $B = -(D + L)^{-1}U$  govern the convergence of the Gauss–Seidel iteration. Note that  $B$  is nonsymmetric and so in general can have complex eigenvalues.

Suppose  $A$  is tridiagonal with negative diagonal elements and nonnegative elements on the superdiagonal and subdiagonal, as is frequently the case in discretizations of partial differential equations, in which  $A$  is typically a Toeplitz matrix. For example,

$$A = \begin{bmatrix} -2 & 1 & & \\ 1 & -2 & 1 & \\ & 1 & -2 & 1 \\ & & 1 & -2 \end{bmatrix} \Rightarrow B = \begin{bmatrix} 0 & 1/2 & 0 & 0 \\ 0 & 1/4 & 1/2 & 0 \\ 0 & 1/8 & 1/4 & 1/2 \\ 0 & 1/16 & 1/8 & 1/4 \end{bmatrix}.$$

The matrix  $(-D - L)^{-1}$  is totally nonnegative by Theorem 5.3, because  $-D - L = M(-D - L)$ , and  $U$  is totally nonnegative by Theorem 5.2. Hence  $B = (-D - L)^{-1}U$  is lower Hessenberg and totally nonnegative. Furthermore,  $B$  is irreducible if the subdiagonal of  $L$  and the superdiagonal of  $U$  are nonzero. Then Theorem 5.6 shows that the eigenvalues of  $B$  are real and nonnegative and the positive eigenvalues are distinct. The eigenvalues of  $B$  can be deduced from the analysis of Young [49], [50, Chap. 5].

**Acknowledgement.** I thank Massimiliano Fasi and Xiaobo Liu for their helpful comments on a draft manuscript.

#### REFERENCES

- [1] Lidia Aceto and Isabel Cação. [A matrix approach to Sheffer polynomials](#). *J. Math. Anal. Appl.*, 446(1): 87–100, 2017.
- [2] Lidia Aceto and Donato Trigiant. [The matrices of Pascal and other greats](#). *Amer. Math. Monthly*, 108(3): 232–245, 2001.
- [3] E. Anderson, Z. Bai, C. H. Bischof, S. Blackford, J. W. Demmel, J. J. Dongarra, J. J. Du Croz, A. Greenbaum, S. J. Hammarling, A. McKenney, and D. C. Sorensen. *LAPACK Users' Guide*. Third edition, Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, 1999. xxvi+407 pp. ISBN 0-89871-447-8.
- [4] Dennis S. Bernstein. *Matrix Mathematics: Theory, Facts, and Formulas*. Second edition, Princeton University Press, Princeton, NJ, USA, 2009. xxxix+1139 pp. ISBN 978-0-691-14039-1.
- [5] Åke Björck. *Numerical Methods for Least Squares Problems*. Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, 1996. xvii+408 pp. ISBN 0-89871-360-9.
- [6] Åke Björck and Victor Pereyra. [Solution of Vandermonde systems of equations](#). *Math. Comp.*, 24(112): 893–903, 1970.
- [7] Samuel D. Conte and Carl de Boor. *Elementary Numerical Analysis: An Algorithmic Approach*. Third edition, McGraw-Hill, Tokyo, 1980. xii+432 pp. ISBN 0-07-066228-2.
- [8] Chandler Davis. [Explicit functional calculus](#). *Linear Algebra Appl.*, 6:193–199, 1973.
- [9] Jorge Delgado, Plamen Koev, Ana Marco, José-Javier Martínez, Juan Manuel Peña, Per-Olof Persson, and Steven Spasov. [Bidiagonal decompositions of Vandermonde-type matrices of arbitrary rank](#). *J. Comput. Appl. Math.*, 426:115064, 2023.
- [10] James W. Demmel and Kahan William. [Accurate singular values of bidiagonal matrices](#). *SIAM J. Sci. Statist. Comput.*, 11(5):873–912, 1990.
- [11] Jean Descloux. [Bounds for the spectral norm of functions of matrices](#). *Numer. Math.*, 5(1):185–190, 1963.
- [12] J. J. Dongarra, J. R. Bunch, C. B. Moler, and G. W. Stewart. *LINPACK Users' Guide*. Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, 1979. ISBN 0-89871-172-X.
- [13] P. J. Eberlein. [A note on the matrices denoted by  \$B\_n\$](#) . *SIAM J. Appl. Math.*, 20(1):87–92, 1971.
- [14] Alan Edelman and Gilbert Strang. [Pascal matrices](#). *Amer. Math. Monthly*, 111(3):189–197, 2004.
- [15] Stanley C. Eisenstat and Ilse C. F. Ipsen. [Relative perturbation techniques for singular value problems](#). *SIAM J. Numer. Anal.*, 32(6):1972–1988, 1995.
- [16] Joseph Frederick Elliott. [The characteristic roots of certain real symmetric matrices](#). M.Sc. Thesis, The University of Tennessee, Knoxville, TN, USA, August 1953. 50 pp.
- [17] Shaun M. Fallat. Totally positive and totally nonnegative matrices. In *Handbook of Linear Algebra*, Leslie Hogben, editor, second edition, Chapman and Hall/CRC, Boca Raton, FL, USA, 2014, pages 29.1–29.17.
- [18] Shaun M. Fallat, Michael I. Gekhtman, and Charles R. Johnson. [Spectral structures of irreducible totally nonnegative matrices](#). *SIAM J. Matrix Anal. Appl.*, 22(2):627–645, 2000.

- [19] Shaun M. Fallat and Charles R. Johnson. *Totally Nonnegative Matrices*. Princeton University Press, Princeton, NJ, USA, 2011. xv+248 pp. ISBN 978-0-691-12157-4.
- [20] Werner L. Frank. Computing eigenvalues of complex matrices by determinant evaluation and by methods of Danilewski and Wielandt. *J. Soc. Indust. Appl. Math.*, 6(4):378–392, 1958.
- [21] Mariano Gasca and Juan M. Peña. On factorizations of totally positive matrices. In *Total Positivity and Its Applications*, Mariano Gasca and Charles A. Micchelli, editors, Springer-Verlag, 1996, pages 109–130.
- [22] G. H. Golub and W. Kahan. Calculating the singular values and pseudo-inverse of a matrix. *SIAM J. Numer. Anal.*, 2(2):205–224, 1965.
- [23] Robert T. Gregory and David L. Karney. *A Collection of Matrices for Testing Computational Algorithms*. Wiley, New York, USA, 1969. ix+154 pp. Reprinted with corrections by Robert E. Krieger, Huntington, New York, 1978. ISBN 0-88275-649-4.
- [24] Desmond J. Higham. Condition numbers and their condition numbers. *Linear Algebra Appl.*, 214:193–213, 1995.
- [25] Nicholas J. Higham. Efficient algorithms for computing the condition number of a tridiagonal matrix. *SIAM J. Sci. Statist. Comput.*, 7(1):150–165, 1986.
- [26] Nicholas J. Higham. Error analysis of the Björck-Pereyra algorithms for solving Vandermonde systems. *Numer. Math.*, 50(5):613–632, 1987.
- [27] Nicholas J. Higham. Stability analysis of algorithms for solving confluent Vandermonde-like systems. *SIAM J. Matrix Anal. Appl.*, 11(1):23–41, 1990.
- [28] Nicholas J. Higham. *Accuracy and Stability of Numerical Algorithms*. Second edition, Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, 2002. xxx+680 pp. ISBN 0-89871-521-0.
- [29] Nicholas J. Higham. *Functions of Matrices: Theory and Computation*. Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, 2008. xx+425 pp. ISBN 978-0-898716-46-7.
- [30] Nicholas J. Higham and Mantas Mikaitis. Anymatrix: An extensible MATLAB matrix collection. *Numer. Algorithms*, 90(3):1175–1196, 2021.
- [31] Leslie Hogben, editor. *Handbook of Linear Algebra*. Second edition, Chapman and Hall/CRC, Boca Raton, FL, USA, 2014. xxix+1904 pp. ISBN 978-1-4665-0728-9.
- [32] Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. Second edition, Cambridge University Press, Cambridge, UK, 2013. xviii+643 pp. ISBN 978-0-521-83940-2.
- [33] M. Kac, W. L. Murdock, and G. Szegő. On the eigen-values of certain Hermitian forms. *Journal of Rational Mechanics and Analysis*, 2:767–800, 1953.
- [34] Plamen Koev. Accurate eigenvalues and SVDs of totally nonnegative matrices. *SIAM J. Matrix Anal. Appl.*, 27(1):1–23, 2005.
- [35] Plamen Koev. Accurate computations with totally nonnegative matrices. *SIAM J. Matrix Anal. Appl.*, 29(3): 731–751, 2007.
- [36] R. B. Lehoucq. The computation of elementary unitary matrices. *ACM Trans. Math. Software*, 22(4):393–400, 1996.
- [37] A. McCurdy, K. C. Ng, and B. N. Parlett. Accurate computation of divided differences of the exponential function. *Math. Comp.*, 43(168):501–528, 1984.
- [38] Efruz Özlem Mersin and Mustafa Bahşi. Sturm theorem for the generalized Frank matrix. *Hacettepe Journal of Mathematics and Statistics*, 50(4):1002–1011, 2021.
- [39] Multiprecision Computing Toolbox. Advanpix, Tokyo. <http://www.advanpix.com>.
- [40] G. Opitz. Steigungsmatrizen. *Z. Angew. Math. Mech.*, 44:T52–T54, 1964.
- [41] Christopher C. Paige and Michael A. Saunders. LSQR: An algorithm for sparse linear equations and sparse least squares. *ACM Trans. Math. Software*, 8(1):43–71, 1982.
- [42] Beresford N. Parlett. *The Symmetric Eigenvalue Problem*. Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, 1998. xxiv+398 pp. Unabridged, amended version of book first published by Prentice-Hall in 1980. ISBN 0-89871-402-8.
- [43] H. Rutishauser. On test matrices. In *Programmation en Mathématiques Numériques, Besançon, 1966*, volume 7 (no. 165) of *Éditions Centre Nat. Recherche Sci., Paris*, 1968, pages 349–365.
- [44] Robert D. Skeel. Scaling for numerical stability in Gaussian elimination. *J. ACM*, 26(3):494–526, 1979.
- [45] Charles F. Van Loan. A study of the matrix exponential. Numerical Analysis Report No. 10, University of Manchester, Manchester, UK, August 1975. Reissued as MIMS EPrint 2006.397, Manchester Institute for Mathematical Sciences, The University of Manchester, UK, November 2006.
- [46] J. M. Varah. A generalization of the Frank matrix. *SIAM J. Sci. Statist. Comput.*, 7(3):835–839, 1986.
- [47] J. H. Wilkinson. Error analysis of floating-point computation. *Numer. Math.*, 2(1):319–340, 1960.
- [48] J. H. Wilkinson. *The Algebraic Eigenvalue Problem*. Oxford University Press, Oxford, UK, 1965. xviii+662 pp. ISBN 0-19-853403-5 (hardback), 0-19-853418-3 (paperback).
- [49] D. M. Young. Iterative methods for solving partial difference equations of elliptic type. *Trans. Amer. Math. Soc.*, 76(1):92–111, 1954.
- [50] David M. Young. *Iterative Solution of Large Linear Systems*. Academic Press, New York, 1971. xxiv+570 pp. ISBN 0-12-773050-8.