

***Universal connection and curvature for statistical  
manifold geometry***

Arwini, Khadiga and Del Riego, L and Dodson, CTJ

2007

MIMS EPrint: **2005.41**

Manchester Institute for Mathematical Sciences  
School of Mathematics

The University of Manchester

Reports available from: <http://eprints.maths.manchester.ac.uk/>

And by contacting: The MIMS Secretary  
School of Mathematics  
The University of Manchester  
Manchester, M13 9PL, UK

ISSN 1749-9097

# Universal connection and curvature for statistical manifold geometry

Khadiga Arwini<sup>1</sup>, L. Del Riego<sup>2</sup> and C.T.J. Dodson<sup>1</sup>

<sup>1</sup>School of Mathematics, University of Manchester, Manchester M60 1QD, UK

arwini2001@yahoo.com    ctdodson@manchester.ac.uk

and

<sup>2</sup>Departamento de Matemáticas, Facultad de Ciencias

Universidad Autónoma de San Luis Potosí, San Luis Potosí, SLP, 78900 México

lilia@ciencias.uaslp.mx

*June 17, 2005*

## Abstract

Statistical manifolds are representations of smooth families of probability density functions that allow differential geometric methods to be applied to problems in stochastic processes, mathematical statistics and information theory. It is common to have to consider a number of linear connections on a given statistical manifold and so it is important to know the corresponding universal connection and curvature; then all linear connections and their curvatures are pullbacks. An important class of statistical manifolds is that arising from the exponential families and one particular family is that of gamma distributions, which we showed recently to have important uniqueness properties in stochastic processes. Here we provide formulae for universal connections and curvatures on exponential families and give an explicit example for the manifold of gamma distributions.

## 1 Introduction

Information geometry is the study of Riemannian geometric properties of statistical manifolds consisting of smooth families of probability density functions. Such manifolds are endowed with the information metric of Rao [22], which arose from the Fisher information matrix [13]. These parts of mathematical statistics have deep relations with general information theory; see eg Roman [23] for a modern account of information theory from a mathematical viewpoint.

For our present purposes we may view a probability density function on  $\Omega \subset \mathbb{R}^m$  as a subadditive measure function of unit weight, namely, a nonnegative map

$$\begin{aligned} f &: \Omega \longrightarrow [0, \infty) \\ \int_{\Omega} f &= 1 \\ \int_{A \cup B} f &\leq \int_A f + \int_B f, \quad \forall A, B \subseteq \Omega. \end{aligned}$$

Usually, a probability density function depends on a set of parameters,  $\theta^1, \theta^2, \dots, \theta^n$  and we say that we have an  $n$ -dimensional family of probability density functions. Let  $\Theta$  be the parameter space of an  $n$ -dimensional smooth such family defined on some fixed event space  $\Omega$

$$\{p_{\theta} | \theta \in \Theta\} \quad \text{with} \quad \int_{\Omega} p_{\theta} = 1 \quad \text{for all } \theta \in \Theta.$$

Then, the derivatives of the log-likelihood function,  $l = \log p_{\theta}$ , yield a matrix with entries

$$g_{ij} = \int_{\Omega} p_{\theta} \left( \frac{\partial l}{\partial \theta^i} \frac{\partial l}{\partial \theta^j} \right) = - \int_{\Omega} p_{\theta} \left( \frac{\partial^2 l}{\partial \theta^i \partial \theta^j} \right),$$

for coordinates  $(\theta^i)$  about  $\theta \in \Theta \subseteq \mathbb{R}^n$ .

This gives rise to a positive definite matrix, so inducing a Riemannian metric  $g$  on  $\Theta$  using for coordinates the parameters  $(\theta^i)$ ; this metric is called the information metric for the family of probability density functions—the second equality here is subject to certain regularity conditions. Amari [1] and Amari and Nagaoka [2] provide modern accounts of the differential geometry that arises from the Fisher information metric.

## 2 Systems of connections and universal objects

The concept of system (or structure) of connections was introduced by Mangiarotti and Modugno [18, 19], they were concerned with finite-dimensional bundle representations of the space of all connections on a fibred manifold. On each system of connections there exists a unique universal connection of which every connection in the family of connections is a pullback. A similar relation holds between the corresponding universal curvature and the curvatures of the connections of the system. This is a different representation of an object similar to that introduced by Narasimhan and Ramanan [20], [21] for  $G$ -bundles, also allowing a proof of Weil's theorem (cf. [16, 14, 8]).

**Definition 2.1** *A **system of connections** on a fibred manifold  $p : E \longrightarrow M$  is a fibred manifold  $p_c : C \longrightarrow M$  together with a first jet-valued fibred morphism*

$$\xi : C \times_M E \longrightarrow JE$$

*over  $M$ , such that each section  $\tilde{\Gamma} : M \longrightarrow C$  determines a unique connection  $\Gamma = \xi \circ (\tilde{\Gamma} \circ p, I_E)$  on  $E$ . Then  $C$  is the **space of connections** of the system.*

In the sequel we are interested in the system of linear connections on a Riemannian manifold. The system of all linear connections is the subject of studies in eg. [16, 14, 17, 8, 9, 10, 7, 12].

**Theorem 2.1** ([18, 19]) *Let  $(C, \xi)$  be a system of connections on a fibred manifold  $p : E \longrightarrow M$ . Then there is a unique connection form  $\Lambda : C \times_M E \longrightarrow J(C \times_M E)$  On the fibred manifold  $\pi_1 : C \times_M E \longrightarrow C$  with the coordinate expression*

$$\Lambda = dx^\lambda \otimes \partial_\lambda + dc^a \otimes \partial_a + \xi_\lambda^i dx^\lambda \otimes \partial_i.$$

*This  $\Lambda$  is called the **universal connection** because it describes all the connections of the system.*

*Explicitly, each  $\tilde{\Gamma} \in \text{Sec}(C/M)$  gives an injection  $(\tilde{\Gamma} \circ p, I_E)$ , of  $E$  into  $C \times E$ , which is a section of  $\pi_1$  and  $\Gamma$  coincides with the restriction of  $\Lambda$  to this section:*

$$\Lambda|_{(\tilde{\Gamma} \circ p, I_E)E} = \Gamma.$$

*A similar relation holds between its curvature  $\Omega$ , called **universal curvature**, and the curvatures of the connections of the system.*

$$\Omega = \frac{1}{2} [\Lambda, \Lambda] = d_\Lambda \Lambda : C \times_M E \longrightarrow \wedge^2(T^*C) \otimes_E V(E).$$

*So the universal curvature  $\Omega$  has the coordinate expression:*

$$\Omega = \frac{1}{2} \left( \xi_\lambda^j \partial_j \xi_\eta^i dx^\lambda \wedge dx^\eta + 2 \partial_a \xi_\eta^i dx^a \wedge dx^\eta \right) \otimes \partial_i.$$

## 3 Exponential family of probability density functions on $\mathbb{R}$

An important class of statistical manifolds is that arising from the so-called exponential family [2] and one case is that of gamma distributions, which we showed recently in [4, 5] to have important uniqueness

properties for near-random stochastic processes. Note also that Hwang and Hu [15] provided an important new characterization of gamma distributions, which helps understanding of their common application in modelling real processes. More details on statistical manifolds in general can be found in [1, 2] and we have provided in [6] explicit geometric neighbourhoods of independence for common bivariate processes. In the present section we shall be concerned with the system of all linear connections on the manifold of an arbitrary exponential family, using the tangent bundle or the frame bundle to give the system space. We provide formulae for the universal connections and curvatures and give an explicit example for the manifold of gamma distributions.

An  $n$ -dimensional set of probability density functions  $S = \{p_\theta | \theta \in \Theta \subset \mathbb{R}^n\}$  for random variable  $x \in \Omega \subseteq \mathbb{R}$  is said to be an **exponential family** [2] when the density functions can be expressed in terms of functions  $\{C, F_1, \dots, F_n\}$  on  $\mathbb{R}$  and a function  $\varphi$  on  $\Theta$  as:

$$p_\theta(x) = e^{\{C(x) + \sum_i (\theta^i F_i(x)) - \varphi(\theta)\}}.$$

Then we say that  $(\theta^i)$  are its **natural** coordinates, and  $\varphi$  is its **potential function**. From the normalization condition  $\int_\Omega p_\theta(x) dx = 1$  we obtain:

$$\varphi(\theta) = \log \int_\Omega e^{\{C(x) + \sum_i (\theta^i F_i(x))\}} dx.$$

From the definition of an exponential family, and putting  $\partial_i = \frac{\partial}{\partial \theta^i}$ , we use the log-likelihood function  $l(\theta, x) = \log(p_\theta(x))$  to obtain

$$\partial_i l(\theta, x) = F_i(x) - \partial_i \varphi(\theta)$$

and

$$\partial_i \partial_j l(\theta, x) = -\partial_i \partial_j \varphi(\theta).$$

The Fisher information metric  $g$  [1, 2] on the  $n$ -dimensional space of parameters  $\Theta \subset \mathbb{R}^n$ , equivalently on the set  $S = \{p_\theta | \theta \in \Theta \subset \mathbb{R}^n\}$ , has coordinates:

$$[g_{ij}] = - \int_\Omega [\partial_i \partial_j l(\theta, x)] p_\theta(x) dx = \partial_i \partial_j \varphi(\theta) = \varphi_{ij}(\theta).$$

Then,  $(S, g)$  is a Riemannian  $n$ -manifold with Levi-Civita connection given by:

$$\begin{aligned} \Gamma_{ij}^k(\theta) &= \sum_{h=1}^n \frac{1}{2} g^{kh} (\partial_i g_{jh} + \partial_j g_{ih} - \partial_h g_{ij}) \\ &= \sum_{h=1}^n \frac{1}{2} g^{kh} \partial_i \partial_j \partial_h \varphi(\theta) = \sum_{h=1}^n \frac{1}{2} \varphi^{kh}(\theta) \varphi_{ijh}(\theta) \end{aligned}$$

where  $[\varphi^{hk}(\theta)]$  represents the inverse to  $[\varphi_{hk}(\theta)]$ .

Next we obtain a family of symmetric connections which includes the Levi-Civita case and has significance in mathematical statistics. Consider for  $\alpha \in \mathbb{R}$  the function  $\Gamma_{ij,k}^{(\alpha)}$  which maps each point  $\theta \in \Theta$  to the following value:

$$\begin{aligned} \Gamma_{ij,k}^{(\alpha)}(\theta) &= \int_\Omega \left( \partial_i \partial_j l + \frac{1-\alpha}{2} \partial_i l \partial_j l \right) \partial_k l p_\theta \\ &= \frac{1-\alpha}{2} \partial_i \partial_j \partial_k \varphi(\theta) = \frac{1-\alpha}{2} \varphi_{ijk}(\theta). \end{aligned}$$

So we have an affine connection  $\nabla^{(\alpha)}$  on the statistical manifold  $(S, g)$  defined by

$$g(\nabla_{\partial_i}^{(\alpha)} \partial_j, \partial_k) = \Gamma_{ij,k}^{(\alpha)},$$

where  $g$  is the Fisher information metric. We call this  $\nabla^{(\alpha)}$  the  $\alpha$ -connection and it is clearly a symmetric connection and defines an  $\alpha$ -curvature. We have also

$$\begin{aligned} \nabla^{(\alpha)} &= (1-\alpha) \nabla^{(0)} + \alpha \nabla^{(1)}, \\ &= \frac{1+\alpha}{2} \nabla^{(1)} + \frac{1-\alpha}{2} \nabla^{(-1)}. \end{aligned}$$




Figure 1: Affine immersion using natural coordinates  $(\mu, \nu)$  in  $\mathbb{R}^3$  for the gamma 2-manifold  $(\mathcal{G}, g)$ . The surface is shaded by the Gaussian curvature  $K_{\mathcal{G}}$  which is independent of  $\mu$  and monotonically decreases from  $-\frac{1}{4}$  to almost  $-\frac{1}{2}$  as  $\nu$  increases to 2.

For a submanifold  $M \subset S$ , the  $\alpha$ -connection on  $M$  is simply the restriction with respect to  $g$  of the  $\alpha$ -connection on  $S$ . Note that the 0-connection is the Riemannian or Levi-Civita connection with respect to the Fisher metric and its uniqueness implies that an  $\alpha$ -connection is a metric connection if and only if  $\alpha = 0$ .

### 3.1 Example: Gamma 2-manifold $(\mathcal{G}, g)$

Gamma distributions form an exponential family with probability density functions:

$$\mathcal{G} = \{p(x; \mu, \nu) = \mu^\nu \frac{x^{\nu-1}}{\Gamma(\nu)} e^{-x\mu} \quad \text{for } \mu, \nu \in \mathbb{R}^+\}. \quad (3.1)$$

The gamma distributions are very important in, information theory, mathematical statistics and stochastic processes. This is because they contain as a special case,  $\nu = 1$ , the negative exponential distribution that represents random states, ie Poisson processes, and because of certain uniqueness properties [5, 15]. It turns out that  $(\theta^i) = (\mu, \nu)$  is a natural coordinate system with corresponding potential function

$$\varphi(\mu, \nu) = \log \Gamma(\nu) - \nu \log \mu. \quad (3.2)$$

The information metric  $g$  has arc length function

$$ds_g^2 = \frac{\nu}{\mu^2} d\mu^2 - \frac{2}{\mu} d\mu d\nu + \psi'(\nu) d\nu^2 \quad \text{for } \mu, \nu \in \mathbb{R}^+$$

where  $\psi(\nu) = \frac{\Gamma'(\nu)}{\Gamma(\nu)}$  is the digamma function. The independent nonzero Levi-Civita connection compo-

| Univariate density | Coordinates     | Mean  | Variance    | $R^{(\alpha)}$   |
|--------------------|-----------------|-------|-------------|--|
| Gaussian           | $(\mu, \sigma)$ | $\mu$ | $\sigma$    | $(\alpha^2 - 1)$   |
| Gamma              | $(\mu, \nu)$    | $\mu$ | $\mu^2/\nu$ | $\frac{(1-\alpha^2)(\psi'(\nu)+\nu\psi''(\nu))}{2(\nu\psi'(\nu)-1)^2}$ |
| Exponential        | $\mu$           | $\mu$ | $\mu^2$     | 0  |

Table 1:  $\alpha$ -Scalar curvature  $R^{(\alpha)}$  of the univariate Gaussian, gamma and exponential statistical manifolds; the logarithmic derivative of the gamma function is denoted by  $\psi = \Gamma'/\Gamma$ . The case  $\alpha = 0$  corresponds to the Levi-Civita connection.

nents with respect to the natural coordinates  $(\mu, \nu)$  are:

$$\begin{aligned}
\Gamma_{11}^1 &= \frac{(1 - 2\nu\psi'(\nu))}{2\mu(-1 + \nu\psi'(\nu))}, \\
\Gamma_{12}^1 &= \frac{\psi'(\nu)}{2(\nu\psi'(\nu) - 1)}, \\
\Gamma_{22}^1 &= \frac{\mu\psi''(\nu)}{2(\nu\psi'(\nu) - 1)}, \\
\Gamma_{11}^2 &= \frac{\nu}{2\mu^2(1 - \nu\psi'(\nu))}, \\
\Gamma_{12}^2 &= \frac{1}{2\mu(\nu\psi'(\nu) - 1)}, \\
\Gamma_{22}^2 &= \frac{\nu\psi''(\nu)}{2(\nu\psi'(\nu) - 1)}.
\end{aligned}$$

The Riemannian 2-manifold  $(\mathcal{G}, g)$  has been shown by Dodson and Matsuzoe [11] to admit an affine immersion in  $\mathbb{R}^3$ . This is depicted in Figure 1, shaded by the Gaussian curvature  $K_{\mathcal{G}}$ , which is independent of  $\mu$  and, with increasing  $\nu$ ,  $K_{\mathcal{G}}$  monotonically decreases from  $-\frac{1}{4}$  to  $-\frac{1}{2}$ :

$$K_{\mathcal{G}} = \frac{\psi'(\nu) + \nu\psi''(\nu)}{4(\nu\psi'(\nu) - 1)^2}.$$

To compute the  $\alpha$ -connection components it is convenient here to change to the orthogonal coordinates  $(\beta = \nu/\mu, \nu)$  for which the metric components are given by

$$ds^2 = \frac{\nu}{\beta^2} d\beta^2 + \left( \psi'(\nu) - \frac{1}{\nu} \right) d\nu^2 \quad \text{for } \beta, \nu \in \mathbb{R}^+.$$

**Proposition 3.1 (Arwini [3])** *The independent nonzero components,  $\Gamma_{jk}^{(\alpha)i}$ , of  $\nabla^{(\alpha)}$  are*

$$\begin{aligned}
\Gamma_{11}^{(\alpha)1} &= -\frac{\alpha + 1}{\beta}, \\
\Gamma_{12}^{(\alpha)1} &= \frac{\alpha + 1}{2\nu}, \\
\Gamma_{11}^{(\alpha)2} &= \frac{(\alpha - 1)\nu}{2\beta^2(\nu\psi'(\nu) - 1)}, \\
\Gamma_{22}^{(\alpha)2} &= \frac{(1 - \alpha)(1 + \nu^2\psi''(\nu))}{2\nu(\nu\psi'(\nu) - 1)}.
\end{aligned}$$

□

**Corollary 3.1** *The Levi Civita connection of  $(\mathcal{G}, g)$  is recovered by  $\Gamma_{jk}^{(0)i}$  in  $(\beta, \nu)$  coordinates and then the curves  $\nu = \text{constant}$  are geodesics.*

□

| Bivariate density                 | Coordinates               | Covariance     | $R^{(\alpha)}$       |
|-----------------------------------|---------------------------|----------------|----------------------|
| <b>McKay</b> $M$                  | $(\alpha_1, c, \alpha_2)$ | $\alpha_1/c^2$ | $R_M^{(\alpha)}$     |
| $M_1 \subset M$ : $\alpha_1 = 1$  | $(c, \alpha_2)$           | $1/c^2$        | $R_{M_1}^{(\alpha)}$ |
| $M_2 \subset M$ : $\alpha_2 = 1$  | $(\alpha_1, c)$           | $\alpha_1/c^2$ | $R_{M_2}^{(\alpha)}$ |
| $M_3$ : $\alpha_1 + \alpha_2 = 1$ | $(\alpha_1, c)$           | $\alpha_1/c^2$ | 0                    |

Table 2:  $\alpha$ -Scalar curvature  $R^{(\alpha)}$  of the McKay bivariate gamma manifold; see § 3.2 for the formulae  $R_M^{(\alpha)}$ ,  $R_{M_1}^{(\alpha)}$ ,  $R_{M_2}^{(\alpha)}$ . The case  $\alpha = 0$  corresponds to the Levi-Civita connection.

| Bivariate density                                | Coordinates                              | Covariance  | $R^{(\alpha)}$               |
|--|--|---|------------------------------|
| <b>Freund</b> $F$                                | $(\alpha_1, \beta_1, \alpha_2, \beta_2)$ | $\frac{\beta_1 \beta_2 - \alpha_1 \alpha_2}{\beta_1 \beta_2 (\alpha_1 + \alpha_2)^2}$ | $\frac{-3(\alpha^2 - 1)}{2}$ |
| $F_1 \subset F$ : $\beta_i = \alpha_i$           | $(\alpha_1, \alpha_2)$                   | 0   | 0                            |
| $F_2$ : $\alpha_1 = \alpha_2, \beta_1 = \beta_2$ | $(\alpha_1, \beta_1)$                    | $\frac{1}{4} \left( \frac{1}{\alpha_1^2} - \frac{1}{\beta_1^2} \right)$               | 0                            |
| $F_3$ : $\beta_i = \alpha_1 + \alpha_2$          | $(\alpha_1, \alpha_2, \beta_2)$          | $\frac{\alpha_1^2 + \alpha_1 \alpha_2 + \alpha_2^2}{(\alpha_1 + \alpha_2)^4}$         | 0                            |

Table 3:  $\alpha$ -Scalar curvature  $R^{(\alpha)}$  of the Freund bivariate exponential manifold. The case  $\alpha = 0$  corresponds to the Levi-Civita connection.

| Bivariate density                     | Coordinates                                       | Covariance    | $R^{(\alpha)}$            |
|---------------------------------------|---|---------------|---------------------------|
| <b>Gaussian</b> $N$                   | $(\mu_1, \mu_2, \sigma_1, \sigma_{12}, \sigma_2)$ | $\sigma_{12}$ | $\frac{9(\alpha^2-1)}{2}$ |
| $N_1 \subset N: \sigma_{12} = 0$      | $(\mu_1, \mu_2, \sigma_1, \sigma_2)$              | 0             | $2(\alpha^2 - 1)$         |
| $N_2: \sigma_i = \sigma, \mu_i = \mu$ | $(\mu, \sigma, \sigma_{12})$                      | $\sigma_{12}$ | $(\alpha^2 - 1)$          |
| $N_3: \mu_1 = \mu_2 = 0$              | $(\sigma_1, \sigma_2, \sigma_{12})$               | $\sigma_{12}$ | $2(\alpha^2 - 1)$         |

Table 4:  $\alpha$ -Scalar curvature  $R^{(\alpha)}$  of the bivariate Gaussian manifold. The case  $\alpha = 0$  corresponds to the Levi-Civita connection.

### 3.2 $\alpha$ -Scalar curvature for common distributions

For convenience of reference we summarize curvature results in Table 1 for univariate Gaussian, gamma and exponential distributions and in Tables 2,3,4 respectively for bivariate gamma, exponential and Gaussian distributions, from recent work of Arwini and Dodson [4, 5, 6]. We have used *Mathematica* for many calculations and we can make available the associated interactive Notebooks. The  $\alpha$ -scalar curvature for the McKay bivariate gamma manifold  $M$  and its submanifolds  $M_1, M_2$ , have long expressions so we give them here:

$$\begin{aligned}
R_M^{(\alpha)} &= (1 - \alpha^2) \left( \frac{\psi'(\alpha_2) (\psi'(\alpha_1) (\psi'(\alpha_1) + \psi'(\alpha_2)) - 2\psi''(\alpha_1)) - 2\psi'(\alpha_1)\psi''(\alpha_2)}{2(\psi'(\alpha_1) + \psi'(\alpha_2) - \psi'(\alpha_1)\psi'(\alpha_2)(\alpha_1 + \alpha_2))^2} \right. \\
&\quad \left. + \frac{(\psi'(\alpha_2)^2\psi''(\alpha_1) + (\psi'(\alpha_1)^2 - \psi''(\alpha_1))\psi''(\alpha_2))(\alpha_1 + \alpha_2)}{2(\psi'(\alpha_1) + \psi'(\alpha_2) - \psi'(\alpha_1)\psi'(\alpha_2)(\alpha_1 + \alpha_2))^2} \right), \\
\psi(\alpha_i) &= \frac{\Gamma'(\alpha_i)}{\Gamma(\alpha_i)}. \\
R_{M_1}^{(\alpha)} &= \frac{(1 - \alpha^2) (\psi'(\alpha_2) + \psi''(\alpha_2) (1 + \alpha_2))}{2(\psi'(\alpha_2) (1 + \alpha_2) - 1)^2} \\
R_{M_2}^{(\alpha)} &= \frac{(1 - \alpha^2) (\psi'(\alpha_1) + \psi''(\alpha_1) (1 + \alpha_1))}{2(\psi'(\alpha_1) (1 + \alpha_1) - 1)^2}.
\end{aligned}$$

## 4 Systems of linear connections

### 4.1 Tangent bundle system: $C_T \times TM \longrightarrow JTM$

The system of all linear connections on a manifold  $M$  has a representation on the tangent bundle

$$E = TM \longrightarrow M$$

with system space

$$C_T = \{\alpha \otimes j\gamma \in T^*M \otimes_M JTM \mid j\gamma : TM \longrightarrow TTM \text{ projects onto } I_{TM}\}$$

Here we view  $I_{TM}$  as a section of  $T^*M \otimes TM$ , which is a subbundle of  $T^*M \otimes TTM$ , with local expression  $dx^\lambda \otimes \partial_\lambda$ .



The fibred morphism for this system is given by

$$\begin{aligned}\xi_T : C_T \times_M TM &\longrightarrow JTM \subset T^*M \otimes_{TM} TTM, \\ (\alpha \otimes j\gamma, \nu) &\longmapsto \alpha(\nu)j\gamma.\end{aligned}$$

In coordinates  $(x^\lambda)$  on  $M$  and  $(y^\lambda)$  on  $TM$

$$\begin{aligned}\xi_T &= dx^\lambda \otimes (\partial_\lambda - \gamma_\lambda^i \partial_i) \\ &= dx^\lambda \otimes (\partial_\lambda - y^j \Gamma_{j\lambda}^i \partial_i) \\ &= dx^\lambda \otimes (\partial_\lambda - y^j (\sum_{h=1}^n \frac{1}{2} \varphi^{jh} \varphi_{j\lambda h}) \partial_i)\end{aligned}$$

Each section of  $C_T \longrightarrow M$ , such as  $\tilde{\Gamma} : M \longrightarrow C_T : (x^\lambda) \longrightarrow (x^\lambda, \gamma_{\eta\theta})$ ; determines the unique linear connection  $\Gamma = \xi_T \circ (\tilde{\Gamma} \circ \pi_T, I_{TM})$  with Christoffel symbols  $\Gamma_{\eta\theta}^\lambda$ .

On the fibred manifold  $\pi_1 : C_T \times_M TM \longrightarrow C_T$ ; the universal connection is given by:

$$\begin{aligned}\Lambda_T : C_T \times_M TM &\longrightarrow J(C_T \times_M TM) \subset T^*C_T \otimes T(C_T \times_M TM), \\ (x^\lambda, v_{\eta\kappa}^\lambda, y^\lambda) &\longmapsto [(X^\lambda, V_{\eta\kappa}^\lambda) \longrightarrow (X^\lambda, V_{\eta\kappa}^\lambda, Y^\eta V_{\eta\kappa}^\lambda X^\kappa)].\end{aligned}$$

briefly,

$$\begin{aligned}\Lambda_T &= dx^\lambda \otimes \partial_\lambda + dv^a \otimes \partial_a + y^\eta v_{\eta\kappa}^i dx^\kappa \otimes \partial_i \\ &= dx^\lambda \otimes \partial_\lambda + dv^a \otimes \partial_a + y^\eta (\sum_{h=1}^n \frac{1}{2} \varphi^{ih} \varphi_{\eta\kappa h}) dx^\kappa \otimes \partial_i.\end{aligned}$$

Explicitly, each  $\tilde{\Gamma} \in \text{Sec}(C_T/M)$  gives an injection  $(\tilde{\Gamma} \circ \pi_T, I_{TM})$ , of  $TM$  into  $C_T \times TM$ , which is a section of  $\pi_1$  and  $\Gamma$  coincides with the restriction of  $\Lambda_T$  to this section:

$$\Lambda_T|_{(\tilde{\Gamma} \circ \pi_T, I_{TM})TM} = \Gamma.$$

and the universal curvature of the connection  $\Lambda$  is given by:

$$\Omega_T = d_{\Lambda_T} \Lambda_T : C_T \times_M TM \longrightarrow \wedge^2(T^*C_T) \otimes_{TM} V(TM).$$

So here the universal curvature  $\Omega_T$  has the coordinate expression:

$$\begin{aligned}\Omega_T &= \frac{1}{2} \left( y^k v_{k\lambda}^j \partial_j y^m v_{m\eta}^i dx^\lambda \wedge dx^\eta + 2 \partial_a y^m v_{m\eta}^i dx^a \wedge dx^\eta \right) \otimes \partial_i \\ &= \frac{1}{2} \left( y^k (\sum_{h=1}^n \frac{1}{2} \varphi^{jh} \varphi_{k\lambda h}) \partial_j y^m (\sum_{h=1}^n \frac{1}{2} \varphi^{ih} \varphi_{m\eta h}) dx^\lambda \wedge dx^\eta \right) \otimes \partial_i \\ &\quad + \left( \partial_a y^m (\sum_{h=1}^n \frac{1}{2} \varphi^{ih} \varphi_{m\eta h}) dx^a \wedge dx^\eta \right) \otimes \partial_i.\end{aligned}$$

## 4.2 Frame bundle system: $C_F \times FM \longrightarrow JFM$

A linear connection is also a principal (i.e. group invariant) connection on the principal bundle of frames  $FM$  with:

$$E = FM \longrightarrow M = FM/G$$

consisting of linear frames (ordered bases for tangent spaces) with structure group the general linear group,  $G = Gl(n)$ . Here the system space is

$$C_F = JFM/G \subset T^*M \otimes_{TM} TFM/G,$$

consisting of  $G$ -invariant jets. The system morphism is

$$\begin{aligned} \xi_F : C_F \times FM &\longrightarrow JFM \subset T^*M \otimes_{TM} TFM, \\ ([js_x], b) &\longmapsto [T_x M \longmapsto T_b FM]. \end{aligned}$$

In coordinates

$$\begin{aligned} \xi_F &= dx^\lambda \otimes (\partial_\lambda - X^\eta \partial_\eta s_\kappa^\lambda) \tilde{\partial}_{\kappa^\lambda} \\ &= dx^\lambda \otimes (\partial_\lambda - X^\eta \Gamma_{\eta\kappa}^\lambda) \tilde{\partial}_{\kappa^\lambda} \\ &= dx^\lambda \otimes (\partial_\lambda - X^\eta \sum_{h=1}^n \frac{1}{2} \varphi^{\lambda h} \varphi_{\eta\kappa h}) \tilde{\partial}_{\kappa^\lambda} \end{aligned}$$

where  $\tilde{\partial}_{\kappa^\lambda} = \frac{\partial}{\partial b_\kappa^\lambda}$  is the natural base on the vertical fibre of  $T_b FM$  induced by coordinates  $(b_\kappa^\lambda)$  on  $FM$ . Each section of  $C_F \rightarrow M$  that is projectable onto  $I_{TM}$ , such as,  $\hat{\Gamma} : M \rightarrow C_F : (x^\lambda) \rightarrow (x^\lambda, [j\gamma_x])$  with  $\Gamma_{\eta\kappa}^\lambda = \partial_\eta s_\kappa^\lambda$ ; determines the unique linear connection  $\Gamma = \xi_F \circ (\hat{\Gamma} \circ \pi_F, I_{FM})$  with Christoffel symbols  $\Gamma_{\eta\kappa}^\lambda$ . On the principal  $G$ -bundle  $\pi_1 : C_F \times_M FM \rightarrow C_F$  the universal connection is given by:

$$\begin{aligned} \Lambda_F : C_F \times_M FM &\longrightarrow J(C_F \times_M FM) \subset T^*C_F \otimes_{FM} T(C_F \times_{FM} FM), \\ (x^\lambda, v_{\eta\kappa}^\lambda, b_\kappa^\eta) &\longmapsto [(X^\lambda, Y_{\eta\kappa}^\lambda) \longrightarrow (X^\lambda, Y_{\eta\kappa}^\lambda, b_\kappa^\eta v_{\eta\theta}^\lambda X^\theta)]. \end{aligned}$$

Briefly,

$$\begin{aligned} \Lambda_F &= dx^\lambda \otimes \partial_\lambda + dv^a \otimes \partial_a + b_\kappa^\eta v_{\eta\theta}^\lambda dx^\theta \otimes \tilde{\partial}_{\kappa^\lambda} \\ &= dx^\lambda \otimes \partial_\lambda + dv^a \otimes \partial_a + b_\kappa^\eta \left( \sum_{h=1}^n \frac{1}{2} \varphi^{\lambda h} \varphi_{\eta\theta h} \right) dx^\theta \otimes \tilde{\partial}_{\kappa^\lambda}. \end{aligned}$$

Explicitly, each  $\tilde{\Gamma} \in \text{Sec}(C_F/M)$  gives an injection  $(\tilde{\Gamma} \circ \pi_F, I_{FM})$ , of  $FM$  into  $C_F \times FM$ , which is a section of  $\pi_1$  and  $\Gamma$  coincides with the restriction of  $\Lambda_F$  to this section:

$$\Lambda_F|_{(\tilde{\Gamma} \circ \pi_F, I_{FM})FM} = \Gamma$$

and the universal curvature of the connection  $\Lambda$  is given by:

$$\Omega = d\Lambda_F \Lambda_F : C_F \times_M FM \longrightarrow \wedge^2(T^*C_F) \otimes_{FM} V(FM).$$

So here the universal curvature form  $\Omega_F$  has the coordinate expression:

$$\begin{aligned} \Omega_F &= \frac{1}{2} \left( b_\kappa^k v_{k\lambda}^\beta \tilde{\partial}_{\kappa^\beta} b_\omega^m v_{m\eta}^\alpha dx^\lambda \wedge dx^\eta + 2 \partial_a b_\omega^m v_{m\eta}^\alpha dx^a \wedge dx^\eta \right) \otimes \tilde{\partial}_{\omega^\alpha} \\ &= \frac{1}{2} \left( b_\kappa^k \left( \sum_{h=1}^n \frac{1}{2} \varphi^{\beta h} \varphi_{k\lambda h} \right) \tilde{\partial}_{\kappa^\beta} b_\omega^m \left( \sum_{h=1}^n \frac{1}{2} \varphi^{\alpha h} \varphi_{m\eta h} \right) dx^\lambda \wedge dx^\eta \right) \otimes \tilde{\partial}_{\omega^\alpha} \\ &\quad + \left( \partial_a b_\omega^m \left( \sum_{h=1}^n \frac{1}{2} \varphi^{\alpha h} \varphi_{m\eta h} \right) dx^a \wedge dx^\eta \right) \otimes \tilde{\partial}_{\omega^\alpha}. \end{aligned}$$

## 5 Universal connection and curvature on the gamma manifold

For the gamma 2-manifold  $(\mathcal{G}, g)$  we give explicit forms for the system space and its universal connection and curvature.

## 5.1 Tangent bundle system on $(\mathcal{G}, g)$

The system space is

$$C_T = \{\alpha \otimes j\gamma \in T^*\mathcal{G} \otimes_{\mathcal{G}} JT\mathcal{G} \mid j\gamma : T\mathcal{G} \longrightarrow TT\mathcal{G} \text{ projects onto } I_{T\mathcal{G}}\}$$

and the system morphism is

$$\begin{aligned} \xi_T &= dx^\lambda \otimes (\partial_\lambda - y^j \Gamma_{j\lambda}^i \partial_i) \\ &= dx^1 \otimes \left( \partial_1 - \left( \frac{(1 - 2\nu \psi'(\nu))}{2\mu(-1 + \nu \psi'(\nu))} y^1 + \frac{\psi'(\nu)}{2(\nu \psi'(\nu) - 1)} y^2 \right) \partial_1 \right) \\ &\quad + dx^1 \otimes \left( \partial_1 - \left( \frac{\nu}{2\mu^2(1 - \nu \psi'(\nu))} y^1 + \frac{1}{2\mu(\nu \psi'(\nu) - 1)} y^2 \right) \partial_2 \right) \\ &\quad + dx^2 \otimes \left( \frac{\psi'(\nu)}{2(\nu \psi'(\nu) - 1)} y^1 + \frac{\mu \psi''(\nu)}{2(\nu \psi'(\nu) - 1)} y^2 \right) \partial_1 \\ &\quad + dx^2 \otimes \left( \partial_2 - \left( \frac{1}{2\mu(\nu \psi'(\nu) - 1)} y^1 + \frac{\nu \psi''(\nu)}{2(\nu \psi'(\nu) - 1)} y^2 \right) \partial_2 \right). \end{aligned}$$

The universal connection on the gamma manifold is given by:

$$\begin{aligned} \Lambda_T &= dx^\lambda \otimes \partial_\lambda + d_{\lambda j}^i \otimes \partial_{\lambda j}^i + y^\tau \Gamma_{\tau\kappa}^i dx^\kappa \otimes \partial_i \\ &= dx^\lambda \otimes \partial_\lambda + d_{\lambda j}^i \otimes \partial_{\lambda j}^i \\ &\quad + \left( \frac{(1 - 2\nu \psi'(\nu))}{2\mu(-1 + \nu \psi'(\nu))} y^1 + \frac{\psi'(\nu)}{2(\nu \psi'(\nu) - 1)} y^2 \right) dx^1 \otimes \partial_1 \\ &\quad + \left( \frac{\nu}{2\mu^2(1 - \nu \psi'(\nu))} y^1 + \frac{1}{2\mu(\nu \psi'(\nu) - 1)} y^2 \right) dx^1 \otimes \partial_2 \\ &\quad + \left( \frac{\psi'(\nu)}{2(\nu \psi'(\nu) - 1)} y^1 + \frac{\mu \psi''(\nu)}{2(\nu \psi'(\nu) - 1)} y^2 \right) dx^2 \otimes \partial_1 \\ &\quad + \left( \frac{1}{2\mu(\nu \psi'(\nu) - 1)} y^1 + \frac{\nu \psi''(\nu)}{2(\nu \psi'(\nu) - 1)} y^2 \right) dx^2 \otimes \partial_2. \end{aligned}$$

The universal curvature on the gamma manifold is:

$$\Omega_T = \frac{1}{2} \left( y^k \Gamma_{k\lambda}^j \partial_j y^m \Gamma_{m\kappa}^i dx^\lambda \wedge dx^\kappa + 2 \partial_a y^m \Gamma_{m\kappa}^i dx^a \wedge dx^\kappa \right) \otimes \partial_i \quad (i = 1, 2).$$

The analytic form of this is known [3] but is omitted here.

## 5.2 Frame bundle system on $(\mathcal{G}, g)$

The system space is  $C_F = JF\mathcal{G}/G$  and the system morphism is

$$\begin{aligned} \xi_F &= dx^\lambda \otimes (\partial_\lambda - X^\tau \Gamma_{\tau\kappa}^\lambda \tilde{\partial}_{\kappa\lambda}) \\ &= dx^1 \otimes \left( \partial_1 - \left( \frac{(1 - 2\nu \psi'(\nu))}{2\mu(-1 + \nu \psi'(\nu))} X^1 + \frac{\psi'(\nu)}{2(\nu \psi'(\nu) - 1)} X^2 \right) \tilde{\partial}_{1^1} \right) \\ &\quad + dx^2 \otimes \left( \partial_2 - \left( \frac{\nu}{2\mu^2(1 - \nu \psi'(\nu))} X^1 + \frac{1}{2\mu(\nu \psi'(\nu) - 1)} X^2 \right) \tilde{\partial}_{1^2} \right) \\ &\quad + dx^1 \otimes \left( \partial_1 - \left( \frac{\psi'(\nu)}{2(\nu \psi'(\nu) - 1)} X^1 + \frac{\mu \psi''(\nu)}{2(\nu \psi'(\nu) - 1)} X^2 \right) \tilde{\partial}_{2^1} \right) \\ &\quad + dx^2 \otimes \left( \partial_2 - \left( \frac{1}{2\mu(\nu \psi'(\nu) - 1)} X^1 + \frac{\nu \psi''(\nu)}{2(\nu \psi'(\nu) - 1)} X^2 \right) \tilde{\partial}_{2^2} \right). \end{aligned}$$

The universal connection on the gamma manifold is:

$$\begin{aligned}
\Lambda_F &= dx^\lambda \otimes \partial_\lambda + d_{\lambda_j}^i \otimes \partial_{\lambda_j}^i + b_\tau^\kappa \Gamma_{\tau\theta}^\lambda dx^\theta \otimes \tilde{\partial}_{\kappa\lambda} \\
&= dx^\lambda \otimes \partial_\lambda + d_{\lambda_j}^i \otimes \partial_{\lambda_j}^i \\
&+ \left( \frac{(1-2\nu\psi'(\nu))}{2\mu(-1+\nu\psi'(\nu))} b_1^1 + \frac{\psi'(\nu)}{2(\nu\psi'(\nu)-1)} b_1^2 \right) dx^1 \otimes \tilde{\partial}_{1^1} \\
&+ \left( \frac{\psi'(\nu)}{2(\nu\psi'(\nu)-1)} b_1^1 + \frac{\mu\psi''(\nu)}{2(\nu\psi'(\nu)-1)} b_1^2 \right) dx^2 \otimes \tilde{\partial}_{1^1} \\
&+ \left( \frac{(1-2\nu\psi'(\nu))}{2\mu(-1+\nu\psi'(\nu))} b_2^1 + \frac{\psi'(\nu)}{2(\nu\psi'(\nu)-1)} b_2^2 \right) dx^1 \otimes \tilde{\partial}_{2^1} \\
&+ \left( \frac{\psi'(\nu)}{2(\nu\psi'(\nu)-1)} b_2^1 + \frac{\mu\psi''(\nu)}{2(\nu\psi'(\nu)-1)} b_2^2 \right) dx^2 \otimes \tilde{\partial}_{2^1} \\
&+ \left( \frac{\nu}{2\mu^2(1-\nu\psi'(\nu))} b_1^1 + \frac{1}{2\mu(\nu\psi'(\nu)-1)} b_1^2 \right) dx^1 \otimes \tilde{\partial}_{1^2} \\
&+ \left( \frac{1}{2\mu(\nu\psi'(\nu)-1)} b_1^1 + \frac{\nu\psi''(\nu)}{2(\nu\psi'(\nu)-1)} b_1^2 \right) dx^2 \otimes \tilde{\partial}_{1^2} \\
&+ \left( \frac{\nu}{2\mu^2(1-\nu\psi'(\nu))} b_2^1 + \frac{1}{2\mu(\nu\psi'(\nu)-1)} b_2^2 \right) dx^1 \otimes \tilde{\partial}_{2^2} \\
&+ \left( \frac{1}{2\mu(\nu\psi'(\nu)-1)} b_2^1 + \frac{\nu\psi''(\nu)}{2(\nu\psi'(\nu)-1)} b_2^2 \right) dx^2 \otimes \tilde{\partial}_{2^2}.
\end{aligned}$$

The universal curvature on the gamma manifold is:

$$\Omega_F = \frac{1}{2} \left( b_\kappa^k \Gamma_{k\lambda}^\eta \tilde{\partial}_{\kappa\eta} b_\omega^m \Gamma_{m\kappa}^\alpha dx^\lambda \wedge dx^\kappa + 2 \partial_a b_\omega^m \Gamma_{m\kappa}^\alpha dx^a \wedge dx^\kappa \right) \otimes \tilde{\partial}_{\omega\alpha}.$$

The analytic form of this is known [3] but is omitted here.

## 6 Universal connection and curvature for $\alpha$ -connections [10]

Consider an exponential family having statistical  $n$ -manifold  $(M, g)$  and the system

$$C \times FM \longrightarrow JFM : (\alpha, b) \mapsto \Gamma^{(\alpha)}(b)$$

where  $C = M \times \mathbb{R}$  is the direct product manifold of  $M$  with the standard real line. So the system space  $C$  consists of a stack of copies of  $M$ . Then every  $\tilde{\Gamma} \in \text{Sec}(C/M)$  is a constant real function on  $M$ , so defining precisely one  $\alpha$ -connection.

In the case of the frame bundle system,  $(M \times \mathbb{R}) \times_M FM \longrightarrow JFM$ , the universal connection on the system of  $\alpha$ -connections is

$$\begin{aligned}
\Lambda &: (M \times \mathbb{R}) \times_M FM \longrightarrow \\
&J((M \times \mathbb{R}) \times_M FM) \subset T^*(M \times \mathbb{R}) \otimes_{FM} T((M \times \mathbb{R}) \times_{FM} FM), \\
(x^\lambda, \alpha, b_\kappa^\eta) &\longmapsto [(X^\lambda, Y_{\eta\kappa}^\lambda) \longrightarrow (X^\lambda, Y_{\eta\kappa}^\lambda, b_\kappa^\eta \alpha X^\theta)].
\end{aligned}$$

briefly,

$$\Lambda = dx^\lambda \otimes \partial_\lambda + d\alpha \otimes \partial_\alpha + b_\kappa^\eta \alpha dx^\theta \otimes \tilde{\partial}_{\kappa\lambda}, \quad \lambda, \eta, \kappa, \theta = (1, \dots, n), \quad \alpha \in \mathbb{R}$$

Explicitly, each  $\tilde{\Gamma} \in \text{Sec}((M \times \mathbb{R})/M)$  gives an injection  $(\tilde{\Gamma} \circ \pi_F, I_{FM})$ , of  $FM$  into  $(M \times \mathbb{R}) \times FM$ , which is a section of  $\pi_1$  and  $\Gamma$  coincides with the restriction of  $\Lambda_F$  to this section.

The connection  $\Lambda$  is universal in the following sense. If  $\tilde{\Gamma} \in \text{Sec}((M \times \mathbb{R})/M)$ , then  $\tilde{\Gamma}$  is a constant real function on  $M$ , so the induced connection  $\Gamma = \xi \circ (\tilde{\Gamma} \circ \pi_F, I_{FM}) : FM \longrightarrow JFM$  coincides with restriction on  $\Lambda$ , on the embedding by  $(\tilde{\Gamma} \circ \pi_F, I_{FM})$  of  $FM$  in  $(M \times \mathbb{R}) \times_M FM$ . So  $\Gamma$  is a pullback of  $\Lambda$ . The universal curvature on the system of  $\alpha$ -connections is

$$\Omega : (M \times \mathbb{R}) \times_M FM \longrightarrow \wedge^2(T^*(M \times \mathbb{R})) \otimes_{FM} V(FM).$$

So  $\Omega$  has the explicit form:

$$\Omega = \frac{1}{2} \left( b_{\kappa}^k \alpha \tilde{\partial}_{\kappa\beta} b_{\omega}^m \alpha dx^{\lambda} \wedge dx^{\eta} + 2 \partial_a b_{\omega}^m \alpha dx^h \wedge dx^{\eta} \right) \otimes \tilde{\partial}_{\omega\alpha},$$

for  $(\kappa, k, \beta, m, \omega, \lambda, \eta, h = 1, \dots, n), \quad \alpha \in \mathbb{R}.$

## References

- [1] S. Amari. **Differential Geometrical Methods in Statistics**, Springer Lecture Notes in Statistics 28, Springer-Verlag, Berlin 1985.
- [2] S. Amari and H. Nagaoka. **Methods of Information Geometry**, American Mathematical Society, Oxford University Press, 2000.
- [3] Khadiga Arwini. **Differential geometry in neighbourhoods of randomness and independence** PhD thesis, Department of Mathematics, UMIST 2004.
- [4] Khadiga Arwini and C.T.J. Dodson. Neighbourhoods of randomness and information geometry of the McKay bivariate gamma 3-manifold. In Proc. **International Mathematica Symposium 2003** Imperial College Press, London 2003 pp. 247-254.
- [5] Khadiga Arwini and C.T.J. Dodson. Information geometric neighbourhoods of randomness and geometry of the McKay bivariate gamma 3-manifold. *Sankhya: Indian Journal of Statistics* 66, 2 (2004) 211-231.
- [6] Khadiga Arwini and C.T.J. Dodson. Neighbourhoods of independence and associated geometry. Preprint 2004. <http://www.ma.umist.ac.uk/kd/PREPRINTS/nhdindep.pdf>
- [7] D.Canarutto and C.T.J.Dodson. On the bundle of principal connections and the stability of b-incompleteness of manifolds. *Math. Proc. Camb. Phil. Soc.* 98, (1985) 51-59.
- [8] L.A. Cordero, C.T.J. Dodson and M. deLeon. **Differential Geometry of Frame Bundles**. Kluwer, Dordrecht, 1989.
- [9] L. Del Riego and C.T.J. Dodson. Sprays, universality and stability. *Math. Proc. Camb. Phil. Soc.* 103(1988), 515-534.
- [10] C.T.J. Dodson. Systems of Connections for Parametric Models. In Proc. Workshop **Geometrization of Statistical Theory**, Lancaster 28-31 October 1987, Ed. C.T.J. Dodson, pp. 153-169, ULDM Publications, Lancaster University 1987.
- [11] C.T.J. Dodson and Hiroshi Matsuzoe. An affine embedding of the gamma manifold. *Appl. Sci.*, 5 (2003) 1-6.
- [12] C.T.J. Dodson and M. Modugno. Connections over connections and universal calculus. In Proc. **VI Convegno Nazionale di Relativita General e Fisic Della Gravitazione** Florence, 10-13 October 1984, Eds. R. Fabbri and M. Modugno, pp. 89-97, Pitagora Editrice, Bologna, 1986.
- [13] R.A. Fisher. Theory of statistical estimation. *Proc. Camb. Phil. Soc.* 122 (1925) 700-725.
- [14] P.L. Garcia. Connections and 1-jet fibre bundles. *Rend.Sem. Mat. Univ.Padova* 47, (1972) 227-242.
- [15] T-Y. Hwang and C-Y. Hu. On a characterization of the gamma distribution: The independence of the sample mean and the sample coefficient of variation. *Annals Inst. Statist. Math.* 51, 4 (1999) 749-753.
- [16] S. Kobayashi and K. Nomizu. **Foundations of differential geometry II**. Interscience New York 1969.
- [17] S. Kumar. A remark on universal connections. *Math. Ann.* 260,4 (1982) 453-462.

- [18] L. Mangiarotti and M. Modugno. Fibred spaces, jet spaces and connections for field theories. In Proc. International Meeting on **Geometry and Physics**, Florence, 12-15 October 1982, ed. M. Modugno, Pitagora Editrice, Bologna, 1983 pp 135-165.
- [19] M. Modugno. Systems of vector valued forms on a fibred manifold and applications to gauge theories. In Proc. Conference **Differential Geometric Methods in Mathematical Physics**, Salamanca 1985, Lecture Notes in Mathematics 1251, Springer-Verlag, Berlin 1987, pp. 238-264.
- [20] M.S. Narasimhan and S. Ramanan. Existence of universal connections I. *Amer. J. Math.* 83, (1961) 563-572.
- [21] M.S. Narasimhan and S. Ramanan. Existence of universal connections II. *Amer. J. Math.* 85, (1963) 223-231.
- [22] C.R. Rao. Information and accuracy attainable in the estimation of statistical parameters. *Bull. Calcutta Math. Soc.* 37, (1945) 81-91.
- [23] S. Roman. **Coding and Information Theory**, Graduate Texts in Mathematics, 134 Springer-Verlag, New York, 1992, cf. Part 1.